



CIAT - COL.
000206.

Working Document
No. 132

Seed Health Testing and
Phytosanitary
Procedures for Tropical
Forages

Compiled by:

Segenet Kelemu
Plant Pathology Section
Tropical Forages Program

Tropical Forages Program
Plant Pathology

November 1993

BIBLIOTECA
CANJE

15952

7 MAR. 1994

14743

CIAT

Centro Internacional de Agricultura Tropical
International Center for Tropical Agriculture

TABLE OF CONTENTS

	Page
I. Introduction.	1
II. Review of literature on seed-borne pathogens of tropical forage plants.	2
Seed-borne or seed associate pathogens of tropical forage legumes.	3
Seed-borne or seed associate pathogens of tropical forage grasses.	6
III. Seed disinfection or decontamination.	10
IV. Seed health testing.	10
i. Methods of detection of seed-borne fungi and bacteria.	10
ii. Selection of tests.	13
1. The blotter or filter paper test.	13
2. The agar plate test.	15
3. Adult plant inspection.	16
iii. Summary on seed health testing.	16
V. General procedures and recommendations.	16
i. General procedures for germplasm introductions.	17
ii. Procedures for adult plant inspections in the glasshouse.	18
References	20
Appendix 1: Definitions	22
Appendix 2: Culture media recipes	23
Appendix 3: Addresses of plant quarantine offices	24

SEED HEALTH TESTING AND PHYTOSANITARY PROCEDURES FOR TROPICAL FORAGES

I. INTRODUCTION

According to Kahn (1989), the word "quarantine" originated from the Latin "quadraginata" and the Italian "quarantina", applied to the 40-day period of isolation required for a ship (passengers and cargo) to remain anchored in port of arrival if the ship arrived from a country where certain epidemic diseases were known to occur. This practice was started to allow the development of symptoms before any passengers debarked.

Plant quarantines regulate the entry of plants and their products, soil, microbial cultures, and commodities in order to protect agriculture and the environment from avoidable pests and pathogens. Organisms of quarantine importance may include pests or pathogens that governments consider threats to the agriculture and environment of countries. These organisms are either foreign to a country or include foreign strains and races of already existing organisms.

In the case of forage legumes and grasses this becomes very complicated by the following factors: 1. there are numerous plant genera and species, 2. there are a number of pathogens which are capable of infecting these wild plants, 3. the lack of appropriate and complete documentation of these pathogens, 4. lack of information and scientific data on the biology, epidemiology, economic importance, complete geographic distribution, etc. of the documented pathogens.

Despite these complications and difficulties, careful tests and treatments should enable us to ensure safe movement of forage germplasm.

Testing, screening, disinfection and seed decontamination are procedures to be considered in phytosanitary programs.

II. REVIEW OF LITERATURE ON SEED-BORNE PATHOGENS OF TROPICAL FORAGE PLANTS

There are many seed-borne pathogens in Gramineae and Leguminosae which are frequently transmitted. Seed transmission of pathogens depends basically on the inherent features of both the host and the pathogen. Members of a plant family transmit groups of related pathogens through seeds; e.g., smut fungi in Graminae, viruses or *Colletotrichum* spp. in Leguminosae.

In the family Gramineae, there is a wide range of seed-borne pathogens such as the smuts, ergots, and seed gall nematodes.

Seed-borne diseases are common in Leguminosae and probably far more severe than in other plant families.

There is very little published information on seed-borne pathogens of tropical forage plants. *Colletotrichum gloeosporioides* and *C. truncatum* were reported to significantly reduce emergence, survival, and root and shoot dry weight of *Stylosanthes hamata* as seed-borne pathogens (Lenné and Sonoda, 1979). *C. gloeosporioides* can also be a seed-borne pathogen of *Centrosema pubescens* and *Pueraria phaseoloides* (unpublished documents of Dr. J. Lenné).

Rhizopus stolonifer, the causal agent of seedling blight and inflorescence diseases of *Stylosanthes* spp., was reported to be the most common fungus associated with seeds of *S. hamata* in Florida (Lenné and Sonoda, 1978). In germination tests, this organism affected seedling emergence and survival.

Seed-borne pathogens of other tropical forage legumes have been reported. *Pseudomonas florescens*, the cause of severe pod rot of *Leucaena leucocephala*, and bacterial leaf spot and dieback of *Centrosema* spp., has been reported to be seed-borne (Lenné *et al.*, 1981a, 1981b). *Pseudomonas syringae* which causes leaf spot disease of *Centrosema pubescens* could be seed-borne (Duveiller, 1987).

Seed-borne or seed associated pathogens of tropical forage legumes.

Host	Pathogen	Country
<i>Arachis</i> **	<i>Cercospora arachidicola</i>	Brazil, Colombia
	<i>Cercospora canescens</i>	Malawi
	<i>Cercospora personata</i>	Brazil, Colombia
	<i>Cochliobolus lunatus</i>	Nigeria
	<i>Colletotrichum capsici</i>	Nigeria
	<i>Colletotrichum gloeosporioides</i>	Brazil, Colombia, Australia
	<i>Colletotrichum truncatum</i>	Colombia
	<i>Fusarium</i> spp.	Colombia
	<i>Leptosphaerulina arachidicola</i>	Brazil, Colombia, Costa Rica, Ecuador, Peru, USA
	<i>Mycosphaerella arachidis</i>	Nigeria
	<i>Myrothecium roridum</i>	India
	Peanut mottle virus	Brazil, Colombia
	<i>Puccinia arachidis</i>	Brazil
	<i>Periconia</i> spp.	Colombia
<i>Rhizoctonia solani</i>	Brazil, Colombia, Costa Rica, Peru	
<i>Sphaceloma arachidis</i>	Brazil, Colombia	
<i>Centrosema</i>	<i>Cercospora canescens</i>	Australia, Barbados, Bolivia, Brazil, Colombia, Costa Rica, Cuba, Ecuador, French Guyana, Malaysia, Panama, Peru, Sudan, USA, Venezuela
	<i>Colletotrichum gloeosporioides</i>	Australia, Barbados, Bolivia, Brazil, Colombia, Costa Rica, Ecuador, Peru, USA, Venezuela
	<i>Colletotrichum truncatum</i>	Colombia, USA
	Mosaic viruses	Australia, Brazil, Colombia, Cuba
	<i>Pseudomonas flourescens</i>	Colombia
	<i>Pseudomonas syringae</i>	Burundi
	<i>Rhizoctonia</i> sp.	Australia, Brazil, Colombia, Costa Rica, Ecuador, French Guyana, Ghana, Indonesia, Malaysia, Papua New Guinea, Peru, Sierra Leone, Seychelles, Solomon Islands, USA, Vanuatu

Host	Pathogen	Country
<i>Desmodium</i> sp.	<i>Cercospora</i> sp.	Australia, Barbados, Brazil, Canada, Colombia, Dominican Republic, El Salvador, Ghana, India, Jamaica, Kenya, Malaysia, Pakistan, Papua New Guinea, Tanzania, USA, Venezuela, Zimbabwe
	<i>Cladosporium cladosporioides</i>	Australia
	<i>Cladosporium</i> sp.	India, Cuba
	<i>Colletotrichum gloeosporioides</i>	Australia, Bolivia, Brazil, Colombia, Ecuador, Peru, USA, Venezuela
	<i>Colletotrichum truncatum</i>	Colombia, USA
	Desmodium mosaic virus	USA
	<i>Rhizoctonia</i> sp.	Brazil, Colombia, Costa Rica, El Salvador, Malaysia, Peru
	<i>Synchytrium desmodii</i>	Colombia, Ecuador, Tanzania
<i>Pueraria</i> sp.	<i>Cercospora</i> sp.	Cambodia, Cuba, Malaysia, Philippines, Puerto Rico, Sierra Leone, Taiwan, Tanzania
	<i>Cladosporium</i> sp.	Sierra Leone
	<i>Colletotrichum gloeosporioides</i>	Brazil, Colombia, Ecuador, Peru, Puerto Rico, Venezuela, Virgin Islands
	<i>Drechslera</i> sp.	Malaysia
	<i>Fusarium</i> sp.	Papua New Guinea, Sierra Leone
	<i>Mycosphaerella puerariicola</i>	Cambodia, Hong Kong, Japan, Malaysia, Philippines, Singapore, Solomon Islands, Taiwan, USA
	<i>Pseudomonas phaseolicola</i>	USA
	<i>Rhizoctonia</i> sp.	Brazil, Colombia, Costa Rica, Ecuador, Malaysia, Peru, Sierra Leone, Solomon Islands
<i>Stylosanthes</i>	<i>Ascochyta</i> spp.	Colombia
	<i>Bipolaris</i> spp.	Colombia
	<i>Botrytis cinerea</i>	Australia, Colombia, Zimbabwe
	<i>Cladosporium</i> spp.	Australia, Brazil

Host	Pathogen	Country
<i>Stylosanthes</i>	<i>Colletotrichum gloeosporioides</i>	Angola, Australia, Belize, Bolivia, Botswana, Brazil, Colombia, Costa Rica, Cuba, Ecuador, Ethiopia, French Guyana, India, Ivory Coast, Kenya, Malaysia, Malawi, Mexico, Mozambique, Nigeria, Panama, Papua New Guinea, Peru, Solomon Islands, Tanzania, Thailand, Venezuela, Zaire, Zambia, Zimbabwe, USA
	<i>Curvularia</i> spp.	Australia, Colombia, USA
	<i>C. truncatum</i>	Australia, Bolivia, Brazil, Colombia, Ecuador, Ethiopia, Nigeria, Peru, Thailand, USA, Venezuela
	<i>Eurotium</i> spp.	Colombia
	<i>Fusarium</i> spp.	Australia*, Brazil, Colombia, Malaysia, Nigeria
	<i>Gloeocercospora</i> spp.	Colombia
	<i>Leptosphaerulina</i> spp.	Colombia, Australia
	<i>Phoma sorghina</i>	Australia, Colombia, Ivory Coast, Nigeria
	<i>Phomopsis</i> spp.	Colombia, Bolivia, Nigeria, Malaysia, Trinidad,
	Potyvirus	Colombia
	<i>Rhizopus stolonifer</i>	Colombia, India, USA
	<i>Rhizoctonia</i> spp.	Australia, Colombia, Malaysia, Papua New Guinea
	<i>Sclerotinia sclerotiorum</i>	India, Zimbabwe
<i>Zornia</i> sp.	<i>Colletotrichum gloeosporioides</i>	Brazil, Colombia, Costa Rica, Ecuador, Panama, Peru, USA
	<i>Colletotrichum truncatum</i>	Bolivia, Brazil, Colombia, Ecuador, Peru, USA, Venezuela,
	<i>Corynebacterium flaccumfaciens</i>	Colombia
	<i>Fusarium</i> sp.	Colombia
	<i>Phoma sorghina</i>	Colombia
	<i>Rhizoctonia solani</i>	Brazil, Colombia, Peru
	<i>Sphaceloma zorniae</i>	Brazil, Colombia, Peru, Venezuela,

* Several species of *Fusarium* reported.

** The pathogens are not necessarily seed-borne or seed associated. These can be transported through planting materials such as cuttings.

Source (fungal pathogens): Lenné, J.M. 1990. A world list of fungal diseases of tropical pasture species. CAB International, Wallingford, U.K.

Seed-borne or seed-associated pathogens of tropical forage grasses.

Host	Pathogen	Country
<i>Andropogon</i>	<i>Balansia</i>	Bengal, China, India, Mexico, Nigeria, Philippines, Puerto Rico, Sierra Leone, West Indies, Zambia,
	<i>Cerebella andropogonis</i>	Burma, Cuba, Ghana, Jamaica, Romania, Sierra Leone, Sri Lanka, Uganda, Zambia, Zimbabwe
	<i>Claviceps</i> , ergot	Bengal, Colombia Kenya, USA,
	<i>Curvularia</i> sp.	Colombia, Nigeria, Sudan
	<i>Phoma sorghina</i>	Colombia, Sri Lanka
	<i>Sorosporium</i>	Argentina, Cambodia, Colombia, Congo, India, Israel, Morocco, Sierra Leone, South Africa, Sri Lanka, USA
	<i>Sphacelia sorghi</i> , ergot	Ethiopia, Ghana
	<i>Sphacelotheca</i> spp.	Bolivia, Brazil, China, Gambia, India, Malawi, Mexico, Pakistan, Sierra Leone, Sudan, Tanzania, Uganda, USA, Venezuela, Zambia, Zimbabwe
	<i>Tolyposporella</i> , smut	Brazil, Dominican Republic, Mexico, Puerto Rico
	<i>Tolyposporium</i> , smut	Nigeria, India
	<i>Tolysporum</i> , smut	India
<i>Ustilago</i> , smut	Brazil, Congo, Ghana, India, Malawi, Mexico, Nigeria, Sierra Leone, Venezuela, West Africa	
<i>Brachiaria</i> sp.	<i>Alternaria alternata</i>	Kenya, Pakistan
	<i>Balansia</i> sp., false smut	Brazil, Fiji, India, Zambia
	<i>Botryodiplodia</i>	Colombia
	<i>Ceratocystis paradoxa</i>	Colombia
	<i>Cercospora</i> sp.	Australia, Bengal, Botswana, Ethiopia, French Guyana, Kenya, Mali, Tanzania, Uganda, Zambia, Zimbabwe
	<i>Cerebella andropogonis</i> , false smut	Bengal, Burma, Kenya, South Africa, Sudan, Tanzania, Uganda, USA, Zambia, Zimbabwe

Host	Pathogen	Country
<i>Brachiaria</i> sp.	<i>Cladosporium</i> sp.	Brazil, Colombia, Ethiopia, Pakistan, Uganda
	<i>Claviceps</i> sp., ergot	Australia, Bengal, Ethiopia, India, Kenya, Malawi, Tanzania, Zambia, Zimbabwe
	<i>Curvularia</i>	Colombia, Fiji
	<i>Ephelis</i> sp.	Australia
	<i>Fusarium acuminatum</i>	Colombia
	<i>Fusarium heterosporum</i>	Ethiopia, Kenya, Tanzania, Uganda, Zambia, Zimbabwe,
	<i>Fusarium</i> sp.*	Brazil, Colombia, Ecuador, French Guyana, Peru, Zambia
	Guineagrass mosaic virus***	Africa, Brazil, Colombia, Peru
	<i>Melanotaenium brachiariae</i> , smut	India
	<i>Phoma sorghina</i>	Colombia
	<i>Pyricularia oryzae</i>	Colombia, Fiji, India, Nepal, Malaysia, Philippines, Thailand
	<i>Rhizoctonia solani</i>	Brazil, Colombia, Costa Rica, Ecuador, Ethiopia, Guadalcanal, India, Malaysia, Peru, Solomon Islands
	<i>Rhizopus stolonifera</i>	Colombia
	<i>Sorosporium brachiariae</i> , smut	Zambia, Zimbabwe
	<i>Sorosporium cryptum</i> , smut	Australia
	<i>Sorosporium</i> sp., smut	Malawi, USA
	<i>Sphacelia</i> sp., ergot	Burma, Sudan, Uganda, Zambia
	<i>Sphacelotheca serrata</i> , smut	Malawi
	<i>Ustilago operta</i> , smut	India, Pakistan, Sudan
	<i>Ustilaginoidea virens</i> , false smut	Kenya, Malawi, Peru, Sudan, Tanzania, Zambia, Zimbabwe,
<i>Xanthomonas</i> sp.**	Colombia, French Guyana	
<i>Panicum</i>	<i>Balansia</i>	Brazil, Costa Rica, Guinea, India, Jamaica, Malaysia, Nicaragua, Panama, Sierra Leone, Trinidad, Venezuela
	<i>Bipolaris</i>	Australia, China, India, South Africa, USA, Venezuela

Host	Pathogen	Country
<i>Panicum</i>	<i>Cercospora</i>	Australia, Bolivia, Brazil, Colombia, Costa Rica, Cuba, Ecuador, El Salvador, Ethiopia, Fiji, French Guyana, Ghana, Guadalupe, Guatemala, India, Ivory Coast, Jamaica, Kenya, Malaysia, Mexico, Nicaragua, Nigeria, Panama, Papua New Guinea, Peru, Puerto Rico, Sierra Leone, Sudan, Tanzania, Thailand, Togo, Trinidad, Uganda, USA, Vanuatu, Venezuela, Virgin Islands, Zambia, Zimbabwe,
	<i>Cerebella</i>	Brazil, Dominican Republic, Ghana, India, Jamaica, Nigeria, Malawi, Puerto Rico, Sierra Leone, South Africa, Sudan, Tanzania, Venezuela, Virgin Islands, Zambia, Zimbabwe,
	<i>Cochliobolus</i>	Australia, Colombia, Cuba, Egypt, French Guyana, India, Japan, Jamaica, Nigeria, Papua New Guinea, Peru, Sierra Leone, Solomon Islands, USA, Zambia, Zimbabwe
	<i>Cladosporium</i>	Cyprus, Bolivia, Sierra Leone, Sudan, Venezuela, Zambia
	<i>Claviceps</i>	Australia, Bengal, Brazil, Colombia, Costa Rica, French Guyana, India, Kenya, Mauritius, Peru, Puerto Rico, Sierra Leone, Sudan, Tanzania, Venezuela, Virgin Islands, Zambia, Zimbabwe
	<i>Fusarium</i>	Bengal, Brazil, Cameroon, Colombia, Costa Rica, Ghana, Kenya, Malawi, Peru, Puerto Rico, Tanzania, Venezuela, Zambia, Zimbabwe
	<i>Helminthosporium</i>	Australia, Guadalupe, India, Malaysia
	<i>Magnaporthe grisea</i>	Australia, Dominican Republic, Egypt, India, Nigeria, Malaysia, Puerto Rico, Tanzania, USA, West Africa, Zambia, Zimbabwe,
	<i>Phoma sorghina</i>	Bolivia, Brazil, China, Ethiopia, India, Ivory Coast, Nigeria,

Host	Pathogen	Country
<i>Panicum</i>	<i>Sorosporium</i>	Angola, Australia, Bengal, Botswana, Congo, Cyprus, China, Ghana, India, Israel, Malawi, Northeast Africa, South Africa, Sri Lanka, Sudan, Tanzania, USA, Zimbabwe,
	<i>Sphacelia</i> sp.	Australia, Bengal, Brazil, Colombia, Costa Rica, Jamaica, Kenya, Malawi, Peru, Venezuela, Zambia
	<i>Sphacelotheca</i>	Australia, Bulgaria, Canada, Czechoslovakia, India, Poland, Yemen, Yugoslavia, Romania, USA, USSR
	<i>Tilletia</i>	Belize, Bolivia, Brazil, Canada, Cameroon, Colombia, Congo, Costa Rica, Cuba, Ecuador, Ghana, India, Ivory Coast, Kenya, Malawi, Mauritius, Mexico, Natal, Nigeria, Pakistan, Panama, Philippines, South Africa, Sri Lanka, Sudan, Tanzania, Uganda, Venezuela, Zambia, Zimbabwe
	<i>Ustilaginoidea</i>	Burma, Congo, Malaysia, Philippines, Puerto Rico, Tanzania, Trinidad & Tobago, Sierra Leone
	<i>Ustilago</i>	Argentina, Brazil, Bulgaria, Canada, Czechoslovakia, England, Guatemala, India, Italy, Kenya, Mauritius, Mozambique, Nigeria, Romania, Uganda, Yugoslavia, Sri Lanka, Sudan, Thailand, Trinidad, USA, Zimbabwe,

* Several species of *Fusarium* reported.

** Potential to be seed-borne.

*** Not seed-borne, but can be introduced through tissue culture or vegetative materials.

Source (fungal pathogens): Lenné, J.M. 1990. A world list of fungal diseases of tropical pasture species. CAB International, Wallingford, U.K.

III. SEED DISINFECTION OR DECONTAMINATION

Seed treatment for eradication of seed-borne pathogens can be used as a precaution in quarantine only if careful consideration is given to the limitations of such procedures. In quarantine, no residual inoculum must remain after treatment. The primary justification for seed treatment as a quarantine precaution is its use as an additional safeguard to kill undetected trace amounts of inoculum in apparently healthy seed lots. **Another acceptable possibility would be to treat seed which carry low amounts of inoculum, in order to save particularly valuable germplasm material.**

A number of countries require that certain seeds and / or propagative materials be treated with broad spectrum fungicides or specific fungicides as a condition of entry. Some of the fungicides most cited are thiram, benomyl, captan and vitavax.

IV. SEED HEALTH TESTING

Seed health testing is conducted to provide information for the following purposes:

1. Seed inspection for quarantine purposes
2. Seed quality evaluation for planting value
3. Seed certification purposes

i. Methods of detection of seed-borne fungi and bacteria

There are several methods of seed testing. Usually there are a number of microorganisms carried in a seed lot, and each of these may have requirements for its growth. The method to be employed will depend mostly on the purpose of the test, the particular pathogen, and the type of seed.

Seed health testing procedures (adopted from Neergaard, 1977)

METHOD	APPLICATION
<p>Direct inspection</p> <p>Examination of the dry seed, with impurities, using a hand lens or, a stereo-microscope. Seeds may be submerged in water to release spores and facilitate detection.</p>	<p>Sclerotia of fungi, smut balls, nematode galls, infected plant debris; e.g., <i>Sclerotinia sclerotiorum</i>, <i>Botrytis cinerea</i>, <i>Claviceps purpurea</i>. Seeds discolored or with lesions produced by fungi, bacteria, or viruses, e.g., anthracnose fungi, some Xanthomonads, viruses such as soybean mosaic virus in leguminous seed.</p>
<p>Examination of suspension from washings of seed</p> <p>An electrical mechanical shaker can be used to obtain standardized washings. Samples of the suspension are examined under compound microscope.</p>	<p>Covered smuts, e.g., <i>Tilletia</i> spp. in monocots, oospores of certain downy mildews; quick orientation on the presence of other fungi that must be adequately detected by incubation procedures, e.g., <i>Drechslera</i> spp.</p>
<p>Whole-embryo count method</p> <p>Soaking grains overnight in 10% NaOH at 22 C, then washing with warm water through sieves of decreasing mesh size. Embryos finally cleared in lactophenol.</p>	<p>Loose smuts of monocots</p>
<p>Blotter method</p> <p>Seeds are incubated on water-moistened blotter, usually for 7 days at 20 C. Sporulation of fungi is stimulated by near-ultra-violet (NUV) irradiation, standard 12/12-hr light/dark cycle. Petri dishes are usually used as containers. To allow penetration of the NUV, plastic or pyrex glass containers should be used. Sometimes blotters are soaked in 0.1-0.2% 2,4-D solution to counteract seed germination, thus aiding recording.</p>	<p>The method is used most commonly for detecting a wide-range of Fungi Imperfecti, including different spp. of <i>Acremonium</i>, <i>Acroconidiella</i>, <i>Alternaria</i>, <i>Ascochyta</i>, <i>Botrytis</i>, <i>Cercospora</i>, <i>Colletotrichum</i>, <i>Diplodia</i>, <i>Drechslera</i>, <i>Fusarium</i>, <i>Macrophomina</i>, <i>Myrothecium</i>, <i>Phoma</i>, <i>Phomopsis</i>, <i>Septoria</i>, and others, and for practically all kinds of seed, including cereals, grasses, ornamentals, forest seeds, and vegetables.</p>
<p>Agar plate method</p> <p>The seeds are plated in Petri dishes on nutrient agar, in particular malt extract agar, potato-dextrose agar. Some selective media are available for specific tests. Light treatment as for the blotter test. Incubation for 5-7 days.</p>	<p>Although slowly growing fungi cannot be adequately detected, the procedure is relatively sensitive for revealing minor amounts of inoculum.</p>

METHOD	APPLICATION
<p>Freezing method</p> <p>A modified blotter method. After 1-2 days at 10-20 C, according to specifications, incubation for some hours or for 1 day at -20 C, then at 20 C in NUV light for 5-7 days.</p>	<p>Sometimes preferred for detecting certain fungi, e.g., <i>Phoma</i>, <i>Septoria</i>, <i>Alternaria</i>.</p>
<p>Ordinary seedling symptom test</p> <p>The seed is sown in autoclaved soil, sand, or similar material and placed under normal daylight conditions for observation of symptoms.</p>	<p>Often used for detecting seedling symptoms which reveal pathogens rather than fungi to be identified. Useful for detecting seedling pathogens.</p>
<p>Water-agar seedling symptom test</p> <p>The seeds are sown on water agar, in 16-mm test tubes, one seed per tube, or in microculture plastic plates or Petri dishes. They are placed under daylight conditions, e.g., 12/12-hr cycle of artificial daylight and darkness. Seedlings are inspected for symptoms, healthy seedlings may be transplanted for further post-entry quarantine cultivation.</p>	<p>Can be used for many kinds of seed as an economical procedure which, in test tubes, secures separation of healthy and infected seedlings. Has been used for detecting <i>Drechslera</i>, <i>Septoria</i>, <i>Macrophomina</i>; and other pathogens and hosts.</p>
<p>Indicator test, inoculation methods</p> <p>A standard technique for identification of viruses but also used for detection of trace amounts of pathogenic bacteria, e.g., by hypodermic injection of indicator plants with material from seeds slightly infected by the pathogen under test.</p>	<p>Used for detection of <i>Xanthomonas</i> in beans, crucifers, etc.</p>
<p>Phage-plaque method</p> <p>Maceration of the seed to be tested followed by incubation by for 24 hr to enable multiplication of bacteria. Samples of this material transferred to sterile flasks, and a standard suspension of phage particles is added. Samples of this mixture are plated immediately and after 6-12 hr on plates with the indicator bacterium. Presence of homologous bacteria is indicated by significant increase in number of phage particles in the second plating.</p>	<p>Used for <i>Pseudomonas phaseolicola</i> and <i>Xanthomonas phaseoli</i> in French bean, <i>Xanthomonas vesicatoria</i> in tomato.</p>

METHOD	APPLICATION
<p>Serological methods</p> <p>An antiserum must be provided and the tests may follow different procedures: the slide agglutination test, the tube precipitin test, the micro-precipitin test, the gel double- diffusion test, the latex flocculation test, the immunofluorescence test.</p>	<p>Used for different seed-borne viruses, and may be used for any pathogen for which an antiserum is provided.</p>

ii. Selection of tests

The test to be used should be capable of revealing the widest possible range of pathogens. The procedures can be combined in different ways. For instance, healthy-looking seedlings can be selected from the blotter and/or agar plate tests and transplanted into soil, sand, or another suitable medium for further growth and observation. The combination of procedures saves time and seed.

In addition to generalized procedures, it may be necessary to include specialized tests to detect specific pathogens: different serological procedures to reveal seed-borne viruses and bacteria, the phage-plaque test for detecting seed-borne bacteria, and the indicator test for detecting both of these categories of pathogens as well as fungi.

Procedural details:

1. The Blotter or Filter Paper Test

The method is widely used to detect a number of fungi. The procedural details of this method are as follows (adopted from International Seed Testing Association):

- a. A working sample of 400 seeds is tested in replicates of 25 seeds per dish of 9 cm diam. Since the

seeds are later incubated in light, the dishes should be of such a material that allows light to pass through, e.g. Petri dishes made from clear plastic, and glass dishes of Pyrex or Corning. In plant quarantine, the samples of seeds may be less than 400 . In such cases the whole sample or part of it may be tested. Only healthy seedlings, free from infection, must be released and grown in the quarantine glasshouse for inspection by the quarantine officer.

- b. Label each dish properly with the accession number of the seed sample, date of examination and the dish number.
- c. Before plating the seeds in the Petri dishes, each dish should be lined with three filter papers (blotters), well soaked in water. The water soaked filter papers for the dishes can be prepared in the following way: Count three filter papers at a time, dip them in water for a few seconds, lift and let the excess water drip off before setting the wet blotters in the dish.
- d. Plate 25 seeds in each dish, 15 seeds in the outer ring, 9 in the middle and one in the centre.
- e. Incubate the seeds at 20-22 C for 7 days in alternating cycles of 12 hr light and 12 hr darkness. Light should be supplied by two fluorescent tubes hanging horizontally, 20 cm apart from each other and the distance between the light tubes and the dishes should be 40 cm.
- f. After 7 days incubation, start examination of seeds under a stereoscopic microscope with magnification at least up to x 50 or x 60. All seeds of the outer ring must be examined first, then the seeds of the second ring and finally the seed in the centre of the dish. Examination of seeds in sequence becomes easier when a line is drawn with a colored pencil. Examine thoroughly the whole seed at different magnifications before proceeding to the next seed.

Once the examination of the first seed is finished rotate the dish gently clockwise with the middle finger and the thumb of the left hand while still looking into the microscope. Follow this procedure in moving from one seed to the other.

- g.** Whenever the growth of an organism of interest is seen, mark the seed infected by writing near to the infected seed.
- h.** Identification of a particular organism needs experience in seed health testing. Whenever a growth is suspected of a particular organism, mount conidia (spores) in water on a glass slide and confirm the conidial morphology of a particular fungus under higher magnifications of a compound microscope.
- i.** Count the total number of seeds infected by a particular organism in each dish and enter the figures in a recording sheet, and calculate per cent seed infection.

2. The Agar Plate Test

Seeds are plated on an appropriate agar medium and spaced according to the size of the seed. Malt extract agar, oatmeal agar, or potato dextrose agar are most commonly used (see media preparations in Appendix (2)).

Pretreatment of the seeds (soaking for 5-10 minutes in 1 per cent sodium hypochlorite) is applied to exclude fast growing saprophytes associated with the seeds. The following procedure is a more rigorous seed pretreatment. **a.** Put seeds in a screw capped tube containing about 25 ml of 30% (v/v) bleach (ie., 5% hypochlorite) 1 ul/ml of 20% Triton-X100, **b.** Shake vigorously so that a layer of foam develops on the solution. Most of the trash (which contains most of the contaminants) will collect in the foam which should be removed by aspiration with a sterile pasteur pipet, **c.** Continue to mix slowly by inversion for 5 to 10 minutes, **d.** Allow the seeds to settle for 1 or 2 minutes and decant the bleach, **e.** Fill the tube with sterile water, let seeds settle and decant, **f.** Repeat this process several times to remove as much bleach as possible. The seeds should be plated within a day. The incubation conditions are the same as described

under the blotter test.

3. Adult Plant Inspection

Certain seed-borne diseases require a longer period of incubation to be detected than is provided by the usual incubation procedure. The seeds are sown in autoclaved soil in suitable containers and submitted to optimum conditions of temperature and humidity protected against contamination.

iii. Summary on seed health testing

Detection of fungi: procedures using the blotter or filter paper method and agar media are usually sufficient. The agar media most commonly used are potato dextrose agar (PDA), malt extract agar, and oat meal agar.

Bacteria: procedures using appropriate culture media are adequate, but specific procedures such as phage-plaque or serological tests (provided that antisera are available) may be used. Nutrient agar and King's medium B are widely used.

Viruses: indicator tests (provided that suitable indicator plants are available), and ELISA can be used.

V. GENERAL PROCEDURES AND RECOMMENDATIONS

All agricultural scientists should understand that plant quarantine is a measure of plant disease control which we all want to achieve. It is our responsibility to take all the necessary precautions to prevent man-assisted pathogen disseminations.

i. General procedures for germplasm introductions.

The following procedures are representative of those followed by most quarantine officers in regulating germplasm of high risk genera.

1. Seeds should be collected from "disease-free" plants.
2. Small, discolored, shrivelled or damaged seeds should be avoided.
3. Seeds should be free of any plant residues such as leaf pieces, roots, glumes, husks or other foreign materials such as stones, soil particles, etc.
4. Seeds should not be packed using plant materials such as straws. All packing materials should be clean.
5. Seeds should be introduced, rather than vegetative materials, unless clonal propagation is a requirement or the plants are collected in the wild when seed is not available.
6. For clonal propagations, unrooted vegetative materials such as scions or unrooted cuttings are preferred to rooted plants.
7. Woody plant introductions should not be more than 2 years old.
8. Consignments of vegetatively propagated clonal germplasm should be small, i.e., limited to a few tubers, cuttings, or scions. If an accession is represented by tubers, cuttings, and scions, each

component should be regarded as separate subclone, particularly for virus indexing.

9. Germplasm that must be introduced as plants should enter free of soil. The original importation should be destroyed once daughter plants have been established.
10. If phytosanitary procedures require that clones, or subclones, be indexed for viruses, only propagations derived from indexed mother plants should be released.
11. With genera that present a high risk of seed-borne organisms, the original seed should be treated and planted to produce a healthy lot of seed.
12. For high-risk genera, whether imported as seed or vegetative material, only the part of the shipment that is passed through quarantine should be released. The part of the original introduction not used as mother or seed plants should be destroyed.

ii. Procedures for adult plant inspections in the glasshouse.

In exchanging genetic resources, the quantities of seed for distribution, and hence for sampling and testing, is very limited.

It seems more appropriate and realistic if all original seed introductions are examined at the Seed Health Lab. (currently under GRU) first, using appropriate seed health testing protocols, and those seeds which show no pathogens be planted in the plant quarantine green house for inspection. Visual plant inspection alone may not be effective and may provide only a false feeling of security.

The following steps should be taken in the phytosanitary glasshouse:

1. Pots, soil, and benches should be sterilized before being placed in the glasshouse.
2. The floor of the entrance cubicle should be treated with a disinfectant.
3. All planting of imported material should be carried out only in the glasshouse.
4. Personnel working in the glasshouse should be careful not to brush against plants or handle any plant unless necessary.
5. Hands and instruments should be washed with soap and water or some disinfectant after trimming, digging, etc., before being moved from one place to another.
6. Distance between potted plants should be sufficient to prevent contact with each other.
7. Airflow within the glasshouse unit should be controlled so that it does not cause the plants to touch each other.
8. All materials have to be inspected on individual plant basis.

REFERENCES

- Agarawal, V. K., Sinclair, J. B. 1987. Principles of seed pathology. Boca Raton, Fla., CRC Press.
- Agarawal *et al.* Seed-borne diseases and seed health testing of rice. Hellerup, Denmark, Danish Government Institute of Seed Pathology for Developing Countries. Surrey, U. K. CAB International Mycological Institute (Technical Bulletin No. 3).
- Charles, R. W., and Hance, R. J. 1991. The pesticide manual, 9th edition. The British Crop Protection Council, U. K.
- Chidambaram *et al.* 1974. Identification of seed-borne *Drechslera* species. Norway, The international seed testing association [Handbook on seed health testing, series 2B (3)].
- Duveiller, E. 1987. A bacterial leaf spot disease on *Centrosema pubescens* caused by *Pseudomonas syringae*. FAO Plant Protection Bulletin. 35: (4) 163.
- FAO digest of plant quarantine regulations. 1990. Rome, Food and Agricultural Organization of the United Nations.
- Frison, *et al.* 1990. FAO/IBPGR technical guidelines for the safe movement of legume germplasm. Research Institute for Plant Protection, Wageningen, the Netherlands.
- Hewitt, W. B., Chiarappa, L. 1977. Plant health and quarantine in international transfer of genetic resources. Cleveland, Ohio, CVC Press.
- Joshi, N. C., Ravi, U. 1989. A guide for the import and export of seeds of ICRISAT. Patancheru, Andhra Pradesh, India, International Crops Research Institute for the Semi-Arid Tropics.
- Kahn, R. P. 1989. Plant protection and quarantine. Boca Raton, Fla., CRC Press.
- Klag, N. 1991. Fungicide benefits assessment regulatory guide. National agricultural pesticide impact assessment program.
- Lenné, J. M. and Sonoda, R. M. 1978. *Colletotrichum* spp. on tropical forage legumes. Plant Disease Reporter 62: 813-817.
- Lenné, J. M. 1979. Pathogenicity of *Sclerotium rolfsii* to *Stylosanthes capitata* and other tropical forage legumes. Plant Disease Reporter 63: 739-741.
- Lenné, J. M. *et al.* 1981b. Bacterial leaf spot and dieback of *Centrosema* spp. Proceedings 5th International Conference on Plant Pathogenic Bacteria, CIAT, Cali, Colombia.
- Lenné, J. M. 1990. A world list of fungal diseases of tropical pasture species. CAB International, Wallingford, UK.
- Lenné, J. M. *et al.* 1981a. Bacterial pod rot of *Leucaena leucocephala* caused by *Pseudomonas fluorescens* Biotype 2. Proceedings 5th. International Conference on Plant Pathogenic Bacteria, CIAT, Cali, Colombia.

- Malone, J. P., Muskett, A. E. 1964. Seed-borne fungi: description of 77 fungus species. Wageningen: International seed testing association [Handbook on seed health testing, series 4 (1)].
- Neergaard, P. 1970. Seed Pathology, international cooperation and organization. Vollebakk, Norway, International seed testing association [Handbook on seed health testing, series 1 (5)].
- Singh, K. G. 1983. Exotic plant quarantine pests and procedures for introduction of plant materials. Serdang, Malaysia, Asian Plant Quarantine Centre and Training Institute.
- Sletten, A. 1989. Detection methods for *Xanthomonas campestris* pv. *graminis* on forage grasses. Bulletin-OEPP 19: (1) 69-72.

Appendix 1 : DEFINITIONS

- 1. Seed health:** According to the definition of the International Seed Testing Association (ISTA, 1985), health of seed refers primarily to the presence or absence of disease-causing organisms such as fungi, bacteria and viruses, and animal pests.
- 2. Plant quarantine:** It is a preventive measure against the introduction of pests and pathogens harmful to agriculture. It is basically the use of exclusion as a control strategy and is applied to both imported and exported materials.
- 3. Incubation:** Seed maintenance in a condition favorable to pathogen growth or disease symptom expression.

Appendix 2 : CULTURE MEDIA RECIPES

Potato dextrose agar

Potato dextrose agar (Difco 0013)	39.0 g
Distilled water	1000 ml
or	
Potatoes, infusion from	200.0 g
Bacto-Dextrose	20.0 g
Bacto-agar	15.0 g
Distilled water to	1000 ml

Malt extract agar

Maltose	12.75 g
Dextrin	2.75 g
Glycerol	2.35 g
Bacto-peptone	0.78 g
Bacto-agar	15.0 g
Distilled water	1000 ml
or	
Malt extract	20.0 g
Peptone	5.0 g
Agar	15.0 g
Distilled water	1000 ml
or	
Malt extract agar (Difco 0112)	33.6 g
Distilled water	1000 ml

Oatmeal agar

Agar	5.0 g
Distilled water	500.0 ml

Melt.

Instant oatmeal (for babies)	40.0 g
Distilled water (cold)	250.0 ml

Mix. Combine mixed oatmeal with melted agar

Nutrient agar

Peptone	5.0 g
Beef extract	3.0 g
Agar	15.0 g
Distilled water	1000 ml

King's medium B

Proteose peptone	20.0 g
Glycerol	15.0 g
K ₂ HPO ₄	1.5 g
M ₈ SO ₄ ·17H ₂ O	1.5 g
Agar	15.0 g
Distilled water	1000 ml

Adjust pH to 7.2 before autoclaving the medium

Notes: Media should be sterilized the same day that they are prepared. Sterilization is completed in 20-25 min. at a temperature of 121 C.

Appendix 3: ADDRESSES OF PLANT QUARANTINE OFFICES¹

Afghanistan

Director
Plant Protection and Quarantine
Ministry of Agriculture
Kabul

Albania

Bureau of Agriculture
Tirana

Algeria

Chef
Service de la Protection des Végétaux
Ministère de l'Agriculture
12, Boulevard Colonel
Amirouche
Algiers

American Samoa

Director
Agricultural Quarantine Services
Department of Agriculture
Pago Pago, 96799

Antigua

Director
Ministry of Agriculture
Lands and Fisheries
St. John's
West Indies

Argentina

Director General
Servicio Nacional de Sanidad Vegetal
Ministerio de Agricultura
Paseo Colon 922, 1o Piso
Oficina No 196
Buenos Aires

Australia

Assistant Director General
Plant Quarantine
Department of Health
Canberra A.C.T.

Austria

Director
Bundesanstalt für Pflanzenschutz
Trunnerstrasse 5
1021 Wien

Bahamas

Director of Agriculture
Ministry of Agriculture and Fisheries
P.O. Box 28
Nassau

Bahrain

Department of Agriculture
Plant Protection Section
P.O. Box 251
Manama

Bangladesh

Department of Agriculture
(Extension and Management)
Director of Agriculture (E&M)
Ansari Building
14/2 Topkhana Road, Dhaka-2

Barbados

Head
Division of Entomology and Quarantine
Ministry of Agriculture,
Science and Technology
P.O. Box 505
Bridgetown

Belgium

Ingenieur en Chef-Directeur
Service de la Protection des Végétaux
Ministère de l'Agriculture
36 Rue de Stassart
1050 Bruxelles

Belize

Director of Agriculture
Belmopan

Benin

Service du Développement
Région Agricole du Sud
B.P. 648
Cotonou

Bermuda

Plant Pathologist
Department of Agriculture
and Fisheries
P.O. Box 834
Hamilton 5

¹

Addresses taken from Plant Production and Protection Division publications of the FAO (1981) and from Export Summaries published by the USDA, Animal and Plant Health Inspection Service, Plant Protection and Quarantine Programs.

Bolivia

Jefe
Departamento de Sanidad Vegetal
Ministerio de Asuntos
Campesinos y Agropecuarios
La Paz

Botswana

Entomologist
Department of Agricultura Research
Private Bag 0033
Gaborone

Brazil

Director
Secretaria de Defensa Sanitaria Vegetal
Ministerio da Agricultura
Esplanada dos Ministerios
Bloco 8, 70.000
Brasilia - D.F.

Brunei

The Director of Agriculture
Department of Agriculture
Bandar Seri Begawan

Bulgaria

Director of Plant Protection
and Fertilizers
BD. Christo Botev 55
Sofia

Burkina Faso

Le Directeur
Section de Lutte Antiacridienne
de Protection des Plantes et des Cultures
Direction des Services Agricoles
B.P. 7082
Ouagadougou

Burma

The Director
Department of Agriculture
Ministry of Agriculture and Forests
Rangoon

Central Agricultural Research Institute
Gyogon, Insein
Rangoon

Burundi

Chef
Institut des Sciences Agronomiques
du Burundi (ISABU)
Groupe de Phytopathologie
et d'Entomologie
B.P. 795
Bujumbura

Cameroon

Chef
Service de la Protection des Végétaux
Ministère de l'Agriculture
Yaound

Canada

The Director
Plant Quarantine Division
Production and Marketing Branch
Agriculture Canada
Sir John Carling Bldg, C.E.F.
Ottawa, Ontario K1A 0C5

Cape Verde Islands

Serviço Nacional de Protecção
des Vegetais
Direcção Nacional de Agricultura
Caixa Postal 50
Praia

Cayman Islands

The Director
Department of Agriculture
Grand Cayman

Central Afrinca Republic

Le Directeur
Direction de l'Agriculture
Service de la Defense des Cultures
162 Bangui

Chad

Service National de la
Protection des Végétaux
B.P. 441
N'Djamena

Chile

Director
Division de Protección Agrícola
Servicio Agrícola y Ganadero (SAG)
Ministerio de Agricultura
Casilla 4088
Santiago

China, People's Republic of

Central China Agricultura Research
and Training Institute
Wuhan

Colombia

Director
División de Sanidad Vegetal
Instituto Colombiano Agropecuario (ICA)
Apartado Aéreo 79480
Calle 37 No. 8-43 - PS 8
Santa Fe de Bogotá

Cook Islands

Director of Agriculture
Ministry of Agriculture and Fisheries
Box 96
Rarotonga

Costa Rica

Jefe
Departamento de Cuarentena y Registro
Ministerio de Agricultura y Ganadería
San José

Côte d'Ivoire

Chef
Service de la Protection des Végétaux
Ministère de l'Agriculture
B.P. V7
Abidjan

Cuba

Director
Dirección Nacional de Sanidad Vegetal
INRA
Departamento de Cuarentena Vegetal
La Habana

Curacao

Plantentuin Cascora
Willemstad

Cyprus

Officer-in-Charge
Plant Protection Section
Department of Agriculture
Ministry of Agriculture
and Natural Resources
Nicosia

Czechoslovakia

Plant Protection Service
Ministry for Agriculture and Food
Ministerstvo Zemedelstvi
a Lesniho Hospodarstvi
Ochrana Rostlin, Tesnov 65
Prague II

Denmark

Statens Plantetilsyn
Plant Protection Service
Gersonsvej 13
DK-2900
Hellerup

Dominica

Ministry of Agriculture,
Land and Cooperatives
Roseau

Dominican Republic

Dirección General de Foresta
Central de los Heroes
Santo Domingo

Ecuador

Jefe
Departamento de Sanidad Vegetal
Ministerio de Agricultura y Ganadería
Quito

Egypt

Ministry of Agriculture
Dokki
Cairo

Plant Quarantine Administration
Customs Gate 6
Alexandria

El Salvador

Jefe
Departamento de Defensa Agropecuaria
Ministerio de Agricultura y Ganadería
San Salvador

Ethiopia

Director
Department of Plant Production
and Protection
Plant Quarantine Service
Ministry of Agriculture and Settlement
Addis Ababa

Federal Republic of Germany

Federal Ministry of Food,
Agriculture and Forestry
Plant Protection Division
Rochusstr. 1
5300 Bonn

Fiji

Director of Agriculture
Department of Agriculture
Rodwell Road
Suva

Finland

Chief
Plant Quarantine Unit
Agricultural Research Centre
P.O. Box 18, SF-01301
Vantaa 30

France

(including overseas Department of
Guadeloupe, Guiana, Martinique, and
Reunion)

Ministère de l'Agriculture
Direction de la Qualité
Service de la Protection des Végétaux
231 Rue de la Convention
75015 Paris

*Martinique, Guadeloupe, and
French Guiana only*
Chef
Service de la Protection des Végétaux
Boite Postale 241
Fort de France
Martinique

Gabon
Direction des Services Agricoles
B.P. 43
Libreville

Gambia
Crop Protection Unit
Department of Agriculture
Cape St. Mary

German Democratic Republic
Staatl. Pfl.sch. u.
Pfl.quarantaendienst
Zentr. Staatl. A.f.
Pfl.sch. u. Pfl.quar.
Minist. f. Land-,
Forst- u. Na.güterwirtsch.
15 Potsdamm
Hermanswerde 20 A

Ghana
Plant Quarantine Service (PQS)
Department of Agriculture
P.O. Box M-37
Accra

Gilbert and Ellice Islands
Senior Agricultural Officer
Department of Agriculture
Bikenibeu, Tarawa

Great Britain

England and Wales
Plant Health Administrative Unit
Ministry of Agriculture,
Fisheries and Food
Eagle House
90-96 Cannon Street
London EC4N 7HT

Northern Ireland
Potatoes, Plant Health and Seeds
Department of Agriculture for
Northern Ireland
Dundonald House
Upper Newtownards Road
Belfast BT4 3SB

Scotland
Potatoes and Plant Health Branch
Department of Agriculture and
Fisheries for Scotland
Chesser House
500 Gorgie Road
Edinburgh EH11 3AW

Greece
The Director
Plant Protection Division
Ministry of Agriculture
2 Acharnon St.
Athens

Grenada
Plant Protection Officer
Plant Protection Division
Ministry of Agriculture,
Forestry and Fisheries
St. George's

Guam
The Director
Department of Agriculture
Agana 96910

Guatemala
Jefe
Departamento de Sanidad
Vegetal y Cuarentena Agrícola
12 Avenida 19-01
Zona 1
Ciudad de Guatemala

Guernsey, Channel Islands
The Secretary
State's Committee for Horticulture
Burnt Lane, St. Martins
Guernsey, C.I.

Guinea
Chef
Service de la Protection des Végétaux
Ministère de l'Economie rurale
et de l'Artisanat
Conakry

Guyana
Chief Agricultural Officer
Plant Quarantine Service
Ministry of Agriculture
Central Agricultural Station
Mon Repos, East Coast
Demerara

Haiti

SERA/DARNDR
Section de Botanique et
de Phytopathologie
Quarantaine Végétale
Damien
Port-au-Prince

Honduras

Programa de Producción y
Protección Vegetal
Ministerio de Recursos Naturales
Boulevard Miraflores
Tegucigalpa, D.C.

Hong Kong

Department of Agriculture
and Fisheries
Government Offices
393 Canton Road
Kowloon

Hungary

Deputy Head
Department of Plant Protection
and Agro-Chemistry
Ministry of Agriculture and Food
Kossuth, Lajes-Ter 11, Pf.1, H-1860
Budapest 55

Iceland

Agricultural Research Institute
Keldnaholt
110 Reykjavik

India

Plant Protection Adviser
to Government of India
Directorate of Plant Protection,
Quarantine and Storage
N.H.IV
Faridabad 121001
Haryana

Director
National Bureau of Plant
Genetic Resources
IARI Campus
New Delhi 110 012

Indonesia

Director General
Food Crops Development
Ministry of Agriculture
Jalan Salemba Raya 16
Jakarta

Iran

The Director
Sazeman Hefze Nabatat
(Plant Protection and
Quarantine Organization)
Ministry of Agriculture
Evin, Teheran

Iraq

The Director
Plant Quarantine Division
D.G. of Plant Protection
Ministry of Agriculture
Sadon Street
Baghdad

Ireland

Plant Protection Service
Department of Agriculture
and Fisheries
Kildare St.
Dublin 2

Israel

Director of the Department of
Plant Protection and Inspection
Ministry of Agriculture
P.O. Box 78
Beit Dagan, 50250

Italy

Chief
Division for the Protection
of Farm Crops
Ministry of Agriculture and Forestry
Directorate General for
Agricultural Production
Division II, Via XX Settembre
Rome

Jamaica

Chief Plant Protection Officer
Plant Protection Division
Ministry of Agriculture
P.O. Box 480
Hope, Kingston

Japan

Chief
Plant Protection Division
Agricultural Production Bureau
Ministry of Agriculture,
Forestry and Fisheries
1-2-1 Kasumigaseki
Chiyoda-ku
Tokyo

Jersey
 Chief Executive Officer
 Department of Agriculture
 6 Bond Street
 St. Helier, Jersey
 Channel Islands

Jordan
 Chief
 Plant Protection Division
 Agriculture Services Department
 Ministry of Agriculture
 Amman

Kampuchea
 Division de la Protection
 des Végétaux
 Director de l'Agriculture
 Department de l'Agriculture
 Phnom-Penh

Kenya
 Chief
 Plant Quarantine Section
 National Agricultural Laboratories
 Ministry of Agriculture
 P.O. Box 30028
 Nairobi

Korea, Republic of
 Ministry of Agriculture,
 Forestry and Fisheries
 Kwachon-City
 Kyunggi-do 171-11

Kuwait
 Plant Protection and Quarantine Section
 Ministry of Public Works
 Agriculture Department
 Kuwait

Laos
 Department de l'Agriculture
 Ministère de l'Agriculture
 Vientiane

Lebanon
 Department of Plant Protection
 and Quarantine
 Ministry of Agriculture
 Beirut

Lesotho
 Agricultural Research Division
 Ministry of Agriculture
 Box 289
 Maseru 100

Liberia
 The National Plant Quarantine Service
 Ministry of Agriculture
 Monrovia

Libya
 Plant Protection Section
 Ministry of Agriculture
 and Agrarian Reform
 Sidi Mesri
 Tripoli

Luxembourg
 Service de la Protection des Végétaux
 Administration des Services
 Techniques de l'Agriculture
 Boite Postale 1904
 Luxembourg - Ville

Madagascar
 Service de la Protection des Végétaux
 Boite Postale 1042
 Antananarivo

Malawi
 Chief
 Bvumbwe Research Station (Plant Imports)
 Ministry of Agriculture and
 Natural Resources
 Department of Agricultural Research
 P.O. Box 5748
 Limbe

Malaysia
 Crop Protection Branch
 Department of Agriculture
 Jalan Gallagher
 Kuala Lumpur

Mali
 Service de la Protection des Végétaux
 B.P. 1908
 Bamako

Malta
 Plant Health Division
 Department of Agriculture and Fisheries
 Government Experimental Farm
 Ghammieri
 Marsa

Mauritania
 Directeur
 Centre National de Recherche Agronomique
 et de Developpement Agricole (CNRA)
 Kaedi

Mauritius
 Plant Pathology Division
 (Plant Protection Service)
 Ministry of Agriculture and
 Natural Resoruces and Environment
 Reduit

Mexico
Dirección General de
Sanidad Vegetal (DGSV)
Av. Gmo. Perez Valenzuela No. 127
Coyoacan, D.F. (S.P. 21)

Montserrat
Ministry of Agriculture,
Lands, Trade and Housing
Plymouth

Morocco
Service de la Protection des Végétaux
Direction de la Recherche
Agronomique (DRA)
B.P. 415
Rabat

Mozambique
Chefe do Serviço
Serviços de Agricultura
Maputo

Nepal
Division of Plant Pathology
Department of Agriculture
P.O. Box 415
Khumaltar
Lalitpur

Netherlands
Director
Plant Protection Service
Geertjesweg 15
P.O. Box 9102
6700 HC Wageningen

New Caledonia
Chef
Service de l'Agriculture
B.P. 34
Noumea

New Zealand
Ministry of Agriculture and Fisheries
P.O. Box 2298
Wellington

Nicaragua
Departamento de Sanidad Vegetal
Ministerio de Agricultura y Ganadería
Managua, D.N.

Niger
Service de la Protection des Végétaux
Direction du Service de l'Agriculture
B.P. 323
Niamey

Nigeria
Project Director
Plant Quarantine Service
PMB 5672
Moor Plantation
Ibadan

Nieuw Island (Channel Islands)
Department of Agriculture
Alofi

Norway
Norwegian Plant Protection Service
B.P. 94
Okern
Oslo 5

Oman
Director
Administration of Plants
General Province of Agriculture
Ministry of Agriculture, Fisheries,
Petroleum and Minerals
P.O. Box 467
Muscat

Pakistan
Director
Department of Plant Protection
Ministry of Agriculture and
Works (Agriculture Division)
Jinnah Avenue, Malir Halt
Karachi 27

Panamá
Departamento de Investigación Agrícola
Sección de Cuarentena Agropecuaria
Ministerio de Agricultura
Comercio e Industrias
Panamá

Papua New Guinea
The Administrator
Department of Agriculture,
Stock and Fisheries
P.O. Box 2417
Konedobu

Paraguay
Jefe
División de Sanidad Vegetal
Ministerio de Agricultura y
Ganadería (MAG)
Asunción

Perú
Unidad de Inspección y Control Fitosanitario
Ministerio de Alimentación
10o. Piso
Ministerio de Trabajo
Av. Salaverry s/n
Lima

Philippines

Plant Quarantine Section
Bureau of Plant Industry (BPI)
692 San Andres
Malate, Manila

Poland

Ministerstwo Rolnictwa
(Ministry of Agriculture)
Department Produkcji Roslinnej
i Ochrony Roslin
(Department of Plant Production and
Protection)
Warszawa, ul.
Wspolna 30

Portugal

Reparticao de Servicos
Fitopatologicos (R.S.F)
Lisbon 3

Puerto Rico

Plant Quarantine Section
Puerto Rico Department
of Agriculture
P.O. Box 101613
Santurce 00908

Reunion Islands

Service de la Protection des Végétaux
Ministère de l'Agriculture
Boite Postal No. 312
Saint-Denis
Reunion

Romania

State Inspectorate for Plant Protection
Ministry of Agriculture
Bd Republicii 24
Bucharest

Rwanda

Bureau de la Protection Végétale
Ministère de l'Agriculture
et de l'Elevage
B.P. 621
Kigali

Saudi Arabia

Director
Plant Protection Branch
Ministry of Agriculture and Water
Riyadh

Senegal

Direction de la Protection des Végétaux
Ministère du Développement
Rurale et de l'Hydraulique
Boite Postale No. 486
Dakar

Seychelles

Director of Agriculture
Victoria
Mahe

Sierra Leone

Director of Agriculture
(Phytosanitary Control Unit)
MANR, Tower Hill
Freetown

Singapore

Senior Primary Production Officer
(Agriculture)
Primary Production Department
Ministry of National Development
Maxwell Road
Singapore 2

Solomon Islands

Ministry of Agriculture and Lands
Plant Quarantine Section
P.O. Box G 11
Honiara

Somalia

Plant Protection and Locust
Control Department
Ministry of Agriculture
Mogadishu

South Africa

Director
Division of Plant and Seed Control
Department of Agriculture
Private Bag X179
0001 Pretoria

Spain

Director
Instituto Nacional de Semillas y
Plantas de Vivero
Camino 2
Ciudad Universitaria
Madrid (3)

Sri Lanka

Director of Agriculture
Department of Agriculture
Ministry of Agricultural
Development and Research
Colombo

St. Kitts and Nevis

Ministry of Agriculture,
Housing and Labour
P.O. Box 39
Basseterre

St. Lucia
 Chief Agricultural Officer
 Department of Agriculture
 Ministry of Agriculture and Lands
 Castries

St. Vincent
 Chief Agricultural Officer
 Ministry of Agriculture, Trade
 and Tourism
 Kingston

Sudan
 Director
 Plant Protection Department
 Ministry of Agriculture and Irrigation
 P.O. Box 14
 Khartoum North

Surinam
 Plant Protection and Production Division
 Ministry of Agriculture,
 Animal Husbandry and Fisheries
 P.O. Box 1153
 Paramaribo

Swaziland
 Director of Agriculture
 Malkerns Research Station
 P.O. Box 4
 Malkerns

Sweden
 Plant Protection Service
 National Board of Agriculture
 S-551 83
 Jonkoping

Switzerland
 Division de l'Agriculture
 Department Federal de
 l'Economie Publique
 Mattenhofstrasse 5 3003
 Berne

Syria
 The Director
 Plant Protection and
 Quarantine Division
 Ministry of Agriculture and Agrarian Reform
 Damascus

Tahiti
 Monsieur le Chef du Service
 de l'Agriculture
 Pirae
 Papeete

Taiwan (Republic of China)
 Chief
 I Division (Plant Quarantine)
 Bureau of Commodity
 Inspection and Quarantine
 Ministry of Economic Affairs
 5 Hsu Chou Road
 Taipei

Tanzania
 Head
 Plant Quarantine Services
 TPRI
 Plant Quarantine
 Arusha

Thailand
 Director General
 Department of Agriculture
 Ministry of Agriculture and Cooperatives
 Bangkok 9

Togo
 Service de la Protection des Végétaux
 B.P. 1263
 Lomé

Tonga
 Plant and Animal Quarantine Service
 Ministry of Agriculture,
 Fisheries and Forestry
 P.O. Box 14
 Nuku'alofa

Trinidad and Tobago
 Technical Officer (Research)
 Ministry of Agriculture,
 Lands and Fisheries
 Centeno

**Trust Territory of
 the Pacific Islands**
 Chief of Agriculture
 Agricultural Division
 Department of Resources
 Saipan 96950
 Commonwealth of the Northern
 Marianas

Tunisia
 Direction de la Production Végétale
 Direction de la Défense des Cultures
 30 Rue Alain Savary
 Tunis

Turkey
 Ministry of Food,
 Agriculture and Animal Husbandry
 General Directorate of
 Plant Protection and Plant Quarantine
 Necatibey Cad. No. 98
 Ankara

**Tuvalu (Gilbert and
Ellice Islands)**
Ministry of Commerce and
Natural Resources
Funafuti

Uganda
Department of Agriculture
Plant Protection Unit
Kawanda Research Station
P.O. Box 7065
Kampala

**Union of Soviet
Socialist Republics**
State Department of
Plant protection
Ministry of Agriculture
State Inspectorate
Orlikov Pereulok 1/11
Moscow, 107139

United States of America
U.S. Department of Agriculture
Animal and Plant health
Inspection Service
Plant Protection Service
Quarantine Programs
(USDA/APHIS/PPQ)
302-E Administration Building
Washington, D.C. 20250

Uruguay
Dirección de Sanidad Vegetal
Ministerio de Agricultura y Pesca
Millan 4703
Montevideo

Vanuatu
Chef
Service de l'Agriculture
Port Villa

Venezuela
Director
Ministerio de Agricultura y Cría
Departamento de Sanidad Vegetal
Torre Este, Piso 20, Parque Central
El Conde
Caracas

Vietnam
Plant Protection Service
Department of Rural Affairs
P.O. Box 427
Hanoi

Virgin Islands
The Director
Department of Agriculture
Road Town
Tortola

Western Samoa
Director of Agriculture
Department of Agriculture and Forests
P.O. Box 206
Apia

Yemen Arab Republic
Plant Protection Division
Ministry of Agriculture
Sana'a

Yemen, People's Democratic Republic of
Plant Protection Project
El Ked Agricultural Research Centre
Ministry of Agriculture
Aden

Yugoslavia
Federal Committee for Agriculture
Department for Plant Protection
and Veterinary Service
Bulevar Avnoj-a 104
11070 Belgrade

Zaire
Inspecteur-Chef
Service du Controle Phytosanitaire
B.P. 8722
Kinshasa I

Zambia
The Phytosanitary Service
Mount Makulu Central Research Station
P.O. Box 7
Chilanga

Zimbabwe
Plant Protection Research Institute
Ministry of Agriculture
P.O. Box 8100
Causeway
Harare

