



CAB INTERNATIONAL

Issued 22nd May 1995

Phytopathological Papers, No. 34

**AN ANNOTATED LIST OF DISEASES,
PATHOGENS AND ASSOCIATED FUNGI
OF THE COMMON BEAN
(*Phaseolus vulgaris*)
IN EASTERN AND SOUTHERN AFRICA**

D.J. ALLEN



REF
SB
327
• A4
c.1

IAT

International Mycological Institute

An Institute of CAB INTERNATIONAL

CAB INTERNATIONAL is an intergovernmental organization providing services worldwide to agriculture, forestry, human health and the management of natural resources.

The information services maintain a computerized database containing over 2.7 million abstracts on agricultural and related research with 150,000 records added each year. This information is disseminated in 47 abstract journals, and also on CD-ROM and online. Other services include supporting development and training projects, and publishing a wide range of academic titles.

The four scientific institutes are centres of excellence for research and identification of organisms of agricultural and economic importance: they provide annual identifications of over 30,000 insect and microorganism specimens to scientists worldwide, and conduct international biological control projects.

International Mycological Institute

An Institute of CAB INTERNATIONAL

The Institute, founded in 1920, and employing over 70 staff:

- Is the largest mycological centre in the world, and is housed in a complex specially designed to support its requirements;
- Carries out research on a wide range of systematic and applied problems involving fungi (including lichens and yeasts), bacteria, and also on the preservation of fungi, the biochemical physiological and molecular characterization of strains, and on crop protection, environmental and industrial mycology, food spoilage, public health, biodeterioration and biodegradation;
- Provides an authoritative identification service, especially for microfungi of economic and environmental importance (other than certain human and animal pathogens), and for plant pathogenic bacteria and spoilage yeasts;
- Undertakes extensive computerized and indexing bibliographic work, including the preparation of 7 serial publications;
- Supports advice and undertakes a wide variety of project and culture work on crop protection, environmental, food spoilage and industrial topics.
- Offers training in pure and applied aspects of mycology and bacteriology, both at the Institute and overseas.

The Institute's dried reference collection numbers in excess of 355,000 specimens representing about 31,500 different species, and a genetic resource collection holds more than 17,000 living isolates by a variety of the most modern methods. The library receives about 600 current journals, and has extensive book and reprint holdings reflecting the Institute's interests.

International Mycological Institute
Bakeham Lane
Egham
Surrey TW20 9TY . ;
UK

Tel: (01784) 470111
Telex: 9312102252
Telecom Gold/Dialcom: 84:CAU009
Fax: (01784) 470909
E-mail: CABI-IMI@CGNET.COM

For full details of the services provided, please contact the Institute Director.

**AN ANNOTATED LIST OF DISEASES,
PATHOGENS AND ASSOCIATED FUNGI
OF
THE COMMON BEAN
(*Phaseolus vulgaris*)
IN
EASTERN AND SOUTHERN AFRICA**

By

D.J. ALLEN


Formerly Plant Pathologist.

Bean Program

*Centro Internacional de Agricultura Tropical (CIAT),
Arusha, Tanzania*

Present Address:

*Higher Quantock, Stockland
Honiton, Devon EX14 9DX
UK*



UNIVERSITY DOCUMENTATION
DOCUMENTATION

020636

CAB INTERNATIONAL

CIAT

28 SET. 1995

CAB INTERNATIONAL
Wallingford
Oxon OX10 8DE
UK

Tel: Wallingford +44 (0)1491 832111
Telex: 847964 (COMAGG G)
E-mail: cabi@cabi.org
Fax: +44 (0)1491 833508

© CAB INTERNATIONAL 1995. All rights reserved. No part of this publication may be reproduced in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without the prior permission of the copyright owners

In Association with
CIAT (Centro Internacional
de Agricultura Tropical)
Apartado aéreo 6713
Cali
Colombia

Tel: 57-23-675050
Fax: 57-23-647243

A catalogue entry for this book is available from the British Library.

ISBN 0 85198 938 1
ISSN 0069-7141

CONTENTS

	Page
Summary	v
Introduction	vi
Acknowledgements	ix
Resume	x
Introduction	xi
Reconnaissance	xv
Resumo	xvi
Introducao	xvii
Agradecimentos	xx
 Annotated territorial lists	
Angola (AN)	1
Burundi (BU)	2
Ethiopia (ET)	3
Kenya (KE)	5
Lesotho (LO)	8
Madagascar (MD)	8
Malawi (MW)	9
Mauritius (MA)	12
Mozambique (MO)	13
Rwanda (RW)	14
Seychelles (SY)	15
Somalia (SO)	16
South Africa (SA)	16
Sudan (SD)	18
Swaziland (WD)	19
Tanzania (TZ)	20
Uganda (UG)	22
Zaire (ZR)	24
Zambia (ZA)	26
Zimbabwe (ZW)	28
 Pathogen indices	
Fungi	30
Bacteria	33
Viruses	34

Distribution of pathogenic variants	35
References	36

SUMMARY

This list brings together many widely scattered unpublished records, from the International Mycological Institute, from the Centro Internacional de Agricultura Tropical, and from field surveys conducted during recent collaborative research between CIAT and Horticulture Research International, Wellesbourne, and the Institute for Biochemistry and Plant Virology, Braunschweig. The literature survey has been extensive but probably not exhaustive: both recent and old checklists of published information have been scanned to provide an historical perspective to the presence of pathogens within a given territory. The list is confined to pathogens and diseases recorded on the common bean (*Phaseolus vulgaris* L.) in 20 countries of eastern, central and southern Africa and its off-shore islands. Information is given on their prevalence and pathogenic variability, with a comprehensive bibliography of more than 150 references. The list should prove valuable to plant pathologists, bean scientists, plant quarantine officers and research directors concerned with crop improvement and the international exchange of genetic resources.

INTRODUCTION

The common bean (*Phaseolus vulgaris* L.) was introduced into Africa from each of two gene centres in Latin America during the past four centuries. Africa is now the second most important bean producing region of the tropics, following Latin America. The total average annual production of dry beans in Africa is estimated at about two million tonnes, some 25 per cent of world production, but as a large part of production is consumed locally, official statistics may underestimate total production. Kenya is the largest producer in Africa, followed in some order by Tanzania, Uganda, Burundi, Rwanda and Zaire, all having more than 270,000 ha under production. Other significant producers in eastern and southern Africa include Angola, Ethiopia, Malawi, Mozambique and Madagascar. Although bean production in Africa has increased in some areas over the past decade, this has been achieved through area increases. Over eastern and southern Africa as a whole, production was stagnant over the 1980s, a period when population grew at an annual rate of over three per cent. The result is that consumption per caput is falling, albeit in an area where beans play a critical role in human nutrition by providing as much as 45 per cent of total protein consumed in some areas (Grisley, 1990; Allen & Smithson, 1991).

Small farmers are the principal producers of beans in most of Africa where the crop is cultivated in a complex array of cropping systems, including the use of cultivar mixtures. Most production is for subsistence and, although its role as a small-scale cash crop is increasing, the use of fertilizers and pesticides is uncommon. Soil infertility, periodic water stress, insect pests and diseases are considered the principal agronomic constraints associated with low yields (Allen *et al.*, 1989; Allen & Edje, 1990). Reviews of the principal bean diseases in Africa and the place of disease among other production constraints of the crop have been compiled recently (Allen, 1987a; Allen *et al.*, 1989; Allen *et al.*, in prep.). Almost 100 pathogens attack the common bean in Africa but only about eight cause substantial economic damage. Diseases like angular leaf spot, anthracnose, rust, common bacterial blight and bean common mosaic are widespread and can decrease seed yield considerably. Other diseases, including ascochyta blight, white mould and halo blight, can also cause significant crop loss, but they tend to be confined to specific environments.

For small scale farmers in sub-Saharan Africa, host plant resistance remains the only feasible means of disease control. At present, many farmers obtain a crop only by planting late so as to avoid the wettest season; improved levels of varietal resistance to disease could give added yield benefits by permitting more timely sowing. Scientists in the national agricultural research systems (NARS) of eastern and southern Africa are devoting varying degrees of effort and resources to the improvement of the common bean, including the evaluation of both local and introduced germplasm. There is growing evidence that immediate, substantial benefit can be gained directly from the large genetic resources and new breeding materials that are available through introduction, especially from the Centro Internacional de Agricultura Tropical (CIAT) in Colombia (Allen & Smithson, 1991), and other centres with well-established breeding programmes. This strategy has enabled many countries in Africa to release new, heavier yielding bean cultivars during the past few years. No international exchange of germplasm can be completely free of risk. East Africa, in particular, has a long history and sound record of operating a closed quarantine service as a safeguard against the accidental introduction of alien pathogens and their races into the region (Sheffield, 1968). Owing to the seed-borne nature of many of the pathogens of the common bean, most of the major diseases occur almost wherever the crop is now cultivated. In some cases, new outbreaks of pests and diseases are attributable not to the

introduction of the causal agent but of novel susceptibility to an indigenous one, presenting a 'new encounter' (Allen, 1983). An example from Africa is the bean stem maggot (*Ophiomyia* sp.) which is an Old World insect pest of a New World introduction, *Phaseolus vulgaris*. New diseases may also arise through changes in crop husbandry or the increasing use of marginal land. For example, root rots, caused by a complex of fungi including *Pythium* species, are apparently increasing through crop intensification (Buruchara, 1992c), while ashy stem blight can be severe in marginal areas subject to drought stress.

It is obviously useful to document what pathogens are present in a given area before substantial introductions are made. Territorial checklists provide a valuable basis for the formulation of regulations governing import by listing those pathogens that are known to be present. As regards the common bean, points of contrast between Africa and Latin America are valuable. Although most of the major pathogens occur in both continents, there are some exceptions: the scab fungus (*Elsinoë phaseoli* Jenk.) is not known on *P. vulgaris* in the Americas, whereas it is fairly widespread in Africa. Conversely, fungi like *Pseudocercospora albida* (Matta & Bell.) Deighton, *Chaetosporia wellmanii* Stev., *Entyloma* sp., *Cercospora vanderystii* Henn. and *Cercospora castellanii* Matta & Bell. are not known to occur in Africa (Allen, 1983; Schwartz, 1989). Viruses not yet recorded from natural infections of beans in Africa include bean mild mosaic, bean rugose mosaic, bean pod mottle, quail pea mosaic, cowpea chlorotic mottle, bean golden mosaic, bean dwarf mosaic and several other geminiviruses that are not seed-borne (Galvez & Morales, 1989; Morales & Gamez, 1989). Among bacteria, there stands alone the strange case of *Curtobacterium flaccumfaciens* (Hedges) Collins & Jones that deserves further comment. *C. flaccumfaciens* has been known for many years to cause a bacterial wilt in common bean in Nebraska and elsewhere in the USA (Mohan & Hagedorn, 1989). Owing to its seed-borne nature and its apparently restricted geographical distribution despite its occurrence in both America and Africa, *C. flaccumfaciens* has been assigned to the high risk category of quarantine objects (Neergaard, 1977) with the result that this pathogen is the 'hot potato' among bean pathogens in the eyes of plant quarantine officials in Africa. With the benefit now of hindsight, it seems timely to review this status. There is little evidence to suggest that bacterial wilt is economically important, even in the USA (J.R. Steadman & M.J. Silbernagel, pers. comm., 1988). *C. flaccumfaciens* is a common soil inhabitant also in Latin America: the bacterium is relatively easy to identify but its role as a pathogen is less easily proven (M.A. Pastor-Corrales, pers. comm., 1987). Furthermore, the presence of the bacterium on other legume genera including *Zornia* seems not necessarily to constitute a threat to the bean crop. These factors together call to question the epidemiological competence of *C. flaccumfaciens* which seems to have been accorded undue prominence in quarantine legislation among pathogens of the common bean.

With the recent development in Africa both of national bean breeding programmes and of regional networks for germplasm exchange, it is no longer only farmers who have interest vested in the transport of bean seed across national boundaries. Yet a lack of knowledge of the risks involved sometimes makes it difficult for countries to collaborate effectively in research on the crop's improvement. At a meeting of African agricultural research directors held in Nairobi in June 1992, it was recognized that "plant quarantine regulations, while intended to prevent the accidental introduction of diseases not present in (a given) country, usually apply knowledge on disease distribution that has not been updated (for many years)... It was agreed that CIAT, in collaboration with pathologists in the various national agricultural research systems, should prepare a distribution map of bean diseases for the attention of NARS directors who would use the

information in following up with the quarantine services". This meeting was preceded by a Pan African Working Group Meeting on Fungal Pathogens of Beans, held in Thika, Kenya in May 1992, when CIAT was given the task of compiling a checklist of bean pathogens, as a first step toward drawing the disease distribution maps that are foreseen as necessary. The present list, then, has the following principal purposes: to provide a sound scientific basis for the preparation of disease distribution maps from which to update quarantine legislation governing the movement of bean seed into and within Africa; to provide plant quarantine officers with the information necessary to assess the level and type of risk associated with each individual introduction so as to guide decisions on procedures appropriate to the case; to provide a basis for priority setting by crop improvement teams by giving access to recent information generated from regional surveys of pathogens and their variants; and to provide a basis for site selection in screening for disease resistance. The scope of the list embraces 20 countries of eastern, central and southern Africa and its off-shore islands which together constitute all countries wherein the common bean is an important crop. Since their inception during 1983-1987, CIAT's three regional bean programmes in Africa have catalyzed the development of a research and training network across eastern and southern Africa, and the majority of the territories included in the list are partners in this bean network. The list omits Botswana and Namibia where the common bean is an insignificant crop and no disease records have been found. The literature survey has been extensive but probably not exhaustive. The pathogen and disease records themselves cover many old as well as new records, to provide an historical perspective of their distribution. Substantial use has been made of territorial checklists, where these exist. Supplementary records have been abstracted from other reliable sources and much use has been made of determinations made at the International Mycological Institute, an Institute of CAB INTERNATIONAL, where collections are assigned IMI numbers. Pathogen nomenclature follows IMI usage as given in Holliday (1980). Records are confined to the single host species, *Phaseolus vulgaris*. In many instances, a particular pathogen (or race) may be present in a certain country on a host plant other than *P. vulgaris*, indicating the importance of distinguishing between the geographical distribution of a disease and the distribution of the causal pathogen; they are not synonymous, as stressed elsewhere (Allen, 1983) and illustrated by the distribution of races of the halo blight bacterium (Table 1). During the course of recent surveys, various pathogens have been recorded from legume hosts other than *P. vulgaris* and these records have been published elsewhere (Allen, 1991). Obvious gaps in our knowledge of bean pathogens in Africa remain, and no checklist can ever claim to be complete. As surveys generate new records, it would be most desirable that these are collated on a regional basis and published, perhaps as an annual supplement in an appropriate journal, like the FAO Plant Protection Bulletin.

ACKNOWLEDGEMENTS

A publication of this type inevitably relies substantially upon the help, support and guidance of many people and agencies. At the outset, the Director and staff of the International Mycological Institute gave me access to unpublished records and gave guidance on pathogen nomenclature, supplemented by my constant reference to Paul Holliday's *Fungus Diseases of Tropical Crops*. Mr Peter Filby greatly assisted my access to the important and under-utilized Buttress Collection of the Scientific Periodicals Library in Cambridge, where Dr S. Dale's help is also acknowledged. Drs C.Y.L. Schotman and R. Ikin kindly gave me access to the FAO Global Plant Quarantine Information System.

Country lists have been circulated in draft to plant pathologists and bean programme leaders who have kindly commented on the list's content. I should like particularly to thank the following for their help which, in some cases, included giving access to unpublished information:

Dr J. Arias, Dr R. Buruchara, Dr N. Govinden, Ato Habtu Assefa, Dr H.C. Haciwa, Dr J. Kannaiyan, Mr J. Kayitare, Dr A. Liebenberg, Dr R.J.M. Melis, Dr O.Z. Mukoko, Mr N. Ntahimpera, Mr M.E. Omunyin, Dr A.F. Opio, Dr D. Perreaux, Dr A. Phillips, Mr Pyndji Mukishi, Mr A.H. Ramos, Prof. Y.P. Rao, Mr S. Saumtally, Dr V. Schmit, Mrs T. Sengooba, Dr J.B. Smithson, Dr N. Spence, Prof. J.M. Teri, Dr D.M. Teverson, Dr P. Trutmann and Dr H.J. Vetten.

Dr B.C. Sutton has provided most valuable guidance during the final stages of production.

At this point, I should state plainly that the responsibility for the technical content of the list rests solely with myself: my comments do not necessarily reflect official opinion either of CABI or of CIAT by whom it is jointly published. Dr R.A. Kirkby has kindly provided the French and Portuguese translations as well as encouragement to the completion of the task which has been supported throughout by funds donated by the Canadian International Development Agency to the Centro Internacional de Agricultura Tropical in its execution of projects for the development of bean research networks in Africa.

Finally, I should add that my largest debt is to my wife, Leonora, who has helped me directly throughout the long period of compilation and editing of the entire booklet.

DJA
Cambridge, January, 1994

RESUME

Cette liste rassemble beaucoup de rapports d'études et résultats bruts fort dispersés et non publiés, provenant de l'Institut International de Mycologie, du Centre International d'Agriculture Tropicale et des inventaires sur terrain menés pendant des recherches récentes conduites en coopération entre CIAT et "Recherche Internationale en Horticulture", Wellesbourne, et l'Institut de Biochimie et de Virologie des plantes, Braunschweig.

La littérature a été intensivement inventoriée mais cela n'a probablement pas été exhaustif: les deux listes de contrôle sur l'information publiée ont été vérifiées en vue de fournir une perspective historique à la présence des pathogènes dans un territoire donné. La liste se limite aux pathogènes et maladies mentionnés sur le haricot commun (*Phaseolus vulgaris* L.) dans vingt (20) pays de l'Afrique orientale, centrale et australe ainsi que dans les îles environnantes. L'information est donnée sur leur prévalence et leur variabilité pathogénique, avec une large bibliographie compréhensible de plus de 150 références. La liste devrait pouvoir présenter une grande valeur pour les phytopathologistes, les spécialistes du haricot, les responsables de la quarantaine des plantes et les directeurs de recherche concernés par l'amélioration des cultures et l'échange international des ressources génétiques.

INTRODUCTION

Le haricot commun (*Phaseolus vulgaris* L.) était introduit en Afrique en provenance de chacun des deux centres de gênes en Amérique Latine durant les 4 derniers siècles. L'Afrique est maintenant la deuxième région tropicale la plus importante du point de vue de la production du haricot après l'Amérique Latine. La moyenne de la production annuelle totale du haricot sec en Afrique est estimée à environ deux millions de tonnes, soit 25 pour cent de la production mondiale; mais comme une grande partie de la production est consommée localement, les statistiques officielles pourraient sous-estimer la production totale.

Le Kenya est le plus grand producteur en Afrique, suivi en quelque ordre par la Tanzanie, l'Uganda, le Burundi, le Rwanda et le Zaïre; tous ayant plus de 270,000 ha sous production. D'autres producteurs importants en Afrique de l'Est et Australe sont: l'Angola, l'Ethiopie, le Malawi, le Mozambique et la Madagascar. Bien que la productivité du haricot en Afrique a augmenté dans quelques parties au courrant de la décennie écoulée, l'augmentation moyenne de la productivité est le résultat des augmentations par zones. À travers l'Afrique orientale et australie d'une manière générale, la production était stagnante durant les années 80, période pendant laquelle la population s'est accrue à un taux annuel supérieur à 3 per cent. En conséquence, la consommation par tête va en décroissant, bien que ce soit une zone où le haricot joue un rôle primordial dans la nutrition humaine, en fournissant autant que 45 per cent des protéines totales consommées dans certains coins (Grisley, 1990; Allen & Smithson, 1991).

Les petits fermiers sont les principaux producteurs des haricots dans la plupart des pays africains où la plante est cultivée dans un arrangement complexe de systèmes cultureaux, incluant l'usage de mélanges variétaux.

La plus grande partie de la production sert à la subsistance et, bien que son rôle en tant que culture de rente à petite échelle s'accroît, l'usage des fertilisants et pesticides est inhabituel. L'infertilité du sol, le stress hydrique périodique, les pestes et les maladies sont considérés comme les principales contraintes agronomiques justifiant les faibles rendements (Allen *et al.*, 1988; Allen & Edje, 1990). Les études des principales maladies du haricot en Afrique et la place des maladies parmi d'autres contraintes à sa production ont été inventoriées récemment (Allen, 1987a; Allen *et al.*, en impression). Près de 100 pathogènes attaquent le haricot commun en Afrique; mais seulement huit (8) d'entre eux causent des dommages économiques substantiels.

Des maladies telles que les taches anguleuses, l'anthracnose, la rouille, la bactériose commune et la mosaïque commune du haricot sont largement répandues et peuvent contribuer à décroître considérablement le rendement en graines. D'autres maladies comme l'ascocytose, la pourriture blanche et la bactériose à halo, peuvent aussi causer d'importantes baisses de productions; mais ces maladies tendent à se confiner dans des environnements bien spécifiques.

Pour les petits fermiers en Afrique sub-saharienne, la résistance de la plante hôte reste le seul moyen possible pour le contrôle des maladies. Pour le moment, beaucoup de fermiers ne peuvent obtenir une récolte qu'en cultivant tard, afin d'éviter la période plus humide. Les niveaux élevés de résistance des variétés aux maladies pourraient permettre d'accroître le rendement, parce qu'on peut alors semer à temps opportun.

Les scientifiques opérant dans des systèmes nationaux de recherches agronomiques (SNRA) de l'Afrique orientale et australe consentent des efforts variés ainsi que des ressources en vue de l'amélioration du haricot commun, y compris l'évaluation du germoplasme local et introduit.

Il est de plus en plus évident qu'un bénéfice substantiel immédiat pourrait être tiré de l'usage de

grandes ressources génétiques disponibles et du nouveau matériel amélioré, spécialement au niveau du CIAT en Colombie (Allen & Smithson, 1991) et d'autres centres ayant des programmes d'amélioration bien établis. Cette stratégie a permis à beaucoup de pays africains de rendre disponibles des nouvelles variétés à plus haut rendement pendant ces dernières années.

Aucun échange international du germoplasme ne peut être dépourvu de risques. L'Afrique de l'Est, en particulier, a une longue histoire et informations appropriées sur son service strict de la quarantaine en guise de sécurité contre l'introduction accidentelle des pathogènes étrangers et leurs races dans la région (Sheffield, 1968).

Dû au fait que beaucoup de pathogènes du haricot commun se transmettent par semences, la plupart des maladies apparaissent presque partout où la plante est maintenant cultivée.

Dans certains cas, les nouvelles éruptions des pestes et des maladies sont causées non pas par l'introduction de l'agent causal, mais plutôt par une susceptibilité singulière du matériel vis-à-vis d'agents locaux du pays hôte (Allen, 1983). Un exemple en provenance de l'Afrique est la mouche du haricot (*Ophiomyia* spp.) qui est une peste du Vieux Monde sur une culture du haricot (*Phaseolus vulgaris*) introduit à partir du Nouveau Monde. Des nouvelles maladies peuvent aussi provenir des modifications des pratiques agricoles ou du recours accru aux sols marginaux. Par exemple, les pourritures racinaires causées par une série de champignons incluant des espèces de *Pythium*, s'accentuent apparemment avec l'intensification de la production du haricot (Buruchara, 1992c) pendant que la pourriture charbonneuse de tiges peut être sévère sur les sols marginaux sujets à la sécheresse.

Il est clairement important de s'informer sur l'existence des pathogènes dans une zone donnée avant de procéder aux introductions substantielles.

Les listes de contrôle par territoires fournissent une base précieuse pour l'élaboration des mesures devant réglementer l'importation, en indiquant les pathogènes qui sont connus être présents.

Concernant le haricot commun, les points de différence entre l'Afrique et l'Amérique Latine sont essentiels. Bien que la plupart des pathogènes apparaissent dans les deux continents, il y a quelques exceptions: le champignon qui cause la gale (*Elsinoë phaseoli* Jenk.) n'est pas connu sur *P. vulgaris* aux Amériques, tandis qu'il est largement répandu en Afrique. Par contre, les champignons tels que *Pseudocercospora albida* (Matta & Bell.) Deighton, *Chaetoseptoria wellmanii* Stev., *Entyloma* sp., *Cercospora vanderystii* Henn. et *Cercospora castellanii* Matta & Bell. n'apparaissent pas en Afrique (Allen, 1983; Schwartz, 1989).

Les virus non encore repérés à partir des infections naturelles des haricots en Afrique comprennent "bean mild mosaic", "bean rugose mosaic", "bean pod mottle", "quail pea mosaic", "cowpea chlorotic mottle", "bean golden mosaic", "bean dwarf mosaic" et plusieurs autres geminivirus qui ne sont pas transmis par les semences (Galvez & Morales, 1989; Morales & Gamez, 1989). Parmi les bactéries, il y a l'étrange cas de *Curtobacterium flaccumfaciens* (Hedges) Collins & Jones qui mérite plus de commentaires. *C. flaccumfaciens* a été connu depuis plusieurs années comme responsable du flétrissement bactérien du haricot commun au Nebraska et ailleurs aux Etats-Unis (Mohan & Hagedorn, 1989). Dû à la nature de sa transmission par graines et de sa distribution géographique apparemment réduites malgré son apparition aussi bien en Amérique qu'en Afrique, *C. flaccumfaciens* a été classé comme un des pathogènes à haut risque mis en quarantaine (Neergaard, 1977) et comme conséquence, ce pathogène est considéré aux yeux des responsables officiels de la quarantaine des plantes en Afrique comme le plus dangereux des pathogènes qui affectent le haricot. Cependant, grâce aux informations rétrospectives, l'on devrait

à présent redéfinir la place de ce pathogène. L'on ne dispose pas de preuves suffisantes pour suggérer que le frétrissement bactérien serait économiquement important, même aux Etats-Unis d'Amérique (J.R. Steadman & M.J. Silbernagel, communication personnelle, 1988). *C. flaccumfaciens* se trouve communément aussi dans le sol de l'Amérique Latine: il est relativement facile d'identifier cette bactérie, cependant son rôle en tant que pathogène est moins facilement prouvé (M.A. Pastor-Corrales, communication pesonnelle, 1987). En plus, la présence de la bactérie dans d'autres genres des légumineuses dont le *Zornia* ne semble pas nécessairement constituer une menace au haricot. Tous ces facteurs mis ensemble remettent en cause la capacité épidémiologique du *C. flaccumfaciens* auquel on semble avoir donné trop d'importance dans la législation en quarantaine parmi les pathogènes du haricot commun.

Avec le développement récent en Afrique aussi bien des programmes nationaux d'amélioration du haricot que des réseaux régionaux d'échange du germoplasme, ce ne sont plus seulement les fermiers qui sont intéressés au transport des semences de haricot au-delà des frontières nationales. Le manque de connaissance des risques qui y interviennent quelque fois continue à rendre difficile la collaboration effective des pays, en matière de recherche pour l'amélioration du haricot. Au cours d'une réunion des directeurs de recherches agronomiques en Afrique tenue à Nairobi au mois de juin 1992, il a été reconnu que "les mesures de quarantaine des plantes, bien qu'elles visent de prévenir l'introduction accidentelle des maladies inexistantes dans un pays donné, utilisent souvent de connaissances sur la propagation des maladies qui ne sont pas actualisées (depuis plusieurs années). Il a été convenu que CIAT, en collaboration avec les phytopathologistes de plusieurs institutions nationales de recherches agronomiques, devra élaborer une carte sur la distribution géographique des maladies du haricot à l'attention des directeurs des services nationaux de recherches agronomiques. Ceux-ci utiliseront cette information dans le suivi des services de quarantaines". Cette réunion a été précédée par un atelier du Groupe Panafricain de Travail sur les champignons pathogènes du haricot, tenu à Thika au Kenya, au mois de mai 1992, qui avait demandé au CIAT d'élaborer une liste de contrôle d'agents pathogènes du haricot, comme première étape dans l'établissement des cartes de distribution des maladies, qui seront très utiles.

La liste actuelle, renferme les préoccupations suivantes:

- fournir une base scientifique pour la préparation des cartes de distribution des maladies devant être utilisées pour actualiser la législation relative à la quarantaine, qui régit le mouvement des semences de haricot de l'extérieur et à l'intérieur de l'Afrique;
- fournir au personnel responsable de la quarantaine des plantes, l'information nécessaire pour pouvoir évaluer le niveau et le type de risque liés à chaque introduction nouvelle, afin de proposer des mesures appropriées à prendre pour chaque cas;
- fournir une base pour l'établissement des priorités par les équipes chargées de l'amélioration des plantes, en leur donnant accès aux informations récentes générées par des enquêtes régionales des pathogènes et de leurs races; et
- fournir une base pour la sélection des sites lorsqu'on veut faire le criblage pour la résistance aux maladies.

La liste embrasse 20 pays de l'Afrique orientale, centrale et australie et les îles environnantes, lesquels dans leur ensemble, constituent des pays où le haricot commun est un produit important. Depuis leur établissement (1983-1987), les trois programmes régionaux de haricot du CIAT en Afrique ont développé un réseau de recherche et de formation à travers l'Afrique orientale et

australe et la majorité des territoires inclus sur la liste sont partenaires de ce réseau. La liste n'inclue pas le Botswana et la Namibie, pays où le haricot commun est un produit négligeable et où aucune maladie n'a été repérée. L'inventaire de la littérature a été intensif, mais certes, pas exhaustif. Les pathogènes et les maladies enregistrés proviennent des informations anciennes aussi bien que récentes, dans le but de donner une perspective historique à leur distribution.

Les listes de contrôle par territoire ont été largement utilisées partout où elles existent. Des résumés d'information supplémentaires ont été élaborés à partir d'autres sources fiables et l'on a beaucoup fait usage des évaluations faites à l'Institut International de Mycologie, un institut faisant partie du CAB INTERNATIONAL, où les collections sont assignées des nombres IMI.

La nomenclature des pathogènes suit le système utilisé par IMI tels qu'indiqué dans Holliday (1980). Les informations se concentrent sur l'unique espèce hôte, *Phaseolus vulgaris*. Dans beaucoup de cas, un pathogène particulier (ou race), peut être présent dans un pays donné sur une plante hôte autre que *P. vulgaris*; ceci montre l'importance qu'il y a de faire la distinction entre la distribution géographique d'une maladie et celle de l'agent causal. Il s'agit de deux choses différentes, comme c'est clarifié ailleurs (Allen, 1983) et illustré par la distribution des races du pathogène de la bactériose à halo (Table 1).

Au cours de récentes enquêtes, plusieurs pathogènes ont été enregistrés sur des légumineuses hôtes autres que *P. vulgaris* et ces résultats ont été publiés ailleurs (Allen, 1991). Toutefois, des limites évidentes demeurent concernant notre connaissance des pathogènes du haricot en Afrique et aucune liste de contrôle ne peut se réclamer être exhaustive.

Comme les études continueront de fournir de nouveaux résultats, il serait très souhaitable que ces informations soient rassemblées sur la base régionale et publiées, peut-être comme un supplément annuel dans un journal approprié tel que le bulletin de la FAO sur la protection des plantes.

RECONNAISSANCE

Une publication de ce genre dépend très sérieusement de l'aide, de l'appui et des conseils de plusieurs personnes et agences. Tout au début, le Directeur et les membres du personnel de l'Institut International de Mycologie m'ont donné accès aux informations non encore publiées et ont fourni des conseils quant à la nomenclature des pathogènes. Ceci a été complété par ma référence constante à l'ouvrage de Paul Holliday sur les *maladies cryptogamiques des plantes tropicales*. M. Pierre Filby m'a grandement assisté pour accéder à l'importante mais sous-utilisée Collection Buttress, à la librairie de Périodiques Scientifiques de l'Université de Cambridge, où je reconnais avoir également reçu l'assistance du Dr S. Dale. Drs C.Y.L. Schotman et R. Ikin, m'ont gentiment donné accès au système d'information global de la FAO sur la quarantaine des plantes.

Les listes par pays avaient été circulées sous forme de brouillon aux phytopathologistes ainsi qu'aux Chefs des programmes haricot, lesquels ont bien gentiment fourni leurs commentaires sur le contenu desdites listes. J'aimerais tout particulièrement remercier les personnes suivantes pour leur assistance qui, dans certains cas, consistait à me faire prendre connaissance des informations non publiées:

Dr J. Arias, Dr R. Buruchara, Dr N. Govinden, Ato Habtu Assefa, Dr H.C. Haciwa, Dr J. Kannaiyan, Mr J. Kayitare, Dr A. Liebenberg, Dr R.J.M. Melis, Dr O.Z. Mukoko, Mr N. Ntahimpera, Mr M.E. Omunyin, Dr A.F. Opio, Dr D. Perreaux, Dr A. Phillips, Mr Mukishi Pyndji, Mr A.H. Ramos, Prof Y.P. Rao, Mr S. Saumtally, Dr V. Schmit, Mrs T. Sengooba, Dr J.B. Smithson, Dr N. Spence, Prof J.M. Teri, Dr D.M. Teverson, Dr P. Trutmann et Dr H.J. Vetten.

Dr B.C. Sutton a fourni des conseils précieux durant l'étape finale de la production du document.

A ce stade, je dois dire pleinement que le contenu technique de la liste n'engage que moi-même: mes commentaires ne reflètent pas nécessairement l'opinion officielle ni de CABI ni du CIAT avec lesquels nous publions conjointement. Dr R.A. Kirkby nous a gentiment encouragé à parachever notre tâche qui a été financé par des fonds offerts par l'Agence Canadienne pour le Développement International (ACDI) au CIAT pour l'exécution des projets d'établissement d'un réseau de recherche sur le haricot en Afrique.

Finallement, je dois ajouter que ma plus grande dette demeure envers ma femme, Leonora, qui m'a aidé durant la longue période de dépouillement et d'édition du livret tout entier.

DJA
Cambridge, January 1994

RESUMO

Esta lista tráz um conjunto de vários registros espalhados na publicados pelo o Instituto Internacional de Micología, o Centro Internacional da Agricultura Tropical e a investigacao de campo conduzida recentemente em colaboracao com o CIAT e a Investigacao Internacional de Horticultura, Wellesbourne, o Instituto Bioquímico e Virólogia das plantas, Braunschweig.

A revisao da bibliografia foi extensiva mas provavelmente nao exautiva, abarcou os registos antigos e recentes da informacao publicada que foi selecionada para dar uma perspectiva histórica da presencia dos patogenos num determinado território.

A lista está confinada aos patogenos e as doenças registradas no Feijao vulgar (*Phaseolus vulgaris* L.) em 20 países da Africa Ocidental, Central e Austral; e as suas ilhas vizinhas.

A informacao é dada conforme a prevalencia e variabilidade do patogeno bem como, a informacao bibliografica compreendida em mais de 150 referencias.

As listas podem ser úteis para os fitopatologistas, os oficiais da quarantena das plantas e os directores de investigacao envolvidos no melhoramento da cultura e a troca os recursos genéticos internacionais.

INTRODUCAO

O feijao vulgar (*Phaseolus vulgaris* L.) foi introduzido em África pelos dois centros genéticos da América Latina há quatro séculos passados. Depois da América Latina, África actualmente ocupa o segundo lugar entre os países tropicais. A produçao média total é de cerca de dois milhões de toneladas, que é 25% de toda a produçao mundial. Porém, como uma grande parte da produçao é consumida localmente, as estatísticas oficiais podem subestimar a produçao total. Em África, o Quénia é o maior produtor seguido pela Tanzânia, Uganda, Burundi, Ruanda e o Zaire; todos estes países em conjunto ocupam 270.000 Ha em produçao. Os outros produtores importantes no Este e Sul de África incluem Angola, Etiópia, Malawi, Mocambique e Madagáscar. A area de produçao do feijao em África aumentou em certas regiões, na década passada. Este aumentou, deveu-se a extensão da area de produçao. No Este e Sul de África, a produçao foi estagnante nos anos 80, período em que, a populacão cresceu numa taxa anual de mais de 3%. O resultado desse aumento foi a diminuição das areas de produçao nos lugares onde o feijao vulgar tem um papel importante na nutrição humana através do fornecimento de 45% de proteína total consumida (Grisley, 1990; Allen & Smithson, 1991).

Os pequenos agricultores são os principais produtores do feijao vulgar em África onde a cultura é cultivada num sistema de cultivo complexo, incluindo o uso das misturas de cultivares. Quase toda a produçao é para a subsistência e a sua comercialização tem vindo a aumentar. O uso de fertilizantes e pesticidas não é muito comum. A fertilidade do solo; a falta periódica das chuvas; as pragas e doenças são consideradas as principais limitantes agronómicas associados a baixa produçao (Allen et al., 1989; Allen & Edje, 1990). A revisão bibliográfica sobre as principais doenças e, distribuição das mesmas entre as outras limitantes de produçao da cultura foi recentemente compilado (Allen, 1987a; Allen et al., 1989; Allen et al., em impressão). Cerca de 100 patógenos atacam o feijao vulgar em África mas, apenas 8% causa de perda económica da produçao. As doenças como a mancha angular da folha, antracose, ferrugem, mancha bacteriana comum, o mosaico comum estão largamente espalhados em várias regiões e podem baixar consideravelmente a produçao do grão. Outras doenças incluem "ascocytha blight", "white mould" e "halo blight" que podem também causar uma perda significativa na produçao, mas elas tendem a confinhar-se a específicas condições ambientais.

Para os pequenos agricultores da África Sub-Sahariana, a planta hospedeira resistente mantêm apenas médias razoáveis de controlo da doença. Neste momento, muitos agricultores semeiam tarde a cultura apenas para evitar a estação muito chuvosa; melhorando o nível de resistência varietal às doenças e adicionando o benefício de permitir que haja um maior período de sementeira. Os cientistas da Investigação Agrícola Nacional (NARs) com grande dedicação tem concentrado os seus esforços em promover o feijao vulgar na África Oriental e Austral incluindo a avaliação de ambos os germoplasmas: introduzido e local. Existe a evidência viva de um substancial benefício que pode ser ganho directamente pela introdução de uma grande diversidade genética, de raças melhoradas disponíveis especialmente no Centro Internacional de Agricultura Tropical (CIAT) em Colômbia (Allen & Smithson, 1991) e de matérias de outros centros com programas de melhoramento bem estabelecidos. Esta foi a estratégia que permitiu que muitos países africanos libertassem novas variedades muito produtivas depois de poucos anos. Nenhuma troca de germoplasma pode evitar completamente o risco. O Este de África, em particular, tem uma longa tradição e um recorde em trabalhar coordenação com o Serviço de Quarentena com vista a evitar a introdução acidental de patógeno estrangeiro e as suas raças dentro da região (Sheffield, 1968). A

maioria das doenças transmitidas pela semente do feijão vulgar ocorrem na maioria das regiões onde a cultura é cultivada. Em alguns casos, o surgimento das doenças e pragas podem ser atribuídas não à introdução casual de um agente mas ao nível de susceptibilidade de uma linhagem indígena quando é confrontada com uma nova introdução (Allen, 1993). Um exemplo em África é a mosca do feijão (*Ophiomyia* spp.) que é uma praga antiga das novas introduções de *Phaseolus vulgaris*. Novas doenças também chegar devido a mudança das práticas culturais ou o uso excessivo de terras marginais. Por exemplo, a podridão da raiz, causada por um complexo de fungos que inclui as espécies de *Pythium*, está aumentando devido ao cultivo intensivo da cultura (Buruchura, 1992c). Enquanto que a mancha cinzenta pode ser séria em solos não férteis submetidos a "stress" hídrico.

E obviamente útil conhecer quais os patógenos estão presentes numa determinada área antes de se fazer uma substancial introdução na região. A lista de guia internacional fornece uma base útil para formular os regulamentos que guiam as importações. Isso pode ser conseguido fazendo uma listagem dos patógenos que são conhecidos e a sua presença nessa região. Nesse contexto, no feijão vulgar os pontos de comparação entre África e América Latina estão disponíveis. Embora esses patógenos encontrem-se nos dois continentes existe algumas exceções: o fungo "scab" (*Elsinoë phaseoli* Jenk.) não é conhecido no *P. vulgaris* da América, entretanto está largamente espalhado em África. Ao contrário, dos fungos como: *Pseudocercosporella albida* (Matta & Bell.) Deighton, *Chaetoseptoria wellmanii* Stev., *Entyloma* sp., *Cercospora vanderystii* Henn. e *Cercospora castellanii* Matta & Bell. ainda não foram identificadas em África. As vírus que ainda não estão identificadas na lista das infecções naturais do feijão em África incluem o mosaico comum, mosaico rugoso, vírose da casca do feijão, "quail pea mosaic", o mosaico clorótico do feijão caupi, o mosaico doirado do feijão, "bean dwarf" e outros gêneros de vírus que não são transmitidos pela semente (Galvez & Morales, 1989; Morales & Gamez, 1989). Entre as bactérias, existem algumas que são diferentes como o caso estranho do *Curtobacterium flaccumfaciens* (Hedges) Collins & Jones que precisa de mais comentários. O *C. flaccumfaciens* é conhecida, a vários anos, como sendo o causador da mancha bacteriana do feijão vulgar em Nebraska e outras regiões dos EUA (Mohan & Hagedorn, 1989). Devido à transmissão da bactéria pela semente, a sua distribuição geográfica é aparentemente limitada em ambos os continentes: em América e África; *C. flaccumfaciens* ficou documentada na categoria de alto risco pela Quarentena (Neergaard, 1977). Segundo a opinião dos oficiais de quarentena em África este patógeno é uma "batata quente" dentro do grupo dos patógenos de feijão. Devido ao facto de agora se conhecer a natureza e a importância da bactéria é importante explicar o seu estado. Existem evidências limitadas para provar que a mancha bacteriana é economicamente importante, mesmo nos EUA (J.R. Steadman & M.J. Silbernagel, com. pessoal, 1988). *C. flaccumfaciens* é também um comum habitante na América Latina: é relativamente fácil identificar a bactéria mas a tarefa de dela como patógeno é mais difícil de se provar (M.A. Pastor-Corrales, com. pessoal, 1987). A sua presença mesmo nos outros legumes incluindo o gênero *Zornia* mostra que a bactéria parece não ser necessariamente perigosa para a cultura do feijão. Estes factores juntos levam a questões de competência epidemiológica do *C. flaccumfaciens* que parece ter sido registrado com uma excessiva prominência entre os patógenos do feijão vulgar.

Com o desenvolvimento recente em África dos programas nacionais de melhoramento e as redes regionais de trabalho para a troca do germoplasma, já não são só os agricultores que têm interesse em transportar semente de feijão através das fronteiras nacionais, mas também os investigadores. A falta do conhecimento sobre os riscos envolvidos por vezes torna difícil para os

países colaborarem eficientemente na investigacao do melhoramento da cultura. No encontro dos directores africanos de Investigacao Agronómica que teve lugar em Nairobi em Junho de 1992, foi reconhecido que "os regulamentos da quarentena das plantas enquanto visa a prevencao da introducao accidental das doenças que nao estao presentes num dado país, usualmente aplicando o conhecimento da distribuicao das doenças que a muitos anos ainda nao foi aplicado ... Foi concordado que o CIAT, em colaboracao com patológitas da vários programas nacionais de investigacao agronómica, deveriam preparar um mapa das doenças do feijao para os directores nacionais poderem estudar a informacao que será utilizada pelos servicos de quarentena". Este encontro foi precedido pela reuniao do Grupo Pan-africano que Trabalha com os patogenos de fungos do feijao que teve lugar em Thika, Quénia em Maio de 1992, quando o CIAT recebeu a incumbencia de compilar a lista de referencia sobre os patogenos do feijao como um primeiro passo para desenhar um mapa de distribuicao das doenças conhecidas.

A presente lista tem os seguintes objectivos:

- fornecer uma base científica para a preparacao do mapa de distribuicao das doenças através da qual a quarentena tem a legislacao que governa o movimento das sementes do feijao dentro e fora de África;
- fornecer aos oficiais da quarentena das plantas informacao necessária do nível e o tipo de risco associado a cada introducao individual assim como orientar as decisões apropriadas a cada um dos casos;
- fornecer um conjunto de prioridades para os melhoradores das culturas através do acesso da recente informacao recolhida da investigacao regional dos patogenos e as suas varientes; e
- fornecer uma base de seleccao e "screening" de resistencia as doenças.

A revisao da lista inclue 20 países da África Oriental, Central e Austral e a ilhas vizinhas onde a cultura do feijao vulgar é regionais de feijao em África, já catalizou o desenvolvimento da investigacao e treinamento na África Ocidental e Oriental, incluindo na lista os membros da rede de trabalho. Esta lista omite o Botswana e a Namíbia onde o feijao vulgar é uma cultura insignificante e portanto, as doenças ainda não foram reportadas. A revisao bibliográfica foi extensiva mas provavelmente não exaustiva. Os registos dos patogenos e das doenças conseguem cobrir muitos registos antigos bem como os recentes, para dar uma perspectiva histórica da distribuicao das mesmas. Foi feito um uso substancial das listas territoriais de consulta, nos sítios onde elas existem. Um registo suplementar de uma fonte disponível no Instituto Internacional Micológico (IMI) e o "CAB International" onde as colecções estão numeradas IMI número. A nomenclatura do patógeno segue IMI como mostra o Holliday (1980). Os registos estão confinados a uma única espécie hospedeira *Phaseolus vulgaris*. Em vários casos, um patógeno (ou raça) particular pode estar presente num determinado país e a planta hospedeira no outro indicando assim, a importância da distinção entre a distribuição geográfica da doença e a importância da distribuição casual dum casual patógeno; eles não são sinônimos, como está enfatizado nas outras partes do mundo (Allen, 1983) e ilustrado na distribuição das raças da mancha bateriana (Tabela 1). Durante a recente revisão, vários patógenos tinham sido compilados nas plantas leguminosas hospedeiras e outros no *P. vulgaris* e por sua vez, estes patógenos já tinham sido compilados noutras partes do mundo e publicados (Allen, 1991). Obviamente que fica uma fenda no conhecimento dos patógenos de feijao vulgar em África, e as consultas precisam de ser completadas. Com a investigação novos registros serão melhorados. Estes registros funcionam na base regional e publicados num suplemento anual num jornal apropriado, como por exemplo o Buletim da Proteção de Plantas da FAO.

AGRADECIMENTOS

Esta publicacao, sem dúvida que deveu-se a ajuda substancial de várias pessoas e agencias. No início, o director e colegas do Instituto Internacional de Micología deram-me o cesso aos registros nao publicados e deram-me o guiao da nomenclatura do patogenos, suplementado pela constante referencia que eu faço a Paul Holliday: Doencas fungosas das culturas trópicais. O Sr Peter Filby deu-me uma grandiosa assistencia em termos de acesso a importante Coleccao de Buttress, que ainda nao foi muito bem utilizada, da biblioteca científica de Cambridge, onde a juda do Dr S. Dale é também reconhecida. Os Drs C.Y.L. Schotman e R. Ikin generosamente deram-me o acesso ao sistema global de informacao da quarentena das plantas da FAO.

O rescunho de uma lista dos países tinha sido circulada a todos os fitopatologistas e responsáveis do programa de feijao que amavelmente fizeram comentários. Quero particularmente agradecer a todos que me adjudaram e em alguns casos inclusivamente deram-me o acesso a informacao ainda nao publicada:

Dr J. Arias, Dr R. Buruchara, Dr N. Govinden, Ato Habtu Assefa, Dr H.C. Haciwa, Dr J. Kannaiyan, Mr J. Kayitare, Dr A. Liebenberg, Dr R.J.M. Melis, Dr O.Z. Mukoko, Mr N. Ntahimpera, Mr M.E. Omunyin, Dr A.F. Opio, Dr D. Perreaux, Dr A. Phillips, Mr Mukishi Pyndji, Mr A.H. Ramos, Prof Y.P. Rao, Mr S. Saumtally, Dr V. Schmit, Mrs T. Sengooba, Dr J.B. Smithson, Dr N. Spence, Prof J.M. Teri, Dr D.M. Teverson, Dr P. Trutmann e Dr H.J. Vetten.

Dr B.C. Sutton que providenciou o mais valioso guia durante o estágio final de producao.

Neste momento poderia dizer que a responsabilidade do conteúdo técnico é inteiramente minha: os meus comentário nao sao necessarios para a reflexao oficial das opiniões do CABI ou CIAT que em conjunto estão publicados. O Dr R.A. Kirkby generosamente encorajou a terminar esta tarefa que foi totalmente financiada pela Agencia de Desenvolvimento Internacional do Canada (CIDA) para o Centro Internacional de Agricultura Tropical que tem Projectos Desenvolvimento da Rede de Trabalho de Investigacão em África.

Finalmente, gostaria de adicionar os meus profundos agradecimentos a minha esposa Leonora, que ajudou-me directamente no longo período de compilacão e correção do livro.

DJA
Cambridge, January 1994

ANNOTATED TERRITORIAL LISTS

ANGOLA

Alternaria leaf spot: *Alternaria tenuissima* (Kunze) Wiltshire.

Local. Huambo.

Serafim & Serafim (1968).

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.

Widespread.

Serafim & Serafim (1968), as *Isariopsis griseola* Sacc.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.

Widespread and important.

Serafim & Serafim (1968).

Ascochyta blight (leaf spot): *Ascochyta phaseolorum* Sacc.

Widespread.

Serafim & Serafim (1968).

Cercospora leaf spot: *Pseudocercospora cruenta* (Sacc.) Deighton.

Local. Huambo and Uige.

Serafim & Serafim (1968), as *Cercospora cruenta* Sacc.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

Chianga, Huambo, Nov. 1988.

Allen (1988).

Fusarium root rot: *Fusarium solani* (Mart.) Sacc. f.sp. *phaseoli* (Burkh.) W.C. Snyder & E.C. Hansen.

Local, minor importance. Bie and Huambo.

Serafim & Serafim (1968).

Mosaic: ? virus.

Camarada (1989).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Widespread and important.

Serafim & Serafim (1968).

Southern blight (stem rot): *Corticium rolfsii* Curzi.

Widespread, minor importance.

Serafim & Serafim (1968), as *Sclerotium rolfsii* Sacc.

Yeast spot (Stigmatomycosis): *Nematospora* sp.

Local, minor importance, associated with the insect *Nezara viridula*. Huambo.

Serafim & Serafim (1968).

BURUNDI

Alternaria leaf spot: *Alternaria alternata* (Fr.) Keissler.
P. Trutmann, unpubl.

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Buyckx (1962), ISABU (1983), as *Isariopsis griseola* Sacc.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
INEAC (1958b), Buyckx (1962). More important at higher altitudes, as at Gisozi (Perreux, 1986).

Ascochyta blight: *Phoma exigua* Desm. var. *diversispora* (Bubák) Boerema, and *Ascochyta phaseolorum* Sacc.
Buyckx (1962), Autrique (1981), Perreux (1986).
Positive identification as var. *diversispora*, coll. at Murongwe, D.J. Allen 1985 (Gerlagh, 1987).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.
Autrique & Perreux (1989).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
Autrique (1981), Munimbazi & Perreux (1986).

Common mosaic: Bean common mosaic virus.
Goethals (1986).

Pathogenicity groups I, III, IVb, Va and VIa. Necrotic strains of the NL3 and NL8 type predominant (Spence, 1992).

Damping off: *Thanatephorus cucumeris* (Frank) Donk.
Buyckx (1962), as *Rhizoctonia solani* Kuhn.

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton. Some damage to beans, especially at altitudes between 1200-1400 m, as in Moso.
Dieudonne (1980), Perreux (1986), as *Ramularia phaseoli* (Drummond) Deighton.

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
Race 3 (IHR 1377A, J.D. Taylor 1985; IHR 1567A and 1568A, D. M. Teverson 1986) and race 4 (IHR 1385A and 1386A, D. Perreux 1985; IHR 2240A, D.M. Teverson 1988; IHR 2274A, A. Autrique).
Teverson (1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Mosaic: Cucumber mosaic virus.
M. Goethals, unpubl.

Mosaic: Bean strains of blackeye cowpea mosaic virus.
Imbo Plains (H.J. Vetten, pers. comm. 1993).

Pythium root rot: *Pythium* spp.
ISABU (1990).

Root rot: associated with a complex of fungi, particularly *Fusarium solani* (Mart.) Sacc. and *Rhizoctonia* sp.
CIAT (1988).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Moso.

INEAC (1958c), Buyckx (1962).

Web blight: *Thanatephorus cucumeris* (Frank) Donk.

Autrique (1981), Perreux *et al.* (1986).

White mould: *Sclerotinia sclerotiorum* (Lib.) de Bary.

Buyckx (1962).

Southern blight (root rot): *Corticium rolfsii* Curzi.

ISABU (1983), as *Sclerotium rolfsii* Sacc.

ETHIOPIA

Alternaria leaf spot: *Alternaria solani* Sorauer.

IMI 207093.

Alternaria tenuissima (Kunze) Wiltshire.

IMI 207094.

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.

IMI 88330, 116687.

Kaffa, Eritrea, Shewa (Stewart & Dagnatchew, 1967), as *Cercospora columnaris* Ellis & Everh.

Didessa (Allen *et al.*, 1976), Awassa (Habtu, 1987).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.

IMI 116596.

Shewa (Stewart & Dagnatchew, 1967).

Bako (Allen *et al.*, 1976).

Didessa (Habtu, 1987).

Ascochyta blight: *Ascochyta phaseolorum* Sacc.

IMI 116509.

Shewa (Stewart & Dagnatchew, 1967).

Bako, Didessa (Habtu, 1987).

Ashy stem blight: *Macrophomina phaseolina* (Tassi) Goid.

Melkassa and Kobbo (Habtu, 1987).

Associated fungi: *Cladosporium oxysporum* Berk. & M.A. Curtis, IMI 207094.

Cladosporium sp., IMI 225599b.

Cercospora leaf spot: *Pseudocercospora cruenta* (Sacc.) Deighton.

Kaffa (Stewart & Dagnatchew, 1967), as *Cercospora cruenta* Sacc.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

IMI B 659, 6930, IMI 136927.

Begemder, Harar, Shewa (Stewart & Dagnatchew, 1967).
Nazreth, Angar Gutin (Allen *et al.*, 1976).

Common mosaic: Bean common mosaic virus.
Bos (1974), Agranovsky (1985).

Recent work has shown that necrotic 'A' serotype strains of BCMV are apparently rare or even absent from Ethiopia, in contrast to elsewhere in eastern and southern Africa.

Pathogenicity groups I and IVb present (Spence & Walkey, 1991; Vetten & Allen, 1991) and possibly also some novel strains (Spence, 1992).

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.
IMI 116536, 207095.

Shewa, Kaffa (Stewart & Dagnatchew, 1967), as *Ramularia phaseoli* (Drummond) Deighton.
Didessa (Allen *et al.*, 1976), Awassa (Habtu, 1987).

From bean plant: unidentified nepovirus.
Alemaya (H.J. Vetten, pers. comm. 1993).

Fusarium root rot: *Fusarium solani* (Mart.) Sacc.
Habtu (1987).

Fusarium wilt: *Fusarium oxysporum* Schlecht.
Habtu (1987).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
IMI B 660.

Shewa, Harar, Sidamo (Stewart & Dagnatchew, 1967). Race 4 (IHR 1448C, Habtu Assefa 1985) and race 6 (IHR 1448A, Habtu Assefa 1985; IHR 1715A, D.J. Allen 1986), Teverson (1991) as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Leaf spot: *Phoma exigua* Desm.
IMI 225546.
Kidane (1982).

Leaf spot: *Phoma phaseolina* Pass.
IMI 225536.
Kidane (1982).

Mosaic: Peanut mottle virus.
Lake Shalla.
Spence (1992).

Mosaic: Unidentified potyvirus with serological and biological affinities to soybean mosaic virus.
Widespread in the Rift Valley (H.J. Vetten, pers. comm. 1993).

Mycosphaerella leaf spot: *Mycosphaerella pinodes* (Berk. & Bloxam) Vestergr.
IMI 225500b.
Kidane (1982).

On leaves: *Cercosporidium* sp.
Kidane (1982).

Periconia leaf spot: *Periconia byssoides* Pers.
IMI 225599c.
Kidane (1982).

Phyllosticta leaf spot: *Phyllosticta phaseolina* Sacc. and *Phyllosticta* sp.
IMI 88301.
Kaffa, Shewa (Stewart & Dagnatchew, 1967).

Pod spot: *Phoma sorghina* (Sacc.) Boerema, Dorenb. & van Kest.
IMI 215673.
Kidane (1982).

Rhizoctonia root rot: *Rhizoctonia solani* Kuhn.
Widespread but of minor importance.
Habtu (1987).

Rust: *Uromyces appendiculatus* (Pers.) Unger.
IMI 88610.
Uredinal and telial states. Widespread.
Stewart & Dagnatchew (1967), as *U. phaseoli* var. *typica* Arthur. Epidemiology studied by Habtu (1994).

Southern blight (root rot): *Corticium rolfsii* Curzi.
Kaffa (Stewart & Dagnatchew, 1967), as *Sclerotium rolfsii* Sacc.

Stem anthracnose: *Colletotrichum truncatum* (Schwein.) Andrus & Moore.
Kaffa (Stewart & Dagnatchew, 1967).

Web blight: *Thanatephorus cucumeris* (Frank) Donk.
Pawe (Smithson, 1986).

White mould (stem rot): *Sclerotinia sclerotiorum* (Lib.) de Bary.
Habtu Assefa, pers. comm. 1985.

Yellow mosaic: Bean yellow mosaic virus.
Jimma (H.J. Vetten, pers. comm. 1993).

KENYA

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
IMI 91195b, 44232, 44233, 276220a, 91195b.
McDonald (1936), as *Isariopsis griseola* Sacc.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
IMI 57460.
McDonald (1936).

Ascochyta blight (Ascochyta leaf spot, phoma blight, black node disease): conventionally referred to *Ascochyta phaseolorum* Sacc., which is often recorded (IMI 44109, 184643a). Is now regarded as a synonym of the weak parasite *Phoma exigua* var. *exigua* Desm. (Boerema, 1972), which is often found on beans in Kenya (IMI 184643, 161806d, 274551; Gatumbi, 1986). However, *P. exigua* var. *diversispora* (Bub.) Boerema is endemic to eastern Africa, including Kenya, and appears to be the taxon of greatest economic importance (Boerema *et al.*, 1981; Stoetzer & Waite, 1984). Records of *Ascochyta holtshauseri* Sacc. in Kenya (Stoetzer & Waite, 1984) are now referable to *Stagonosporopsis hortensis* (Sacc. & Malbr.) Petr.

Ashy stem blight (charcoal rot): *Macrophomina phaseolina* (Tassi) Goid.

IMI 62566, 114782.

Nattrass (1961), as *M. phaseoli* (Maubl.) Ashby.

Associated fungi: *Alternaria cucumerina* (Ellis & Everh.) J.E. Elliot, IMI 206021.

Alternaria sp., IMI 213685.

Gatumbi (1986).

Chaetomium spirale Zopf, IMI 170327.

Gatumbi (1986).

Cladosporium sp., IMI 161806g.

Epicoccum purpurascens Ehrenb. ex Schlecht.

Under *E. nigrum* Link, IMI 161806a.

Fusarium semitectum Berk. & Rav., IMI 158641.

Gatumbi (1986).

Penicillium cyclopium Westling, IMI 68119.

Nattrass (1961).

Bacterial wilt: *Curtobacterium* (=*Corynebacterium*) *flaccumfaciens* (Hedges) Collins & Jones.

A few isolated records from the period 1968-1973. Presumed eradicated (Hubbeling, 1973; A.H. Ramos, pers. comm. 1988).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.

Kaiser & Ramos (1980).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

Very common (Hubbeling, 1973; Mukunya *et al.*, 1981).

Common mosaic, black root: Bean common mosaic virus.

'Mosaic' attributed to virus by McDonald (1936). BCMV positively identified by Kulkarni (1973). See also van Rheenen & Muigai (1984). Strain variation first studied by Omunyin (1979) and presence of both serotypes A and B confirmed by Vetten & Allen (1991). Pathogenicity groups Va and VIa, the former apparently confined to western Kenya, near Kisii (Spence, 1992).

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.

IMI 91195a, 151003.

Ondieki (1973), as *Ramularia phaseoli* (Drummond) Deighton.

Fusarium root rot: *Fusarium solani* (Mart.) Sacc. f.sp. *phaseoli* (Burkh.) W.C. Snyder & E.C. Hansen.

Important (Hubbeling, 1973; Njugunah *et al.*, 1980; Mutitu *et al.*, 1983).

Fusarium wilt, yellows: *Fusarium oxysporum* Schlecht. f.sp. *phaseoli* J.B. Kendr. & W.C. Snyder.
Mutitu *et al.* (1988).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

McDonald (1936), as *Bacterium medicaginis* var. *phaseolicola* (Burkh.) Link & Hall; Mukunya *et al.* (1981).
Studies on pathogenic variation have shown the presence of at least three races: race 4 (IHR 1356A, G.K. Kinyua 1985), race 6 (IHR 1435A, J.D. Taylor 1985; IHR 2253A, 2254A, 2254C, D.M. Teverson 1988) and race 7 (IHR 1354A, D.J. Allen 1985).

Kinyua *et al.* (1981), Teverson (1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Leptosphaerulina leaf spot: *Leptosphaerulina trifolii* (Rostr.) Petr.
IMI 161806b, 124259.

Phyllosticta leaf spot: *Phyllosticta phaseolina* Sacc.
IMI 161806c.

Powdery mildew: *Erysiphe polygoni* DC. ex St.-Am.

Moderately important (Hubbeling, 1973; Njugunah *et al.*, 1980).

Pythium root rot: *Pythium* sp.

Hubbeling (1973), Njugunah *et al.* (1980).

Recently found severe in western districts, associated with heavy rainfall (Buruchara, 1992a).

Rhizoctonia root rot: *Rhizoctonia solani* Kuhn.

Widespread (Hubbeling, 1973; Njugunah *et al.*, 1980).

Rust: *Uromyces appendiculatus* (Pers.) Unger, and apparently also *U. viciae-fabae* (Pers.) Schroet.

U. appendiculatus recorded by McDonald (1936) and pathogenic variation studied by Howland & Macartney (1966). *U. viciae-fabae* recorded by Nattrass (1961) but this record should be regarded with suspicion.

Scab: *Elsinoë phaseoli* Jenk.

IMI 21136.

Mutitu (1979), Gatumbi (1986).

Southern blight: *Sclerotium rolfsii* Sacc.

Moderately important (Hubbeling, 1973; Njugunah *et al.*, 1980).

Tobacco mosaic: Tobacco mosaic virus.

One record (Hollings *et al.*, 1981).

White mould: *Sclerotinia sclerotiorum* (Lib.) de Bary.

IMI 141593, 141594, 142922.

Ondieki (1973).

Yeast spot: *Nematospora coryli* Peglion.

Nattrass (1961).

Yellow mosaic: Bean yellow mosaic virus.

An early record of yellow mosaic (Wallace, 1941) was presumed to have been BYMV (Zaumeyer & Thomas,

1957). Recorded from Embu and Njoro (Hubbeling, 1973); its presence in Njoro has been confirmed recently (Vetten & Allen, 1991).

LESOTHO

Ashy stem blight (Charcoal rot): *Macrophomina phaseolina* (Tassi) Goid.

IMI 322519, 322522, 322523, 322524, 322525.

Maseru, Mafeteng.

Smithson (1988).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.

IHR 2673A-D, 2675C, 2676C, 2677C, 2682C, J.D. Taylor 1990.

Teverson (1991), D.M. Teverson, pers. comm. 1993.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

Widespread (Smithson, 1988).

Common mosaic: Bean common mosaic virus.

Pathogenicity group VIa (Spence & Walkey, 1991).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

Maseru, Mafeteng and Machache.

Race 1 (IHR 1967A, J.B. Smithson 1988; IHR 2662A, 2675A, 2676A, 2681A, 2682A, J.D. Taylor 1990),

race 2 (IHR 2671A, J.D. Taylor 1990), race 6 (IHR 1965A, J.B. Smithson 1988; IHR 2367A, 2368A, D.J.

Allen 1989; IHR 2654A, 2673A, J.D. Taylor 1990) and race 8 (IHR 2654C, 2656A, 2656B, 2656C *et seq.*,

J.D. Taylor 1990), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Teverson (1991).

Root rot and wilt: *Neocosmospora vasinfecta* E.F. Sm. var. *vasinfecta*

IMI 322520.

Maseru (Smithson, 1988).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Leribe, Maseru and Teyateyaneng (Allen & Smithson, 1989).

MADAGASCAR

Alternaria leaf spot: *Alternaria solani* Sorauer and, to a lesser extent, also *A. longipes* (Ellis & Everh.) Mason. *A. solani* is found all over the island and almost all bean cultivars on which both leaves and pods are attacked (Dadant *et al.*, 1961).

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.

Found around Tananarive, Bealanana and Lac Alaotra (Bouriquet, 1946). Common but regarded of minor importance (Dadant *et al.*, 1961).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.

Found first in February 1937 (Bouriquet, 1946).

Common and fairly important (Dadant *et al.*, 1961).

Ascochyta blight: presumably *Phoma exigua* Desm. var. *diversispora* (Bubák) Boerema Buruchara (1992b).

Bacterial wilt: *Burkholderia solanacearum* (E.F. Sm.) Yabuuchi *et al.*

Widely distributed and damaging, 20-50 per cent incidence.

Dadant *et al.* (1961), as *Pseudomonas solanacearum* (E.F. Sm.) E.F. Sm.

Cercospora leaf spot: *Pseudocercospora cruenta* (Sacc.) Deighton.

Damage apparently insignificant (Bouriquet, 1946), as *Cercospora cruenta* Sacc.

Frequent but unimportant (Dadant *et al.*, 1961).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

Found near Lac Itasy in 1932 (Bouriquet, 1946), as *Bacterium phaseoli* Smith.

Common mosaic: presumed to be bean common mosaic virus.

Rasolofo-Razafindramamba & Ralilarison (1976).

Fusarium wilt: *Fusarium* sp.

Uncommon and of little importance (Dadant *et al.*, 1961).

Mosaic: Peanut mottle virus.

H.J. Vetten, pers. comm. 1993.

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Found first in 1929 but thought then to have been present for a long time (Bouriquet, 1946).

Widespread; important on very susceptible cultivars (Dadant *et al.*, 1961).

Sclerotium root rot (southern blight): *Corticium rolfsii* Curzi.

Found in 1934 on the East coast at Fenerive (Bouriquet, 1946), as *Sclerotium rolfsii* Sacc. Causes important damage under conditions of poor drainage or heavy shade (Dadant *et al.*, 1961).

Web blight: *Thanatephorus cucumeris* (Frank) Donk.

Frequent; severe under humid conditions (Dadant *et al.*, 1961), as *Corticium solani* (Prill. & Delacr.) Bourd. & Galz.

MALAWI

Alternaria leaf spot: *Alternaria* sp.

Sometimes causes severe defoliation.

Peregrine & Siddiqi (1972).

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.

IMI 156668b, 164268, 89206, 130925, 137407.

Peregrine & Siddiqi (1972).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.

IMI 114415, 124356a, 124357a, 125115.

The most serious disease in Malawi (Peregrine, 1971; Peregrine & Siddiqi, 1972).

Pathogenic variation studied by Ayonoado (1974) who found that the gamma race group predominated among 16 isolates obtained from three sites in Malawi. Three new races were identified.

Ascochyta leaf spot: *Ascochyta* sp. and *A. phaseolorum* Sacc.

IMI 92151.

Peregrine & Siddiqi (1972).

Associated fungi: *Epicoccum nigrum* Link, IMI 124357b.

Peregrine & Siddiqi (1972).

Cercospora leaf spot: *Cercospora* sp. and *Cercospora canescens* Ellis & G. Mart.

IMI 164267, 119349d.

Peregrine & Siddiqi (1972).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

IMI B 3340, B 3643.

Widespread (Peregrine & Siddiqi, 1972), as *X. phaseoli* (E.F. Sm.) Dowson, and includes isolates referable to fuscous blight.

Common mosaic: Bean common mosaic virus.

Peregrine & Siddiqi (1972). Necrosis-inducing strain identified from distorted apical leaves from Bunda, Feb. 1989 (H.J. Vetten, unpubl. 1989). Pathogenicity groups IVb and VIa (Spence, 1992).

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.

IMI 156668, IMI 155355.

Serious, Central Region (Peregrine & Siddiqi, 1972), as *Ramularia phaseoli* (Drummond) Deighton.

Fusarium root rot: *Fusarium solani* (Mart.) Sacc.

Mughogho, 1970 in Allen (1975a).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

IMI B 3376.

Peregrine & Siddiqi (1972), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.* Pathogenic variation of halo blight in Malawi studied by Msuku (1985). Subsequent studies have identified race 2 (IHR 2710A, J.D. Taylor 1990), race 5 (IHR 2708A, J.D. Taylor 1990) and race 9 (IHR 2709A, J.D. Taylor 1990). Teverson (1991).

Mild mottle: cowpea mild mottle virus.

Bunda, nr Lilongwe.

Vetten & Allen (1991).

Mycosphaerella leaf spot: *Mycosphaerella pinodes* (Berk. & Bloxam) Vestergr.

IMI 86575b.

Peregrine & Siddiqi (1972).

On roots: *Rhizopus stolonifer* (Ehrenb.) Lind.

IMI 159467b.

Peregrine & Siddiqi (1972).

Periconia leaf spot: *Periconia byssoides* Pers.

IMI 86575a.

Peregrine & Siddiqi (1972).

Phyllosticta leaf spot: *Phyllosticta* sp.

Peregrine & Siddiqi (1972).

Root rot: *Gibberella fujikuroi* (Sawada) Ito (anamorph: *Fusarium moniliforme* J. Sheld.).

IMI 159467a.

Peregrine & Siddiqi (1972).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

IMI 155355c.

A serious disease with heavy yield losses in dry season irrigated crops; particularly white-seeded cultivars.

Peregrine & Siddiqi (1972).

Scab: *Elsinoë phaseoli* Jenk.

IMI 81050, 164269.

Peregrine & Siddiqi (1972).

Southern blight: *Corticium rolfsii* Curzi.

Not serious.

Peregrine & Siddiqi (1972).

Stem anthracnose: *Colletotrichum truncatum* (Schwein.) Andrus & Moore.

IMI 155356b.

Peregrine & Siddiqi (1972).

Web blight, collar rot and stem blight: *Thanatephorus cucumeris* (Frank) Donk.

IMI 124773.

Peregrine & Siddiqi (1972), as *Corticium solani* (Prill. & Delacr.) Bourd. & Galz.

Yellow mosaic: Bean yellow mosaic virus.

Peregrine & Siddiqi (1972).

Dubious, not confirmed during recent surveys (Vetten & Allen, 1991).

MAURITIUS

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
E.F.S. Shepherd, 1937 (Wiehe, 1948), as *Isariopsis griseola* Sacc.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
E.F.S. Shepherd, 1937 (Wiehe, 1948).

Bacterial wilt: *Burkholderia solanacearum* (E.F. Sm.) Yabuuchi *et al.*
IMI B 1728, 1729.

Beau Bassin, 1963 (Orieux & Felix, 1968), as *Pseudomonas solanacearum* (E.F. Sm.) E.F. Sm.

Also *Curtobacterium flaccumfaciens* (Hedges) Collins & Jones, as *Corynebacterium flaccumfaciens* (Hedges) Dowson, the latter reported for the first time from Beau Bassin and Black River in 1963. In 1964, found at Mapou. In the summer of 1965, *C. flaccumfaciens* was reported as severe in a bean crop of Local Red, at an incidence of 10-36% (Anon., 1965-67).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.
Autrey & Saumtally (1990).

Cercospora leaf spot: *Cercospora canescens* Ellis & G. Mart.
IMI 76665.
Reduit, 1959 (Orieux & Felix, 1968).

Charcoal rot (ashy stem blight): *Macrophomina phaseolina* (Tassi) Goid.
IMI 90236, 90237.
Richelieu, 1961 (Orieux & Felix, 1968).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
First recorded 1981. Its fuscous variant found more recently.
Autrey & Saumtally (1990).

Fusarium root rot: *Fusarium solani* (Mart.) Sacc.
Rojoa (1989).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
1950 (Orieux & Felix, 1968), as *P. phaseolicola* (Burkh.) Dowson.
Race 2 (IHR 1502A, 1986), Teverson (1991).

Leaf spot: *Pleospora* sp.
IMI 76664b.
Orieux & Felix (1968).

Mosaic: ? Bean common mosaic virus.
Wiehe (1948), Anon. (1986).

Phyllosticta leaf spot: *Phyllosticta phaseolina* Sacc.
IMI 76664a.
Reduit, 1959 (Orieux & Felix, 1968).

Powdery mildew: *Erysiphe polygoni* DC. ex St-Am.
F.A. Stockdale, 1915 (Wiehe, 1948).

Rhizoctonia root rot (as wilt): *Thanatephorus cucumeris* (Frank) Donk.
Orieux & Felix (1968), as *Corticium solani* (Prill. & Delacr.) Bourd. & Galz.

Rust: *Uromyces appendiculatus* (Pers.) Unger.
E.F.S. Shepherd, 1937 (Wiehe, 1948).

Southern blight (as root rot): *Corticium rolfsii* Curzi.
1958 (Orieux & Felix, 1968).

White mould (as pod rot): *Sclerotinia sclerotiorum* (Lib.) de Bary.
Reduit, G. Orian, 1945 (Wiehe, 1948).

Yellow mosaic: ? Bean yellow mosaic virus.
Beau Plan (Anon., 1986).

MOZAMBIQUE

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Plumb-Dhindsa & Mondjane (1984).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
Plumb-Dhindsa & Mondjane (1984).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
Plumb-Dhindsa & Mondjane (1984).

Damping off: causal fungi apparently undetermined.
Common in Lichinga.
J. Arias, pers. comm. 1993.

Fusarium root rot: *Fusarium* sp.
Plumb-Dhindsa & Mondjane (1984).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
Plumb-Dhindsa & Mondjane (1984), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*
Races not yet identified.

Leaf spot: *Phoma exigua* Desm.
Plumb-Dhindsa & Mondjane (1984).

Mosaic: Complex of viruses.
Plumb-Dhindsa & Mondjane (1984).

Rust: *Uromyces appendiculatus* (Pers.) Unger.
Plumb-Dhindsa & Mondjane (1984).

Seed decay: Complex of fungi.
Plumb-Dhindsa & Mondjane (1984).

Stunt: associated with cowpea mild mottle virus
Umbeluzi, March 1989 (Vetten & Allen, 1991).

Web blight: *Thanatephorus cucumeris* (Frank) Donk (sclerotial state *Rhizoctonia solani* Kuhn).
Plumb-Dhindsa & Mondjane (1984).

RWANDA

Alternaria leaf spot: *Alternaria alternata* (Fr.) Keissl.
P. Trutmann, unpubl.

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Nyabyenda *et al.* (1981), as *Isariopsis griseola* Sacc.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
Goeteyn (1975), Nyabyenda *et al.* (1981), Kamanzu (1982).
Pathogenic variation studied by Nkezabera (1987).

Ascochyta blight: *Phoma exigua* Desm. var. *diversispora* (Bubák) Boerema, and *Ascochyta phaseolorum* Sacc.
Durnez (1983), Price & Cishahayo (1986), Munyemana (1987).
Positive identification as var. *diversispora*, coll. P. Trutmann 1986 (Gerlagh, 1987).

Ashy stem blight: *Macrophomina phaseolina* (Tassi) Goid.
P. Trutmann & W. Graf, unpubl. 1987, Rusuku (1991), Buruchara & Rusuku (1992).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.
IHR 1378A-D, J.D. Taylor 1985; IHR 1912A-C, 1913A-D, 1915A-D, 1917A-D, 1918A-D, 1922A-D,
1928A-D, J.D. Taylor 1987.
Teverson (1991), D.M. Teverson, pers. comm. 1993.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
Durnez (1983).

Common mosaic, black root: Bean common mosaic virus.
Pathogenicity groups III, IVb and VIa (Spence, 1992).

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.
Durnez (1983).

Fusarium root rot: *Fusarium solani* (Mart.) Sacc.
P. Trutmann, unpubl. 1987, Rusuku (1991), Buruchara & Rusuku (1992).

Fusarium wilt: *Fusarium oxysporum* Schlecht. f.sp. *phaseoli* J.B. Kendr. & W.C. Snyder.
Butare and Gikongoro (Rusuku, 1991; Buruchara, 1992c; Buruchara & Rusuku, 1992).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
ISAR (1985), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*
Race 1 (IHR 1500A, P. Trutmann 1985), race 2 (IHR 1612A, D.M. Teverson 1986) and race 4 (IHR 1302A,
J.H.C. Davis 1984; IHR 1348A, 1349A, P. Trutmann 1984; IHR 1357A, P. Trutmann 1985; IHR 1392A,
1403A, 1404A, J.D. Taylor 1985; IHR 1410A, D.J. Allen 1985; IHR 1413A, M.A. Pastor-Corrales 1985;
IHR 1485A, D.J. Allen 1985; IHR 1499A, P. Trutmann 1985; IHR 1613A, 1614A, D.M. Teverson 1986;
IHR 1909A, J.D. Taylor 1987; IHR 2233A, 2252A, D.M. Teverson 1988).
Teverson (1991).

Mosaic: Southern bean mosaic virus.
Rubona (H.J. Vetten, unpubl. 1988).

On leaves: *Phoma sorghina* (Sacc.) Boerema, Dorenbr. & van Kest.
Coll. D.J. Allen, Rubona 1985 (det. M. Gerlagh).

Powdery mildew: *Erysiphe polygoni* DC. ex St-Am.
Haut Terres Volcaniques, P. Trutmann, unpubl. 1987.

Pythium root rot: *Pythium* spp.
Durnez (1983), Rusuku (1991), Buruchara (1992c), Buruchara & Rusuku (1992).

Rhizoctonia root rot: *Thanatephorus cucumeris* (Frank) Donk.
Nyabyenda *et al.* (1981), as *Rhizoctonia solani* Kuhn.
Rusuku (1991), Buruchara & Rusuku (1992).

Rust: *Uromyces appendiculatus* (Pers.) Unger.
Lac Mohasi, Mildbraed, 1911 (Hendrickx, 1948).

Southern blight: *Corticium rolfsii* Curzi.
Nyabyenda *et al.* (1981), as *Sclerotium rolfsii* Sacc.
Buruchara & Rusuku (1992).

White mould: *Sclerotinia sclerotiorum* (Lib.) de Bary.
Durnez (1983), R.A. Buruchara, unpubl. 1992.

SEYCHELLES

Cercospora leaf spot: *Cercospora ? canescens* Ellis & G. Mart.
On 'runner bean', so host *Phaseolus* species in question.
Mathias (1971).

Collar rot (wilt ?): *Fusarium oxysporum* Schlecht. f.sp. *phaseoli* J.B. Kendr. & W.C. Snyder.
Frequent heavy losses (Mathias, 1971).

SOMALIA

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Golato (1967a, 1967b), as *Cercospora columnaris* Ellis & Everh.

Bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
As *Pseudomonas phaseoli*, recorded by Curzi (1934) but from *Lablab purpureus* (as *Dolichos*) not from
Phaseolus vulgaris.

Rust: *Uromyces appendiculatus* (Pers.) Unger.
Golato (1967a).

Wilt: ? *Fusarium* sp.
C. Wortmann, pers. comm. 1989.

SOUTH AFRICA

Alternaria leaf spot (black blight): *Alternaria tenuissima* (Kunze) Wiltshire, and ? *Alternaria* sp., latter
species as *Macrosporium phaseoli* Fautrey.
Durban and Natal (Doidge *et al.*, 1953).

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Natal, Transvaal (Doidge *et al.*, 1953), as *Cercospora columnaris* Ellis & Everh.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
Widespread, common; often epiphytic (Doidge *et al.*, 1953).
Races not yet identified.

Ascochyta blight: *Phoma exigua* Desm. var. *exigua* and as *Ascochyta phaseolorum* Sacc.
Transvaal low veld and Natal.
Trench *et al.* (1986), A.J. Liebenberg, pers. comm. 1993.

Ashy stem blight: *Macrophomina phaseolina* (Tassi) Goid.
Vorster, 1964 cited by Gorter (1977).

Bacterial brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.
Transvaal; damaging in 1992 (Serfontein, 1994).

Cercospora leaf spot (leaf blotch): *Pseudocercospora cruenta* (Sacc.) Deighton.
Cape, Transvaal and especially Natal where it is most damaging (Doidge *et al.*, 1953), as *Cercospora cruenta*
Sacc.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
Doidge *et al.* (1953), Gorter (1977), as *X. phaseoli* (E.F. Sm.) Dowson and *X. phaseoli* var. *fusca* (Burkh.)
Starr & Burkh.

Common mosaic: Bean common mosaic virus.

Common and widespread (Doidge *et al.*, 1953).

Isolates from Transvaal and Natal identified as belonging to pathogenicity group V (Edington & Whitlock, 1988).

Fusarium root rot: *Fusarium solani* (Mart.) Sacc.

Doidge *et al.* (1953), Gorter (1977).

Fusarium wilt: *Fusarium oxysporum* Schlecht.? f.sp. *phaseoli* J.B. Kendr. & W.C. Snyder.

Doidge *et al.* (1953), Gorter (1977).

Grey mould: *Botrytis cinerea* Pers.

Seasonal and unimportant (A.J.L. Phillips, unpubl.).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

Doidge *et al.* (1953), as *Pseudomonas medicaginis* f.sp. *phaseolicola* (Burk.) Link & van Hall.

Races not yet identified.

Leaf scorch: attributed to the effects of high temperature and strong winds on plants suffering from manganese deficiency.

Southwestern Cape (Doidge *et al.*, 1953).

Mosaic: Cucumber mosaic virus.

M. Koch, unpubl.

Mosaic: Peanut mottle virus.

M. Koch, unpubl.

Mottle leaf: Non-parasitic.

Very common on Cape Flats resulting from manganese deficiency (Doidge *et al.*, 1953).

Phyllosticta leaf spot: *Phyllosticta phaseolina* Sacc.

Pretoria (Doidge *et al.*, 1953).

Pod blight: *Phoma subcircinata* Ellis & Everh.

Zululand, unimportant (Doidge *et al.*, 1953).

Pythium root rot (wilt): *Pythium debaryanum* Hesse and *Pythium ultimum* Trow.

Wager, 1941 cited by Gorter (1977).

Rhizoctonia root rot: *Thanatephorus cucumeris* (Frank) Donk.

Doidge & Bottomley, 1931 cited by Gorter (1977).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Doidge *et al.* (1953).

Scab: *Sphaceloma* anamorph of *Elsinoë phaseoli* Jenk.

Natal and Transvaal, often epiphytic.

Phillips (1994).

Virus-like disease: associated with a geminivirus.
Eastern Transvaal low veld.
G. Pietersen, pers. comm. 1994.

White mould (stem rot): *Sclerotinia sclerotiorum* (Lib.) de Bary.
Gorter (1977).

Wilt: associated with *Fusarium oxysporum* Schlecht. and *F. solani* (Mart.) Sacc. and their varieties as well as
with *Pythium debaryanum* Hesse, *P. spinosum* Sawada and *P. ultimum* Trow.
Unimportant except when growing conditions are unfavourable.
Dodge *et al.* (1953).

Yeast spot: *Nematospora coryli* Peglion and *Nematospora gossypii* Ashby & Nowell.
Dodge *et al.* (1953), as *Ashbya gossypii* (Ashby & Nowell) Guill.

Yellow dot, yellow mosaic: Alfalfa mosaic virus.
Minor (Neveling, 1956).

Yellow mosaic: Bean yellow mosaic virus.
M. Koch, unpubl.

SUDAN

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
IMI 323.
Occasional in Equatoria but rarely severe.
Kagelu, 1944 (Tarr, 1955), as *Isariopsis griseola* Sacc.

Ascochyta blight: *Ascochyta phaseolorum* Sacc.
Tarr (1955).

Ashy stem blight (charcoal rot): *Macrophomina phaseolina* (Tassi) Goid.
Tarr (1955), as *Macrophomina phaseoli* (Maubl.) Ashby.

Bacterial blight: Tarr (1955) suspected the presence of bacterial diseases on "various species of *Phaseolus*"
(which may have included spp. now consigned to the genus *Vigna*) "but little is known of them and
investigation is needed". *Xanthomonas phaseoli* was reported from the Sudan by Sabet (1959) and Sabet *et al.* (1969) but from legume hosts other than *P. vulgaris*.

Cercospora leaf spot: *Cercospora canescens* Ellis & G. Mart.
Tarr (1955).

Curly top: Cowpea mild mottle virus.
Attributed to a virus by Tarr (1955) who described the symptoms. Considered serious, since severely infected
plants are strongly distorted and sterile. Widespread in the North of the country but particularly severe
North of Khartoum where losses may be heavy, especially in hot weather. Regarded as the most important
disease of beans in the Sudan, especially when populations of the whitefly vector are dense (Salih & Salih,
1985). Recently found strongly associated with cowpea mild mottle carlavirus but field symptoms at Ed-
Damer could not be reproduced under greenhouse conditions (Vetten & Allen, 1991; H.J. Vetten, pers.
comm. 1993).

Mosaic: ? Virus.

Plants stunted, leaves being dwarfed, asymmetrical in shape with distinct light-dark green mosaic. Leaf laminae narrowed, plants sterile. Nuba Mountains (Tarr, 1955). BCMV not found in recent surveys (Vetten & Allen, 1991).

On leaves: *Periconia byssoides* Pers.

IMI 48174.

Katire, 1951, Tarr (1955).

On seed: *Aspergillus flavus* Link.

Habish (1972).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Widespread in the southern and central Sudan south of the Gezira, often causing severe damage (Tarr, 1955).

Sooty mould: *Cladosporium* sp.

Red Sea Coast (Tokar).

Tarr (1963).

Wilt: attributed to sodium toxicity but associated also with *Fusarium* sp.

Salih & Salih (1985).

SWAZILAND

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.

Minor importance.

Lin (1984), Y.P. Rao, pers. comm. 1987.

Ashy stem blight, charcoal rot: *Macrophomina phaseolina* (Tassi) Gooid.

Big Bend.

Allen & Ampofo (1992).

Bacterial wilt of seedlings: *Burkholderia solanacearum* (E.F. Sm.) Yabuuchi *et al.*

Y.P. Rao, pers. comm. 1987, as *Pseudomonas solanacearum* (E.F. Sm.) E.F. Sm.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

Lin (1984), Allen (1987b).

Common mosaic: Bean common mosaic virus.

Lin (1984), Y.P. Rao, pers. comm. 1987.

Pathogenicity groups IVb and VIa (Spence & Walkey, 1991).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

Y.P. Rao, pers. comm. 1987.

Race 6 (IHR 2688A, J.D. Taylor 1990).

Teverson (1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Rust: *Uromyces appendiculatus* (Pers.) Unger.
Lin (1984), Allen (1987b).

Seedling mortality: *Rhizoctonia* sp.
Y.P. Rao, pers. comm. 1987.

TANZANIA

Alternaria leaf spot: *Alternaria tenuissima* (Kunze) Wiltshire, IMI 119349a,d, *A. alternata* (Fr.) Keissler, IMI 159445, 159944, *A. macrospora* Zimm., IMI 110997, 112075, and *Alternaria* sp., IMI 110717.
Ebbels & Allen (1979).

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Wallace & Wallace (1949), as *Isariopsis griseola* Sacc.
Ebbels & Allen (1979). A more aggressive variant reported from Arusha by Hocking (1967).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
IMI 21059, 106512, 112167, 112168, 118796, 118797, 118798, 118799, 118803.
Wallace & Wallace (1949).

Ascochyta blight (leaf blotch, leaf spot): *Phoma exigua* Desm. var. *diversispora* (Bubák) Boerema.
Wallace & Wallace (1953), as *Ascochyta phaseolorum* Sacc. Positively identified as var. *diversispora*, coll.
D.J. Allen 1985 (Gerlagh, 1987).

Ashy stem blight (charcoal rot, as 'root disease'): *Macrophomina phaseolina* (Tassi) Goid.
IMI 22161, 281338.
Wallace & Wallace (1949), Riley (1960), as *M. phaseoli* (Maubl.) Ashby.

Associated fungi: *Acremonium* sp., IMI 245763.
Arthrinium phaeospermum (Corda) M.B. Ellis, IMI 159443.
Ebbels & Allen (1979).
Botrytis cinerea Pers., IMI 73755, 118653.
Cladosporium sp., IMI 76591.
. *Cladosporium sphaerospermum* Penz., IMI 91250b, 91251b.
Ebbels & Allen (1979).
Colletotrichum dematium (Pers.) Grove, IMI 91245.
Fusarium sulphureum Schlecht., IMI 96598.
Hyalodendron sp., IMI 279911.
Periconia byssoides Pers., IMI 76591c.
Verticillium sp., IMI 91257.
Ebbels & Allen (1979).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.
IHR 2566A-B, 2567A-B, D.J. Allen 1989 (D.M. Teverson, pers. comm. 1993).

Cercospora leaf spot: *Cercospora canescens* Ellis & G. Mart.
IMI 112081, 119349d.
and *Cercospora* sp., IMI 8178a, 118680, 106747, 134068, 133602.
Ebbels & Allen (1979).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

IMI B 2702 and probably IMI B 2636, 2637, 2638, 2639, 2702.

First recorded on seed in 1964 (Ebbels & Allen, 1979) as *X. phaseoli* (E.F. Sm.) Dows. and *X. phaseoli* var. *fuscanus* (Burkh.) Starr & Burkh.

Common mosaic: Bean common mosaic virus.

First reported by Wallace & Wallace (1949) and confirmed by Kulkarni (1973). Strains identified by Mink (1985), Silbernagel *et al.* (1986) and H.J. Vetten (unpubl.).

Pathogenicity groups I, III, IVb and VIa (Spence, 1992).

Damping off: *Thanatephorus cucumeris* (Frank) Donk.

Wallace & Wallace (1953), Riley (1960), as *Corticium solani* (Prill. & Delacr.) Bourd. & Galz.

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.

Wallace & Wallace (1949), as *Erostrotheca multiformis* G.H. Martin & Charles.

Riley (1960), as *Hyalodendron album* (Dowson) Diddens.

Ebbels & Allen (1979).

From stem pith and cotyledons: *Erwinia* sp., probably *E. carotovora* (Jones) Bergey *et al.*

IMI B 2835, 2836.

Ebbels & Allen (1979).

Fusarium root rot (as collar rot): *Fusarium* sp.

Riley (1960).

Grey mould: *Botrytis cinerea* Pers.

Riley (1960).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

Wallace & Wallace (1949), Riley (1960), as *Pseudomonas phaseolicola* (Burkh.) Dows.

Pathogenic variation studied by Gondwe (1987), Teverson (1991) and Mabagala & Saettler (1992), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Race 2 (IHR 2568A, D.J. Allen 1989), race 3 (IHR 1301A, J.H.C. Davis 1984; IHR 1427A, J.D. Taylor 1985), race 4 (IHR 1727A, 1729A, D.M. Teverson 1986; IHR 2569A, D.J. Allen 1989), race 5 (IHR 99/NCPPB 1341, 1962) and race 6 (IHR 1299A, 1300A, J.H.C. Davis 1984; IHR 1303A, P. Trutmann 1984; IHR 1420A, J.D. Taylor 1985; IHR 1514B, D.J. Allen 1986; IHR 1725A, D.M. Teverson 1986; IHR 1987A, J.B. Smithson 1988).

Leaf spot: *Mycosphaerella pinodes* (Berk. & Bloxam) Vestergr.

Wallace & Wallace (1949), Riley (1960).

Mild mottle: Cowpea mild mottle virus.

Morogoro (Mink, 1985) and Lushoto, Tanga (Vetten & Allen, 1991).

Oily pod: Considered to be a physiological disorder by Riley (1960), but it seems possible that 'oily pod' was in fact a symptom from infection either with bean common mosaic virus or *Xanthomonas campestris* pv. *phaseoli*. Zaumeyer & Thomas (1948) referred to 'greasy pod' being caused by a necrosis-inducing strain of BCMV, and a greasiness of pods in common bacterial blight is well-known.

Pythium root rot (Wilt): *Pythium ultimum* Trow.
Wallace & Wallace (1953).

Powdery mildew: *Leveillula taurica* (Lév.) G. Arnaud.
IMI 73753, 8178b.
Riley (1960).

Rust: *Uromyces appendiculatus* (Pers.) Unger.
IMI 65423, 112080, 126948, 126954, 126955, 126956, 126957.
Wallace & Wallace (1949), Riley (1960).
Races identified by Howland & Macartney (1966), Allen (1975b), Mmbaga & Stavely (1987). At least 25 races now recognized in Tanzania (M.T. Mmbaga, pers. comm. 1992).

Scab: *Elsinoë* sp., presumably *E. phaseoli* Jenkins.
IMI 237697.
Uyole, Mbeya 1979; also Arusha and Kilimanjaro Regions (D.J. Allen, unpubl.).

Schorch: non-parasitic.
Wallace & Wallace (1949), Riley (1960).

Southern blight: *Corticium rolfsii* Curzi.
IMI 119385, 121765, 259574.
Ebbels & Allen (1979).

Web blight: *Thanatephorus cucumeris* (Frank) Donk.
IMI 92529, 76533, 279916.

White mould: *Sclerotinia sclerotiorum* (Lib.) de Bary.
Observed on beans for the first time in 1942 in Arusha. In the same year, the fungus caused considerable damage to beans at Oldeani and near Moshi (Wallace, 1944). The disease continues to be damaging in Arusha and Kilimanjaro Regions (Msuya *et al.*, 1992).

Wilt: *Fusarium* spp.
Wallace & Wallace (1953), Riley (1960).

Yeast spot (Stigmatomycosis): *Nematospora coryli* Peglion.
Wallace & Wallace (1949).

Yellow mosaic: Bean yellow mosaic virus.
Recorded by Wallace & Wallace (1949) but, since subsequent surveys have not confirmed the presence of BYMV in Tanzania, the record has to be regarded with suspicion.

UGANDA

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
IMI 96166b, 102842b, 102825b, 7158b, 85772, 144489, 91775b, 93654, 93602a, 91347b, 93593a, 98742, 98777, 98778a, 98749c.
Hansford (1937), as *Isariopsis griseola* Sacc.

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
 IMI 102877, 102878, 102879, 102880, 102881, 102882, 102883, 102884, 102885, 102887, 103577, 93602b,
 98794, 102886.

Hansford (1937), as *Glomerella lindemuthianum* Shear.
 Pathogenic variation studied by Leakey & Simbwa-Bunya (1972).

Ascochyta blight (leaf spot): *Phoma exigua* Desm. var. *diversispora* (Bubák) Boerema.
 IMI 91773, 91741, 91750a, 93582, 93622a, 93591, 93602b, 98796.

Hansford (1937), as *Ascochyta* sp. and *A. phaseolorum* Sacc.
 Positive identification as var. *diversispora*, coll. nr Kabale, D.J. Allen 1985 (Gerlagh, 1987).

Ashy stem blight (charcoal rot): *Macrophomina phaseolina* (Tassi) Goid.
 Hansford (1937), as *M. phaseoli* (Maubl.) Ashby.

Associated fungi: *Alternaria alternata* (Fr.) Keissler, IMI 164951, 268982.

Alternaria sp., IMI 123945.

Cladosporium sp., IMI 164959.

Colletotrichum truncatum (Schwein.) Andrus & Moore, IMI 164955.

Epicoccum nigrum Link, as *E. purpurascens* Ehrenb. ex Schlecht., IMI 93622b, 164961, 165518.

Gibberella sp., IMI 91381.

Glomerella cingulata (Stonem.) Spauld. & Schrenk, IMI 95999.

Mycosphaerella pinodes (Berk. & Bloxam) Vestergr., Hansford (1943).

Nematospora coryli Pegl., IMI 77422.

Phoma sorghina (Sacc.) Boerema, Dorenb. & van Kest., IMI 165526, 165517.

Phoma sp., IMI 102865, 102866, 102867, 102869, 102870, 102871, 102872, 102873, 102875, 102876,
 164956.

Pleospora sp., Hansford (1937).

Speira punctulata Cooke & Ellis var. *latebrosa* Bizz., Hansford (1943).

Xylaria hypoxylon (Fr.) Grev., IMI 165539.

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.

IHR 1931A-D, 1932A-D, 1943A-D, 1945A-D, J.D. Taylor 1987; IHR 2795A-B, 2800A-B, 2801A-B, 2811B-C, D.M. Teverson 1991.

D.M. Teverson, pers. comm. 1993.

Cercospora leaf spot: *Pseudocercospora cruenta* (Sacc.) Deighton and *Cercospora canescens* Ellis & G. Mart.

Hansford (1937), former species as *Cercospora cruenta* Sacc.

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F.Sm.) Dye.

Leakey (1973). Pathogenic variation studied by Opio (1993).

Common mosaic, black root: Bean common mosaic virus.

Kulkarni (1973). Necrosis-inducing strains predominant (Owera, 1990) and pathogenicity groups I, IVb and VIa identified (Spence & Walkey, 1991; Spence, 1992).

Dactuliophora leaf spot: *Dactuliophora tarpii* Leakey.

IMI 98272.

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.
IMI 7158a, 91750b, 101370, 102825a, 102842a, 91775a, 98749b, 96166a.
Hansford (1937), as *Cladosporium album* Dowson.

Golden mosaic: ? Whitefly-transmitted geminivirus.
Kabanyolo (H.J. Vetten, pers. comm. 1993).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
Probably of minor importance in the hotter and lower areas (Leakey, 1970). Common and damaging in the
Kigezi highlands of Kabale.
Race 3 (IHR 1334A, T.N. Sengooba 1984) and race 4 (IHR 1460A and 1490A, D.J. Allen, 1985). Teverson
(1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Mild mottle: Cowpea mild mottle virus.
Kabanyolo, Kampala.
Vetten & Allen (1991).

Mosaic: Bean strains of blackeye cowpea mosaic virus.
Kabanyolo (H.J. Vetten, pers. comm. 1993).

Powdery mildew: *Oidium* sp., suspected to be *Erysiphe polygoni* DC. ex St-Am.
Hansford (1937).

Root rot: *Rhizoctonia solani* Kuhn.
Hansford (1937).

Rust: *Uromyces appendiculatus* (Pers.) Unger.
IMI 91349, 91357, 74271, 85783, 91772, 93604, 98796, 98749, 98740, 162848, 96166.
Hansford (1937).
Pathogenic variation studied by Howland & Macartney (1966).

Scab: *Elsinoë phaseoli* Jenkins.
IMI 251363.

Southern blight: *Corticium rolfsii* Curzi.
Causes damage in early growth and later to pods carried near ground under wet conditions (Leakey, 1970), as
Pellicularia rolfsii (Sacc.) Westend.

Web blight: *Thanatephorus cucumeris* (Frank) Donk.
Kinkizi, Kabale, May 1985 (Allen & Trutmann, 1985).

ZAIRE

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
Mulungu, South Kivu (Hendrickx, 1939; Hendrickx, 1948), as *Isariopsis griseola* Sacc.
Louto, Ndihira, North Kivu (P. Trutmann and M. Pyndji, unpubl., Pyndji, 1988).
Bas Zaire, 1985 (P. Trutmann, unpubl.).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
 Mulungu (Hendrickx, 1939); Veamba, Kivu, 1942 (Steyaert, 1948).
 Goma, Loutu, Ndihira, North Kivu (P. Trutmann and M. Pyndji., unpubl.).

Ascochyta blight: *Ascochyta phaseolorum* Sacc.
 Mulungu, Kivu, D.L. Hendrickx, 1939 (Hendrickx, 1948).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.
 IHR 2242A-B, D.M. Teverson, 1988 (Teverson, 1991).

Cercospora leaf spot: *Cercospora canescens* Ellis & G. Mart.
 Mvuazi, Bas Zaire (INEAC, 1959).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
 North Kivu, 1985 (P. Trutmann, unpubl.)
 Bas Zaire, 1987 (P. Trutmann, unpubl.)

Common mosaic: Bean common mosaic virus.
 An early report (INEAC, 1939) mentions attack by a virus disease transmitted by a jassid and a capsid at Nioka in North Kivu, and there is reference to mosaic at Mulungu, Kivu (INEAC, 1949). BCMV positively identified much more recently, and pathogenicity groups III and VIa found (Spence & Walkey, 1991; Spence, 1992).

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.
 Mulungu, South Kivu; and Rushuru, Ndihira, North Kivu, 1985 (P. Trutmann, unpubl.).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
 Race 2 (IHR 1507A, P. Trutmann 1986), race 3 (IHR 1412A, M.A. Pastor-Corrales 1985) and race 4 (IHR 1414A, M.A. Pastor-Corrales, 1985). Teverson (1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

'On bean': *Phytophthora phaseoli* Thaxt.
 Mulungu, Kivu (Hendrickx, 1939; Hendrickx, 1948).

'On bean': *Stagonosporopsis hortensis* (Sacc. & Malbr.) Petr.
 Mulungu, Kivu (Hendrickx, 1939), as *Ascochyta boltshauseri* Sacc.

Rust: *Uromyces appendiculatus* (Pers.) Unger.
 Kasai (Steyaert, 1927 in Hendrickx, 1948); severe in Gandajika (INEAC, 1944). Mulungu, Kivu (Hendrickx, 1939).

Southern blight: *Corticium rolfsii* Curzi.
 South Kivu and Bas Zaire, 1987 (P. Trutmann, unpubl.).

Web blight: *Thanatephorus cucumeris* (Frank) Donk.
 Yangambi, 1956 (INEAC, 1958a), as *Rhizoctonia solani* Kuhn.
 Prevalent and damaging during periods of rain (Lama, 1989).

ZAMBIA

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.
IMI 103514, 87901, 55938, 85670, 89743, 89729, 90008, 90009, 95374, 130491b.
Central & Eastern Provinces (Riley, 1956), as *Isariopsis griseola* Sacc.

Alternaria leaf spot: *Alternaria tenuissima* (Kunze) Wiltshire, *A. solani* Sorauer, *A. tomato* (Cooke) G.F. Weber and *Alternaria* sp.
IMI 163680, 169475, 121311b, 68953.
Angus (1962-66).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.
IMI 68953a, 166107b.
Central & Eastern Provinces (Riley, 1956).

Ascochyta blight (leaf spot): *Phoma exigua* Desm. var. *diversispora* (Bubák) Boerema, and *Ascochyta phaseolorum* Sacc.
IMI 69319, 85698, 89773, 89777, 89789, 899986, 90009, 90010, 93582, 93591, 93602b, 93622a, 162994.
Angus (1962-66). Positive identification as var. *diversispora*, coll. at Nakonde, D.J. Allen 1985 (Gerlagh, 1987).

Ashy stem blight (charcoal rot): *Macrophomina phaseolina* (Tassi) Goid.
Central Province (Riley, 1956), as *M. phaseoli* (Maubl.) Ashby.

Associated fungi: *Aspergillus flavus* Link, IMI 92573.
Cladosporium herbarum (Pers.) Link, IMI 287637.
Colletotrichum truncatum (Schwein.) Andrus & Moore, IMI 68165, 89724, 90022.
Corticium sp., IMI 74082.
Epicoccum nigrum Link, as *E. purpurascens* Ehrenb., IMI 85685e.
Fusarium sp., IMI 85685b.
Gloeosporium sp., IMI 90019.
Phoma sp., IMI 287649.
Pleospora sp., IMI 69305a.
Pseudoplea trifolii (Rostr.) Petr., IMI 85695.
Angus (1962-66).
Pythium myriotylum Drechsl., IMI 184696.

Brown spot?: *Pseudomonas syringae* pv. *syringae* van Hall.
Angus (1962-66).

Cercospora leaf spot: *Cercospora canescens* Ellis & G. Mart. and *Cercospora* sp.
IMI 90020, 93860, 130493.
Angus (1962-66).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.
IMI B 12509, B 12501, 75700, 89777.
Angus (1962-66), J. Kannaiyan (pers. comm. 1989).

Floury leaf spot: *Mycovellosiella phaseoli* (Drummond) Deighton.

Angus (1962-66), as *Ramularia deusta* (Fuckel) Karakulin.

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

Western & Central Provinces (Riley, 1956).

Eastern Province, race 6 (IHR 1304A and 1361A, J. Kannaiyan 1985). Teverson (1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Leptosphaerulina leaf spot: *Leptosphaerulina trifolii* (Rostr.) Petr.

IMI 85695, 85685a.

Mosaic, black root: Bean common mosaic virus.

Central Province (Riley, 1956).

Northern Province (Angus, 1962-66).

Eastern Province.

Both serotypes A and B present (Vetten & Allen, 1991) and strains from pathogenicity groups IVb and VIa identified. Sohati *et al.* (1992), Spence (1992).

Mosaic: Cucumber mosaic virus.

Msekera, Chipata, Eastern Province (Vetten & Allen, 1991).

Mosaic: Peanut mottle virus.

Lusaka and Msekera, Chipata, Eastern Province (Vetten & Allen, 1991).

Mycosphaerella leaf spot: *Mycosphaerella pinodes* (Berk. & Bloxam) Vestergr.

IMI 93861b, 287635.

Angus (1962-66).

Phyllosticta leaf spot: *Phyllosticta phaseolina* Sacc. and *Phyllosticta* sp.

IMI 93858, 166107a.

Angus (1962-66).

Powdery mildew: *Oidium* sp. and *Phyllactinia* ? *corylea* (Pers.) Karst.

IMI 119634, 70818.

Angus (1962-66).

Red leaf spot: *Phoma* ? *sorghina* (Sacc.) Boerema, Dorenbr. & van Kest.

IMI 277713.

Nr Nakonde (Greenberg *et al.*, 1973).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

IMI 89744, 89772, 90001c, 85713, 68164, 103514, 110743, 115404, 165201a.

Central & Western Provinces (Riley, 1956).

Northern & Eastern Provinces (Angus, 1962-66).

Scab: *Elsinoë phaseoli* Jenkins.

IMI 77503, 130493a, 130471.

Central Province (Angus, 1962-66). Northern Province, where sources of resistance were identified (Kannaiyan *et al.*, 1990).

Southern blight (stem rot or collar rot): *Corticium rolfsii* Curzi.

IMI 90021.

Angus (1962-66), as *Sclerotium rolfsii* Sacc.

Stem anthracnose: *Colletotrichum truncatum* (Schwein.) Andrus & Moore.

Angus (1962-66).

Web blight: *Thanatephorus cucumeris* (Frank) Donk.

Nr Nakonde (Greenberg *et al.*, 1983).

Wilt: *Fusarium equiseti* (Corda) Sacc.

Southern Province (Riley, 1956).

Yeast spot: *Nematospora coryli* Peglion.

Central Province (Riley, 1956).

Yellow mosaic: Bean yellow mosaic virus.

Riley (1956).

Dubious. Presence of BYMV not confirmed in recent virus surveys (Vetten & Allen, 1991).

ZIMBABWE

Alternaria leaf spot: *Alternaria solani* Sorauer, *Alternaria* sp. and *Alternaria tomato* (Cooke) G.F. Weber.

IMI 68953b.

Whiteside (1966), Rothwell (1983).

Angular leaf spot: *Phaeoisariopsis griseola* (Sacc.) Ferraris.

IMI 59542, 99839b.

Whiteside (1966), Rothwell (1983).

Anthracnose: *Colletotrichum lindemuthianum* (Sacc. & Magn.) Briosi & Cavara.

IMI 99839a, 21062.

Whiteside (1966), Rothwell (1983).

Ashy stem blight: *Macrophomina phaseolina* (Tassi) Goid.

IMI 92595.

Whiteside (1966), Rothwell (1983), as *M. phaseoli* (Maubl.) Ashby.

Bald Head: Mechanical injury to seed.

Whiteside (1966).

Brown spot: *Pseudomonas syringae* pv. *syringae* van Hall.

IHR 2692C, 2703C, J.D. Taylor 1990 (Teverson, 1991).

Cercospora leaf spot: *Cercospora canescens* Ellis & G. Mart.

Whiteside (1966), Rothwell (1983).

Common bacterial blight: *Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.) Dye.

Whiteside (1966), Rothwell (1983), as *X. phaseoli* (E. F. Sm.) Dows, and *X. phaseoli* var. *fusca*s (Burkh.) Starr & Burkh.

Common mosaic: Bean common mosaic virus.

Whiteside (1966), Rothwell (1983).

Necrosis-inducing strains predominant. Pathogenicity groups I, IVa, IVb and VIa identified.

Spence & Walkey (1991), Mukoko (1992), Spence (1992).

Crown rot: *Sclerotium rolfsii* Sacc.

Archibald (1974).

Damping-off: *Pythium* sp. and *Rhizoctonia solani* Kuhn.

Whiteside (1966), Rothwell (1983).

Fusarium wilt: *Fusarium* sp.

Whiteside (1966), Rothwell (1983).

Halo blight: *Pseudomonas savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*

Whiteside (1966), Rothwell (1983), as *Pseudomonas phaseolicola* (Burkh.) Dows.

Race 2 (IHR 2370A, J.B. Smithson 1989; IHR 2698B and 2700A, J.D. Taylor 1990) and race 6 (IHR 2692A, 2693A, 2694A, 2698C and 2703A, J.D. Taylor 1990). Teverson (1991), as *Pseudomonas syringae* pv. *phaseolicola* (Burkh.) Young *et al.*

Intumescence: Physiological.

Whiteside (1966).

Mosaic: Cucumber mosaic virus, often in mixed infections with peanut mottle virus.

Gwebi, Harare and Shamva (Vetten & Allen, 1991).

Pink rot: *Trichothecium roseum* (Pers.) Link.

Whiteside (1966), Rothwell (1983).

Rhizoctonia root rot: *Thanatephorus cucumeris* (Frank) Donk, as *Rhizoctonia solani* Kuhn.

Whiteside (1966), Rothwell (1983).

Rust: *Uromyces appendiculatus* (Pers.) Unger.

Whiteside (1966), Rothwell (1983).

Scab: *Elsinoë phaseoli* Jenkins.

IMI 56161, 56102.

Whiteside (1966), Rothwell (1983).

Southern blight (root rot): *Sclerotium rolfsii* Sacc.

Whiteside (1966), Rothwell (1983).

White mould: *Sclerotinia sclerotiorum* (Lib.) de Bary.

Rothwell (1983).

Yeast spot: *Nematospora coryli* Peglion.

Follows injury by sucking bug.

Whiteside (1966), Rothwell (1983).

FUNGAL PATHOGEN INDEX

- Acremonium* sp.
TZ
- Alternaria alternata* (Fr.) Keissler
BU, RW, TZ, UG
- A. cucumerina* (Ellis & Everh.) J.E. Elliot
KE
- A. longipes* (Ellis & Everh.) Mason
MD
- A. macrospora* Zimm.
?ZA, TZ
- A. solani* Sorauer
ET, MD, ZA, ZW
- A. tenuissima* (Kunze) Wiltshire
AN, ET, SA, TZ, ZA
- A. tomato* (Cooke) G.F. Weber
ZA, ZW
- Alternaria* sp.
KE, MW, ?SA, TZ, UG, ZA, ZW
- Arthrinium phaeospermum* (Corda) M.B. Ellis
TZ
- Ascochyta boltshauseri* Sacc.
(See *Stagonosporopsis hortensis*
(Sacc. & Malbr.) Petr.)
KE, ZR
- A. phaseolorum* Sacc.
(See *Phoma exigua* Desm. var. *exigua*)
AN, BU, ET, KE, MW, RW, SA, SD, TZ, UG,
ZA, ZR
- Ascochyta* sp.
MW, UG
- Aspergillus flavus* Link.
ZA, SD
- Botrytis cinerea* Pers.
(Teleomorph *Sclerotinia fuckeliana*
(de Bary) Fuckel)
SA, TZ
- Cercospora canescens* Ellis & G. Mart.
MA, MW, ?SY, SD, TZ, UG, ZA, ZW, ZR
- C. columnaris* Ellis & Everh.
(See *Phaeoisariopsis griseola* (Sacc.) Ferraris)
ET, SO, SA,
- C. cruenta* Sacc.
(See *Pseudocercospora cruenta*
(Sacc.) Deighton)
AN, ET, MD, SA, UG
- Cercospora* sp.
MW, TZ, ZA
- Cercosporidium* sp.
ET
- Chaetomium spirale* Zopf.
KE
- Cladosporium album* Dowson
(See *Mycovellosiella phaseoli* (Drummond)
Deighton)
- UG
- C. herbarum* (Pers.) Link
ZA
- C. oxysporum* Berk. & M.A. Curtis
ET
- C. sphaerospermum* Penz.
TZ
- Cladosporium* sp.
ET, KE, SD, TZ, UG
- Colletotrichum dematium* (Pers.) Grove
TZ
- C. lindemuthianum* (Sacc. & Magn.) Briosi & Cavara
AN, BU, ET, KE, MD, MW, MA, MO, RW, SA,
TZ, UG, ZR, ZA, ZW
- C. truncatum* (Schwein.) Andrus & Moore
ET, MW, UG, ZA
- Corticium rolfsii* Curzi
(Sclerotial state *Sclerotium rolfsii* Sacc.)
AN, BU, ET, MW, MD, MA, RW, TZ, UG, ZA,
ZR
- C. solani* (Prill. & Delacr.) Bourd. & Galz.
(See *Thanatephorus cucumeris* (Frank)
Donk; Sclerotial state *Rhizoctonia solani* Kuhn)
AN, KE, MD, MA, MW, MA, TZ
- Corticium* sp.
ZA
- Dactuliothora tarrii* Leakey
UG
- Elsinoë phaseoli* Jenkins
(Anamorph *Sphaceloma* sp.)
KE, MW, SA, TZ, UG, ZA, ZW
- Epicoccum nigrum* Link
(Syn. *E. purpurascens* Ehrenb.)
KE, UG, ZA
- E. purpurascens* Ehrenb.
KE, MW, UG, ZA
- Erostrotheca multiformis* G.H. Martin & Charles
(See *Mycovellosiella phaseoli* (Drummond)
Deighton)
- TZ

- Erysiphe polygoni* DC. ex St-Am.
KE, MA, RW, UG
- Fusarium equiseti* (Corda) Sacc.
ZA
- F. moniliforme* J. Shield.
(Teleomorph *Gibberella fujikuroi* (Sawada) Ito)
MW
- F. oxysporum* Schlecht.
ET, ZA
- F. oxysporum* Schlecht. f.sp. *phaseoli* J.B. Kendr. &
W.C. Snyder
KE, RW, SY, ?SA
- F. semitectum* Berk. & Rav.
KE
- F. solani* (Mart.) Sacc.
(Teleomorph *Nectria haematococca* Berk. &
Broome)
BU, ET, MW, MA, RW, SA
- F. solani* (Mart.) Sacc. f.sp. *phaseoli* (Burkh.) W.C.
Snyder & E.C. Hansen
AN, KE
- F. sulphureum* Schlecht.
TZ
- Fusarium* sp.
MD, MO, ?SO, SD, TZ, ZA, ZW
- Gibberella fujikuroi* (Sawada) Ito
(Teleomorph *Fusarium moniliforme* J. Shield.)
MW
- Gibberella* sp.
UG
- Gloeosporium* sp.
ZA
- Glomerella cingulata* (Stonem.) Spauld. & Schrenk.
UG
- G. lindemuthianum* Shear
(Putative teleomorph of *Colletotrichum*
lindemuthianum (Sacc. & Magn.) Briosi &
Cavara)
UG
- Hyalodendron album* (Dowson) Diddens
(See *Mycovellosiella phaseoli* (Drummond)
Deighton)
TZ
- Hyalodendron* sp.
TZ
- Isariopsis griseola* Sacc.
(See *Phaeoisariopsis griseola* (Sacc.) Ferraris)
AN, BU, KE, MA, RW, SD, UG, ZR, ZA
- Leptosphaerulina trifolii* (Rostr.) Petr.
KE, ZA
- Leveillula taurica* (Lév.) G. Arnaud
TZ
- Macrophomina phaseoli* (Maubl.) Ashby
(See *M. phaseolina* (Tassi) Goid.)
KE, SD, TZ, UG, ZA, ZW
- M. phaseolina* (Tassi) Goid.
ET, KE, LO, MA, RW, SA, SD, WD, TZ, UG,
ZA, ZW
- Macrosporium phaseoli* Fautrey
(See *Alternaria macrospora* Zimm. ?)
SA
- M. pinodes* (Berk. & Bloxam) Vestergr.
ET, MW, TZ, UG, ZA
- Mycovellosiella phaseoli* (Drummond) Deighton
BU, ET, KE, MW, RW, TZ, UG, ZR, ZA
- Nematospora coryli* Peglion
KE, SA, TZ, UG, ZA, ZW
- N. gossypii* Ashby & Nowell
SA
- Nematospora* sp.
AN
- Neocosmospora vasinfecta* var. *vasinfecta* E.F.
Sm.
(Anamorph *Cephalosporium* sp. =
Acremonium sp.)
LO
- Oidium* sp.
UG, ZA
- Pellicularia rolfsii* (Sacc.) Westend.
(See *Corticium rolfsii* Curzi)
UG
- Penicillium cyclopium* Westling
KE
- Periconia byssoides* Pers.
ET, MW, SD, TZ
- Phaeoisariopsis griseola* (Sacc.) Ferraris
AN, BU, ET, KE, MD, MW, MA, MO, RW, SO,
SA, SD, WD, TZ, UG, ZR, ZA, ZW
- Phoma exigua* Desm.
ET, KE, MO, SA
- P. exigua* Desm. var. *diversispora* (Bubák) Boerema
BU, KE, ?MD, RW, TZ, UG, ZA
- P. phaseolina* Pass.
ET
- P. sorghina* (Sacc.) Boerema, Dorenb. & van Kest.
ET, RW, UG, ?ZA
- P. subcircinata* Ellis & Everh.
SA
- Phoma* sp.
UG, ZA

- Phyllactinia ? corylea* (Pers.) Karst.
ZA
- Phyllosticta phaseolina* Sacc.
ET, KE, MA, SA, ZA
- Phyllosticta* sp.
ET, MW, ZA
- Phytophthora phaseoli* Thaxt.
ZR
- Pleospora* sp.
MA, UG, ZA
- Pseudocercospora cruenta* (Sacc.) Deighton
(Teleomorph *Mycosphaerella cruenta* Latham)
AN, ET, MD, SA, UG
- Pseudoplea trifolii* (Rostr.) Petr.
(See *Leptosphaerulina trifolii* (Rostr.) Petr.)
ZA
- Pythium debaryanum* Hesse
SA
- P. myriotylum* Drechs.
ZA
- P. spinosum* Sawada
SA
- P. ultimum* Trow
SA, TZ
- Pythium* sp.
BU, KE, RW, ZW
- Ramularia deusta* (Fuckel) Karakulin
(See *Mycovellosiella phaseoli* (Drummond)
Deighton)
ZA
- R. phaseoli* (Drummond) Deighton
(See *Mycovellosiella phaseoli* (Drummond)
Deighton)
BU, ET, KE, MW
- Rhizoctonia solani* Kuhn
(Teleomorph *Thanatephorus cucumeris* (Frank)
Donk)
BU, ET, KE, MO, RW, SA, UG, ZR, ZA, ZW
- Rhizoctonia* sp.
BU, WD
- Rhizopus stolonifer* (Ehrenb.) Lind
MW
- Sclerotinia sclerotiorum* (Lib.) de Bary
BU, ET, KE, MA, RW, SA, TZ, ZW
- Sclerotium rolfsii* Sacc.
(Teleomorph *Corticium rolfsii* Curzi)
AN, BU, ET, KE, MD, RW, ZA, ZW
- Speira punctulata* Cooke & Ellis var. *latebrosa* Bizz.
UG
- Sphaceloma* sp.
SA
- Stagonosporopsis hortensis* (Sacc. & Malbr.) Petr.
KE, ZR
- Thanatephorus cucumeris* (Frank) Donk
(Sclerotial state *Rhizoctonia solani* Kuhn)
BU, ET, KE, MD, MW, MA, MD, MO, RW, SA,
TZ, UG, ZR, ZA, ZW
- Trichothecium roseum* (Pers.) Link
ZW
- Uromyces appendiculatus* (Pers.) Unger
AN, BU, ET, KE, LO, MD, MW, MA, MO, RW,
SO, SA, SD, WD, TZ, UG, ZR, ZA, ZW
- U. phaseoli* (Pers.) Wint. var. *typica* Arthur
(See *U. appendiculatus* (Pers.) Unger)
ET
- U. viciae-fabae* (Pers.) Schroet.
?KE
- Verticillium* sp.
TZ
- Xylaria hypoxylon* (Fr.) Grev.
UG

BACTERIAL PATHOGEN INDEX

- Bacterium medicaginis* var. *phaseolicola* (Burkh.)
 Link & van Hall
 (See *Pseudomonas savastanoi* pv. *phaseolicola*
 (Burk.) Gardan *et al.*)
 KE, SA
- Bacterium phaseoli* E.F. Sm.
 (See *Xanthomonas campestris* pv. *phaseoli*
 (E.F. Sm.) Dye)
 MD
- Burkholderia solanacearum* (E.F. Sm.) Yabuuchi *et al.*
 MD, MA, WD
- Corynebacterium flaccumfaciens* (Hedges) Dowson
 (See *Curtobacterium flaccumfaciens* (Hedges)
 Collins & Jones)
 KE, MA
- Curtobacterium flaccumfaciens* (Hedges) Collins &
 Jones
 KE, MA
- Erwinia* ? *carotovora* (Jones) Bergey *et al.*
 TZ
- Pseudomonas phaseolicola* Burkh.) Dowson
 (See *Ps. syringae* pv. *phaseolicola* (Burkh.)
 Gardan *et al.*)
 MA
- Ps. savastanoi* pv. *phaseolicola* (Burkh.) Gardan *et al.*
 BU, ET, KE, LO, MW, MA, MO, RW, SA, TZ,
 UG, WD, ZA, ZR, ZW
- Ps. solanacearum* (E.F. Sm.) E.F. Sm.
 (See *Burkholderia solanacearum* (E.F. Sm.)
 Yabuuchi *et al.*)
 MD, MA, WD
- Ps. syringae* pv. *phaseolicola* (Burk.) Young *et al.*
 (See *Ps. savastanoi* pv. *phaseolicola* (Burkh.)
 Gardan *et al.*)
 BU, ET, KE, LO, MW, MA, MO, RW, SA, TZ,
 UG, WD, ZA, ZR, ZW
 Race distribution, see Table I.
- Ps. syringae* pv. *syringae* van Hall
 BU, KE, LO, MA, RW, SA, TZ, UG, ?ZA, ZR,
 ZW
- Xanthomonas campestris* pv. *phaseoli* (E.F. Sm.)
 Dye
 AN, BU, ET, KE, LO, MD, MW, MA, MO, RW,
 SA, TZ, UG, WD, ZA, ZR, ZW
- X. phaseoli* (E.F. Sm.) Dowson
 (See *X. campestris* pv. *phaseoli* (E.F. Sm.) Dye)
 MW, SA, TZ, ZW
- X. phaseoli* var. *fusca* (Burkh.) Starr & Burk.
 (See *X. campestris* pv. *phaseoli* (E.F. Sm.) Dye).
 MW, SA, TZ, ZW

VIRUS INDEX

Alfalfa mosaic virus	Cucumber mosaic cucumovirus
SA	BU, SA, ZA, ZW
Bean common mosaic potyvirus	Geminivirus
BU, ET, KE, LO, ?MD, MW, ?MA, RW, SA,	UG
WD, TZ, UG, ZR, ZA, ZW	Nepovirus
For distribution of pathogenicity groups, see Table	ET
2.	Peanut mottle potyvirus
Bean yellow mosaic potyvirus	ET, MD, SA, ZA, ZW
ET, KE, ? MA, SA; dubious ancient records from	Potyvirus aff. soybean mosaic
MW, TZ, ZA	ET
Blackeye cowpea mosaic potyvirus	Southern bean mosaic sobemovirus
BU, UG	RW
Cowpea mild mottle carlavirus	Tobacco mosaic tobamovirus
MW, MO, SD, TZ, UG	KE

DISTRIBUTION OF PATHOGENIC VARIANTS

Table 1. Geographical distribution¹ of races of *Pseudomonas savastanoi* pv. *phaseolicola* in Africa (after Teverson, 1991)

Country	Races on <i>Phaseolus vulgaris</i>					Additional races on other hosts	
Burundi		3	4				
Ethiopia			4	6		2	7
Kenya			4	6	7		
Lesotho	1	2		6	8		
Madagascar						1	
Malawi		2		5		9	
Mauritius		2					
Rwanda	1	2		4			
Swaziland					6		
Tanzania		2	3	4	5	6	
Uganda			3	4		7	8
Zaire					6		
Zambia		2	3	4			
Zimbabwe		2			6		

¹ Countries in which races have been identified.

Table 2. Geographical distribution¹ of the pathogenicity groups² comprising bean common mosaic virus in Africa³

Country	Pathogenicity groups					
Burundi	I	III		IVb	Va	VIa
Ethiopia	I			IVb		
Kenya					Va	VIa
Lesotho						VIa
Malawi				IVb		VIa
Rwanda		III		IVb		VIa
South Africa					V	
Tanzania	I	III		IVb		VIa
Uganda	I			IVb		VIa
Zaire		III				VIa
Zambia				IVb		VIa
Zimbabwe	I		IVa	IVb		VIa

¹ Only countries from where BCMV strains have been positively identified.

² Strains representative of pathogenicity groups I(NL1), II(NL7), III(NL8), IVa(Florida), IVb(NL6), Va(NY15), Vb(NL2), VIa(NL3), VIb(NL5) and VII(NL4). Serotype A consists of temperature independent necrosis inducing strains belonging to pathogenicity groups III, VIa and VIb, whereas B serotypes belong to pathogenicity groups I, II, IVa, Va and VII which do not induce necrosis, and groups IVb and Vb which may induce necrosis at high temperatures.

³ Sources: Edington & Whitlock (1988), Spence & Walkey (1991), Mukoko (1992), Spence (1992).

REFERENCES

- AGRANOVSKY, A.A. (1985). Identification of viruses infecting pulse crops in Ethiopia. Phytopathology Committee, Addis Ababa, pp. 73-80. *Proceedings of the 10th Annual Meeting Ethiopia*.
- ALLEN, D.J. (1975a). Additions to the fungi and plant diseases of Malawi. *Society of Malawi Journal* **28**: 35-44.
- ALLEN, D.J. (1975b). Breeding beans (*Phaseolus vulgaris* L.) for resistance to rust (*Uromyces appendiculatus* (Pers.) Ung.). PhD Thesis, University of Cambridge.
- ALLEN, D.J. (1983). *The Pathology of Tropical Food Legumes : Disease Resistance in Crop Improvement*. Chichester, John Wiley & Sons.
- ALLEN, D.J. (1987a). *Principal diseases of beans in Africa*. An audiotutorial unit, series 04EB-06.13. Cali, Colombia, Centro Internacional de Agricultura Tropical.
- ALLEN, D.J. (1987b). Trip to Swaziland, 17-19 September, 1987. Unpublished report, Centro Internacional de Agricultura Tropical, 10 pp.
- ALLEN, D.J. (1988). Trip to Angola, 15-21 November, 1988. Unpublished report, Centro Internacional de Agricultura Tropical, 7 pp.
- ALLEN, D.J. (1991). New disease records from legumes in Tropical Africa. *FAO Plant Protection Bulletin* **39**: 112-113.
- ALLEN, D.J. & AMPOFO, J.K.O. (1992). Trip to Swaziland, 8-11 April, 1992. Unpublished report, Centro Internacional de Agricultura Tropical.
- ALLEN, D.J., BURUCHARA, R. & SMITHSON, J.B. (unpubl.). Diseases of *Phaseolus* beans.
- ALLEN, D.J., DESSERT, M., TRUTMANN, P. & VOSS, J. (1989). Common Beans in Africa and their Constraints. In *Bean Production Problems in the Tropics*, 2nd Edn (ed. H.F. Schwartz & M.A. Pastor-Corrales), pp. 9-31. Cali, Colombia, Centro Internacional de Agricultura Tropical.
- ALLEN, D.J. & EDJE, O.T. (1990). Common Bean in African Farming Systems. In Bean Research, Volume 5. Progress in Improvement of Common Bean in Eastern and Southern Africa (ed. J.B. Smithson). *CIAT Africa Workshop Series No. 12* : 20-32.
- ALLEN, D.J., NWANZE, K.F. & WATT, E.E. (1976). Trip to Ethiopia, 7-22 September, 1976. Unpublished report, International Institute of Tropical Agriculture, Ibadan, 24 pp.
- ALLEN, D.J. & SMITHSON, J.B. (1989). Travel Report : Lesotho, 29 January-5 February, 1989. Unpublished report, Centro Internacional de Agricultura Tropical, 18 pp.
- ALLEN, D.J. & SMITHSON, J.B. (1991). The Regional Program on Beans, *Phaseolus vulgaris* L., in Southern Africa : Strategy and Progress in Plant Improvement. In *Crop Genetic Resources of Africa, I* (ed. F. Attere, H. Zedan, N.Q. Ng & P. Perrino), pp. 205-214. IBPGR/IITA/UNEP/CNR.
- ALLEN, D.J. & TRUTMANN, P. (1985). Trip to South-western Uganda, 26-29 May, 1985. Unpublished report, Centro Internacional de Agricultura Tropical, 10 pp.
- ANGUS, A. (1962-66). Plant Pests and Diseases in Zambia, Parts 1-7 and supplement. Mt Makulu Research Station, Zambia (mimeo).
- ANON. (1965-67). Bean. Annual Reports for the Years 1963, 1964 and 1965. Ministry of Agriculture and Natural Resources and the Environment. Port Louis, Mauritius.
- ANON. (1986). Annual Report for 1985, Mauritius Sugar Industry Research Institute. Reduit, Mauritius.
- ARCHIBALD, R.I.M. (1974). Production of seed beans, *Phaseolus vulgaris*, in the lowveld. *Rhodesian agricultural Journal* **71**: 61-64.
- AUTREY, L.J.C. & SAUMTALLY, S. (1990). Identification of components of the bacterial blight complex of bean in Mauritius, their relative importance and measures adopted for their control. In Smithson, J.B. (ed.). Proceedings of the Second Workshop on Bean Research in Eastern Africa, Nairobi, Kenya, 5-8 March 1990. *CIAT Africa Workshop Series No.7* : 236-250.

- AUTRIQUE, A. (1981). Principaux ennemis des cultures de la région des Grands Lacs d'Afrique Centrale. *Publication AGCD*.
- AUTRIQUE, A. & PERREAUX, S. (1989). Maladies et ravageurs des cultures de la région des Grands Lac d'Afrique Centrale. *Publication AGCD No. 24*.
- AYONOADU, U.W.U. (1974). Races of bean anthracnose in Malawi. *Turrialba* **24**: 311-314.
- BOEREMA, G.H. (1972). *Ascochyta phaseolorum* synonymous with *Phoma exigua*. *Netherlands Journal of Plant Pathology* **71**: 79-89.
- BOEREMA, G.H., CRUGER, G., GERLAGH, M. & NIRENBERG, H. (1981). *Phoma exigua* var. *diversispora* and related fungi on *Phaseolus* beans. *Zeitschrift fur Pflanzenkrankheiten und Pflanzenschutz* **88**: 597-607.
- BOS, L. (1974). FAO Working Paper AGP : DP/ETH/71/533.
- BOURIQUET, G. (1946). Maladies du haricot. In *Les maladies des plantes cultivees a Madagascar. Encyclopedie Mycologique* **12**: 462-470. Paul Lechevalier, Paris.
- BURUCHARA, R.A. (1992a). Report on travel to Kenya, 25 September - 02 October 1992. Unpublished report, Centro Internacional de Agricultura Tropical, Rubona, Rwanda, 5 pp.
- BURUCHARA, R.A. (1992b). Report on travel to Madagascar. Unpublished report, Centro Internacional de Agricultura Tropical.
- BURUCHARA, R.A. (1992c). Annual Report Bean Program 1992. Part 1 : Strategic Research. *Draft activity report*. Centro Internacional de Agricultura Tropical, Cali, 343 pp.
- BURUCHARA, R.A. & RUSUKU, G. (1992). Root rot research in the Great Lakes Region. In Buruchara, R.A. & Scheidegger, U.C. (ed.). Proceedings of the Pan-Africa Bean Pathology Working Group Meeting, Thika, Kenya, 26-30 May 1992. *CIAT Africa Workshop Series No. 23* : 49-55.
- BUYCKX, E.J.E. (1962). *Precis des maladies et des insectes nuisibles rencontres sur les plants cultivees au Congo, au Rwanda et au Burundi*. Hors serie, Institut National pour l'Etude Agronomique du Congo (INEAC), 708 pp.
- CAMARADA, C.P. (1989). Angola. In Smithson, J.B. (ed.). Proceedings of a Workshop on Bean Varietal Improvement in Africa, Maseru, Lesotho, 30 January – 02 February, 1989. *CIAT Africa Workshop Series No. 4*, 219-226.
- CIAT (1988). Annual Report Bean Program 1988, *Working Document No. 53*. Centro Internacional de Agricultura Tropical, Cali, 398 pp.
- CURZI, M. (1934). De fungis et morbis africanis. II. De *Pseudomonas plantarum* parasitis *Somaliae*. *Boll. R. Staz. Pat. Veg.*, N.S., XIV, 1, pp. 173-184.
- DADANT, R., RASOLOFO & BAUDIN, P. (1961). *Liste des maladies des plantes cultivees a Madagascar jusqu'en 1960*. Institut de Recherches Agronomiques de Madagascar.
- DIEUDONNÉ, C. (1980). Bean production in Burundi. In *Potential for Field Beans in Eastern Africa*, pp. 31-34, Proceedings of a Regional Workshop held in Lilongwe, 9-14 March 1980, CIAT Series No. 03EB-1.
- DOIDGE, E.M., BOTTOMLEY, A.M., VAN DER PLANK, J.E. & PAUER, G.D. (1953). A revised list of plant diseases occurring in South Africa. *Science Bulletin, Department of Agriculture, Union of South Africa*, No. 346, 122pp.
- DURNEZ, C. (1983). Les maladies du haricot commun (*Phaseolus vulgaris* L.) au Rwanda. Journées d'études du Département des Productions Végétales. *Compte-rendu. Institut des Sciences Agronomiques du Rwanda (ISAR)*, pp. 60-72.
- EBBELS, D.L. & ALLEN, D.J. (1979). A supplementary and annotated list of plant diseases, pathogens and associated fungi in Tanzania. *Phytopathological Papers* **22**: 1-89.
- EDINGTON, B.R. & WHITLOCK, V.H. (1988). Identification of an isolate of bean common mosaic virus from the Transvaal and Natal. *Annals of applied Biology* **113**: 645-648.
- GALVEZ, G.E. & MORALES, F.J. (1989). Whitefly-transmitted viruses. In *Bean Production Problems in the Tropics*, 2nd edn. (ed. H.F. Schwartz & M.A. Pastor-Corrales), pp. 379-406. Cali, Colombia, Centro Internacional de Agricultura Tropical.

- GATUMBI, R.W. (1986). Host lists of Kenya fungi and bacteria (a ten-year supplement : 1971-1980). *East African Agricultural and Forestry Journal* **45**: Special issue.
- GERLAGH, M. (1987). Ascochyta blight: classification, distribution, prevalence, economic importance, survival, spread, prevention and cure. In Smithson, J.B. & Trutmann, P. (ed.) Proceedings of the First African Bean Pathology Workshop, Kigali, Rwanda 14-16 November 1987, CIAT Africa Workshop Series No. 20 : 22-26.
- GOETEYN, R. (1975). Note Technique No. 13. *Institut des Sciences Agronomique du Rwanda*.
- GOETHALS, M. (1986). Les viroses des legumineuses. *Rapport Annuel 1986*, Institut des Sciences Agronomiques du Burundi (ISABU), Bujumbura.
- GOLATO, C. (1967a). Malattie delle Plante Coltivate in Somalia. Instituto Agronomico per L'Oltremare, Firenze.
- GOLATO, C. (1967b). Cercosporiosi del fagiolo in Somalia. *Rivista di Agricoltura Subtropicale e Tropicale* **61**: 159-162.
- GONDWE, B. (1987). Halo blight of *Phaseolus* beans. *Bean Research* 1, 70-74. Proceedings of the Fifth Bean Research Workshop, Sokoine University of Agriculture, Morogoro, 9-12 September 1986.
- GORTER, G.J.M.A. (1977). Index of plant pathogens and the diseases they cause in cultivated plants in South Africa. Department of Agricultural Technical Services.
- GREENBERG, D.C., ALLEN, D.J. & MBEWE, M. (1983). Trip to Northern Zambia and Southern Tanzania. Unpublished report, Msekera Research Station, Ministry of Agriculture.
- GRISLEY, W. (1990). An overview of Bean Production in Sub-Saharan Africa. In *Bean Research* **5**. Progress in Improvement of Common Bean in Eastern and Southern Africa (ed. J.B. Smithson). CIAT Africa Workshop Series No. 12 : 9-19.
- HABISH, H.A. (1972). Aflatoxin in haricot bean and other pulses. *Experimental Agriculture* **8**: 135-137.
- HABTU ASSEFA (1987). Haricot bean diseases and their importance in Ethiopia. *Ethiopia Journal of Agricultural Sciences* **9**: 55-66.
- HABTU ASSEFA (1994). Epidemiology of bean rust in Ethiopia. PhD Thesis. Wageningen Agricultural University.
- HANSFORD, C.G. (1937). Annotated host list of Uganda parasitic fungi and plant diseases, Part III. *East African agricultural Journal* **3**: 79-84.
- HANSFORD, C.G. (1943). Host list of the parasitic fungi of Uganda, Part 11. *East African agricultural Journal* **9**: 50-55.
- HENDRICKX, F.L. (1939). Observations phytopathologiques a la station de Mulungu en 1938. *Rapport Annuel pour l'Exercice 1938, 2 part*, pp. 117-128. Institut National pour l'Etude Agronomique du Congo Belge (INEAC).
- HENDRICKX, F.L. (1948). *Sylloge fungorum congensem*. Catalogue des champignons signales au Conge Belge et au Ruanda-Urundi. Publications a l'Institut National pour l'Etude Agronomiques du Congo Belge (INEAC), Serie Scientifique No. 35, 216 pp.
- HOCKING, D. (1967). A new virulent form of *Phaeoisariopsis griseola* causing circular leaf spot of French bean. *Plant Disease Reporter* **51**: 276-278.
- HOLLIDAY, P. (1980). *Fungus Diseases of Tropical Crops*. Cambridge University Press.
- HOLLINGS, M., PAWLEY, R.R. & BOCK, K.R. (1981). French bean. *Report of the Glasshouse Crops Research Institute for 1979*, 159.
- HOWLAND, A.K. & MACARTNEY, J.C. (1966). East African bean rust studies. *East African Agricultural and Forestry Journal* **32**: 208-210.
- HUBBELING, N. (1973). Report on bean diseases in Kenya: survey of diseases occurring and recommendations about the methods of control. IPO, Wageningen, 27 pp.
- INEAC (1939). *Rapport Annuel pour l'Exercice 1938 1 part*. Institut National pour l'Etude Agronomique du Congo Belge.
- INEAC (1944). *Rapport pour les Exercices 1942 et 1943*. Institut National pour l'Etude Agronomique du Congo Belge.

- INEAC (1949). *Rapport Annuel pour l'Exercice 1948*. Institut National pour l'Etude Agronomique du Congo Belge.
- INEAC (1958a). *Rapport Annuel pour l'Exercice 1956*. Institut National pour l'Etude Agronomique du Congo Belge.
- INEAC (1958b). Haricots. *Rapport Annuel 1958*. Burundi, Station Experimentale de Kisozi. Institut National pour l'Etude Agronomique du Congo Belge.
- INEAC (1958c). Haricots. *Rapport Annuel 1958*. Burundi, Station du Mosso. Institut National pour l'Etude Agronomique du Congo Belge.
- INEAC (1959). *Rapport Annuel pour l'Exercice 1957*. Institut National pour l'Etude Agronomique du Congo Belge.
- ISABU (1983). Legumineuses : le haricot. *Rapport Annuel (Septembre 1982- Aout 1983)*. Institut des Sciences Agronomiques du Burundi, Bujumbura.
- ISABU (1990). *Rapport Annuel*. Institut de Sciences Agronomiques du Burundi, Bujumbura.
- ISAR (1985). *Rapport Annuel 1984*, Programme Legumineuse, Institut des Sciences Agronomique du Rwanda.
- KAISER, W.J. & RAMOS, A.H. (1980). Occurrence of *Pseudomonas syringae* on bean and soybean in Kenya, *Plant Disease* **64**: 593-595.
- KANNAIYAN, J., HACIWA, H.C. & MBEWE, M.N. (1990). Screening bean genotypes for scab resistance in Zambia. In Smithson, J.B. (ed.). Proceedings of the Ninth SUA/CRSP and Second SADC/CIAT Bean Research Workshop, Morogoro, Tanzania, 17-22 September 1990. *CIAT Africa Workshop Series No. 12*: 280- 293.
- KAMANZI, A. (1982). Contribution a L'etude de *Colletotrichum lindemuthianum* (Sacc. & Magn.) Bri. & Cav., agent de l'anthracnose. *Memoire Universite National du Rwanda*, Butare.
- KIDANE AWGICHEW (1982). *Additional index of plant diseases in Ethiopia*. Institute of Agricultural Research, Addis Ababa.
- KINYUA, G.K., MUKUNYA, D.M. & VAN BREUKELEN, E.M. (1981). Studies on genetic resistance of beans to *Pseudomonas phaseolicola* in Kenya. *Proceedings of the Fifth International Conference of Plant Pathogenic Bacteria*, Cali.
- KULKARNI, H.Y. (1973). Notes on East African Plant virus diseases. 4. Bean common mosaic virus. *East African agricultural and Forestry Journal* **39**: 72-76.
- LAMA, L. (1989). Zaire. In Smithson, J.B. (ed.). Proceedings of a Workshop on Bean Varietal Improvement in Africa, Maseru, Lesotho, 30 January - 02 February 1989. *CIAT Africa Workshop Series No.4* : 156-160.
- LEAKEY, C.L.A. (1970). The improvement of beans (*Phaseolus vulgaris*) in East Africa, In Leakey C.L.A. (ed.). *Crop Improvement in East Africa*, pp. 99-128. Commonwealth Agricultural Bureaux, Farnham Royal.
- LEAKEY, C.L.A. (1973). A note on *Xanthomonas* blight of beans (*Phaseolus vulgaris* (L.) Savi) and prospects for its control by breeding for tolerance. *Euphytica* **22**: 132-140.
- LEAKEY, C.L.A. & SIMBWA-BUNNYA, M. (1972). Races of *Colletotrichum lindemuthianum* and implications for bean breeding in Uganda. *Annals of applied Biology* **70**: 25-34.
- LIN, E. (1984). Development, evaluation and utilization of germplasm in Swaziland. In Reunion de Trabajo sobre Ensayos Internacionales de Frijol, pp. 286-304. Centro Internacional de Agricultura Tropical, Cali.
- MABAGALA, R.B. & SAETTLER, A.W. (1992). Races and survival of *Pseudomonas syringae* pv. *phaseolicola* in northern Tanzania. *Plant Disease* **76**: 678-682.
- MATHIAS, P.L. (1971). Pest and disease problems in the Seychelles. *Proceedings Vol.3, 6th British Insecticide and Fungicide Conference*, Brighton, 15-18 November 1971, pp. 835-841.
- MCDONALD, J. (1936). A revised list of plant diseases in Kenya Colony. *East African agricultural Journal* **1**: 463-468.
- MINK, G.I. (1985). *Bean virus serological survey, Kenya and Tanzania*. Unpublished report, Washington State University, Prosser.
- MMBAGA, M.T. & STAVELY, J.R. (1987). Pathogenic specialization in *Uromyces appendiculatus* and rust resistance in Tanzania bean cultivars. *Bean Research I*, 66-69. Proceedings of the Fifth Bean Research Workshop, Sokoine University of Agriculture, Morogoro, 9-12 September, 1986.

- MOHAN, S.K. & HAGEDORN, D.J. (1989). Additional bacterial diseases. In *Bean Production Problems in the Tropics* (ed. H.F. Schwartz & M.A. Pastor-Corrales), pp. 303-319. Cali, Colombia, Centro Internacional de Agricultura Tropical.
- MORALES, F.J. & GAMEZ, R. (1989). Beetle-transmitted viruses. In *Bean Production Problems in the Tropics*, 2nd Edn. (ed. H.F. Schwartz & M.A. Pastor-Corrales), pp. 363-377. Cali, Colombia, Centro Internacional de Agricultura Tropical.
- MSUKU, W.A.B. (1985). Pathogenic variation and virulence of isolates of *Pseudomonas syringae* pv. *phaseolicola* in Malawi. *Annual Report of the Bean Improvement Cooperative* **28**: 125-126.
- MSUYA, D., MUSHI, C.S. & ALLEN, D.J. (1992). Progress in identifying sources of resistance to white mould (*Sclerotinia sclerotiorum*) in beans in Tanzania. *Annual Report of the Bean Improvement Cooperative* **35**: 122-123.
- MUKOKO, O.Z. (1992). Breeding beans (*Phaseolus vulgaris* L.) for resistance to bean common mosaic virus in Zimbabwe. PhD Thesis, University of Cambridge.
- MUKUNYA, D.M., MUTHANGYA, P.M. & ESELE, J.P. (1981). Bacterial blights of beans (*Phaseolus vulgaris* L.) in Kenya. *Proceedings of the Fifth International Conference of Plant Pathogenic Bacteria*, Cali.
- MUNIMBAZI, C. & PERREAUX, D. (1986). Screening for resistance to common and halo bacterial blights. *Phaseolus Beans Newsletter for Eastern Africa* **5**: 5-6.
- MUNYEMANA, A. (1987). Contribution à l'étude de *Ascochyta phaseolorum* (Sacc.), agent de l'ascochytose du haricot au Rwanda. *Mémoire Université National du Rwanda*, Butare.
- MUTITU, E.W. (1979). Etiology, epidemiology and control of bean caused by *Elsinoe phaseoli* Jenkins f.sp. *vulgare* in Kenya. MSc. Thesis, University of Nairobi.
- MUTITU, E.W., MUKUNYA, D.M. & GATHURU, E. (1983). Fusarium root rot on beans caused by *Fusarium solani* f.sp. *phaseoli* in Kenya. In Webster, B.D. (ed.). *Improvement of drought and heat tolerance of disease resistant beans in semi-arid regions of Kenya*. Report of Research, 1982-83. East Lansing, Bean- Cowpea Collaborative Research Support Program, pp. 74-94.
- MUTITU, E.W., MUKUNYA, D.M. & KEYA, S.O. (1988). Biological control of fusarium yellows on beans caused by *Fusarium oxysporum* Schl. f.sp. *phaseoli* Kendrick & Snyder using organic amendments locally available in Kenya. *Acta Horticulturae* **218**: 267-274.
- NATTRASS, R.M. (1961). Host lists of Kenya fungi and bacteria. *Mycological Papers* **81**: 1-46.
- NEERGAARD, P. (1977). Quarantine policy for seed in transfer of genetic resources. In *Plant Health and Quarantine in International Transfer of Genetic Resources* (ed. W.B. Hewitt & L. Chiarappa), pp. 309-314. Cleveland, USA, CRC Press.
- NEVELING, C.H. (1956). Report of the Secretary for Agriculture for the year ended 31 August 1955. *Farming in South Africa* **31**: 359.
- NJUGUNAH, S.K., NDEGWA, A.M.M., VAN RHEENEN, H.A. & MUKUNYA, D.M. (1980). Bean production in Kenya. In *Potential for Field Beans in Eastern Africa*. Proceedings of a Regional Workshop, Lilongwe, Malawi. 9-14 March 1980. CIAT, pp. 35-53.
- NKEZABERA, M. (1987). Etude de la pathogénicité des souches de *Colletotrichum* du Rwanda à l'égard de certains variétés de haricot. *Mémoire, Université National du Rwanda*, Butare.
- NYABYENDA, P., SEKANABANGA, C. & NYANGURUNDI, L. (1981). Bean production in Rwanda. In *Potential for Field Beans in Eastern Africa*, pp. 99-121. Centro Internacional de Agricultura Tropical, Cali.
- OMUNYIN, M.E. (1979). Screening for resistance to bean common mosaic virus in food beans (*Phaseolus vulgaris* L.). *Proceedings of a Symposium on Grain Legume Improvement in Eastern Africa*, Nairobi, 20-24 August, 1979.
- ONDIEKI, J.J. (1973). Host lists of Kenya fungi and bacteria (a ten-year supplement 1961-1970). *East African agricultural and Forestry Journal* **28**: spec. suppl., 31 pp.
- OPIO, A.F. (1993). Pathogenic variation, seed transmission and survival of (*Xanthomonas campestris* pv. *phaseoli*) in Uganda. PhD Thesis, Sokoine University of Agriculture, Morogoro.

- ORIEUX, L. & FELIX, S. (1968). List of plant diseases in Mauritius. *Phytopathological Papers* 7: 1-48.
- OWERA, S.A.P. (1990). Analysis of serotypes and strains of bean common mosaic virus (BCMV) in countries within CIAT's Regional Programme on Beans in Eastern Africa. In Smithson, J.B. (ed.). Proceedings of the Second Workshop on Bean Research in Eastern Africa, Nairobi, Kenya, 5-8 March 1990. *CIAT Africa Workshop Series No. 7* : 63-74.
- PEREGRINE, W.T.H. (1971). A preliminary note on chemical control of bean anthracnose in Malawi. *Pest Articles and News Summaries* 17: 47-50.
- PEREGRINE, W.T.H. & SIDDIQI, M.A. (1972). A revised and annotated list of plant diseases in Malawi. *Phytopathological Papers* 16: 1-51.
- PERREAUX, D. (1986). Ecology and importance of bean diseases in Burundi. *Phaseolus Beans Newsletter for Eastern Africa* 5: 4-5.
- PERREAUX, D., BAGAMBAKE, D., GAHUNGU, E., MIKOKORO, C., NKUBAYE, E. & WAKANA, E. (1986). Epidemiologie des maladies du haricot en milieu rural et pertes de rendement. *Rapport Annuel (Septembre 1985 -Aout 1986)*. Institut des Sciences Agronomiques du Burundi (ISABU), Bujumbura.
- PHILLIPS, A.J.L. (1994). Occurrence of scab of *Phaseolus vulgaris* caused by *Elsinoë phaseoli* in South Africa. *Plant Pathology* 43: 417-419.
- PLUMB-DHINDSA, P. & MONDJANE, A.M. (1984). Index of Plant Diseases and Associated Organisms of Mozambique. *Tropical Pest Management* 30: 407-429.
- PRICE, M. & CISAHAYO, D. (1986). Ascochyta blight on major food crops in Rwanda, East Africa. *Plant Disease* 70: 1159 (Abstr.).
- PYNDJI, M.M. (1988). Progres de recherche sur la maladie des taches anguleuses du haricot commun dans la region des Grands Lacs. In Pyndji, M., Davis, J. & Scheidegger, U. (ed.). *Actes du Quatrieme Seminaire Regional sur l'Amelioration du Haricot dans la Region des Grands Lacs*, Bukavu, Zaire, 21-25 November 1988. *CIAT Africa Workshop Series No. 9* : 39-43.
- RASOLOFO-RAZAFINDRAMAMBA & RALIARISON, V.A. (1976). *Liste des maladies des plantes cultivees a Madagascar de 1961 a 1971*. Centre National de la Recherche Appliquee au Developpement Rural (CENRADERU-FOFIFA).
- RILEY, E.A. (1956). A preliminary list of plant diseases in Northern Rhodesia. *Mycological Papers* 63: 1-28.
- RILEY, E.A. (1960). A revised list of plant diseases in Tanganyika Territory. *Mycological Papers* 75: 1-42.
- ROJOA, H. (1989). Mauritius. In Smithson, J.B. (ed.). Proceedings of a Workshop on Bean Varietal Improvement in Africa, Maseru, Lesotho, 30 January - 02 February 1989. *CIAT African Workshop Series No. 4* : 128-132.
- ROTHWELL, A. (1983). A revised list of plant diseases occurring in Zimbabwe. *Kirkia* 12, II : 233-351.
- RUSUKU, G. (1991). Incidence des maladies a pourritures des racines du haricot au Rwanda. In Nyabyenda, P. & Scheidegger, U. (ed.). *Actes du Sixieme Seminaire Regional sur L'Amelioration du Haricot dans la Region des Grands Lacs*, Kigali, Rwanda, 21-25 January 1991. *CIAT Africa Workshop Series No.17* : 92-103.
- SABET, K.A. (1959). Studies on the bacterial diseases of Sudan crops. III. On the occurrence, host range and taxonomy of the bacteria causing leaf blight diseases of certain leguminous plants. *Annals of applied Biology* 47: 318-331.
- SABET, K.A., ISHAG, F. & KHALIL, O. (1969). Studies on the bacterial diseases of Sudan crops. VII. New records. *Annals of applied Biology* 63: 357-369.
- SALIH, S.H. & SALIH, F.A. (1985). Bean production in the Sudan. In Potential for Field Beans (*Phaseolus vulgaris* L.) in West Asia and North Africa, pp. 87-100. Proceedings of a Regional Workshop, Aleppo, Syria, 21-23 May, 1983. Centro Internacional de Agricultura Tropical, Cali.
- SCHWARTZ, H.F. (1989). Additional fungal pathogens. In *Bean Production Problems in the Tropics*, 2nd edn. (ed. H.F. Schwartz & M.A. Pastor-Corrales), pp. 231-259. Cali, Colombia, Centro Internacional de Agricultura Tropical.
- SENGOOBA, T.N.K. (1994). Identification of alternative natural hosts of bean common mosaic virus and

- assessment for resistance in 'African' bean genotypes. PhD Thesis, Sokoine University of Agriculture, Morogoro.
- SERAFIM, F.J.D & SERAFIM, M.C. (1968). Lista das Doencas de Culturas de Angola. Instituto de Investigacao Agronomica de Angola. Serie Tecnica, No.2. 16 pp.
- SHEFFIELD, F.M.L. (1968). Closed quarantine procedures. *Review of applied Mycology* **47**: 1-8.
- SILBERNAGEL, M.J., MILLS, L.J. & WANG, W.-Y. (1986). Tanzanian strains of bean common mosaic virus. *Plant Disease* **70**: 839-841.
- SMITHSON, J.B. (1986). Tour Report to Ethiopia, August 1986. Unpublished report, Centro Internacional de Agricultura Tropical.
- SMITHSON, J.B. (1988). Travel Report : Lesotho, 7-14 February, 1988. Unpublished report. Centro Internacional de Agricultura Tropical. 25 pp.
- SOHATI, P., SITHANANTHAM, S., KANNAIYAN, J., HACIWA, H.C. & KABUNGO, D.A. (1992). A review of the SADC/CIAT aphid/BCMV sub-project in Zambia and Tanzania. In Allen, D.J. (ed.). Proceedings of the Third Regional Workshop on Bean Research in Southern Africa, Mbabane, Swaziland, 5-7 October 1992. *CIAT Africa Workshop Series*.
- SPENCE, N.J. (1992). The identification, distribution and ecology of bean common mosaic virus in Africa. PhD Thesis, University of Birmingham.
- SPENCE, N.J. & WALKEY, D.G.A. (1991). Identification of strains of bean common mosaic virus occurring in different regions of Africa. *Annual Report of the Bean Improvement Cooperative* **34**: 5-6.
- STEWART, R.B. & DAGNATCHEW YIRGOU (1967). *Index of plant diseases in Ethiopia*. Haile Selassie I University College of Agriculture Experimental Station Bulletin No. 30, 95 pp.
- STEYAERT, R.L. (1948). Contribution a l'étude des parasites des vegetaux du Congo Belge. *Bulletin de la Societe Royale de Botanique de Belgique* **80**: 11-58.
- STOETZER, H.A.I. & WAITE, B.H. (1984). Evaluation of bean diseases in the Grain Legume Project Trials, Katumani Research Station. National Horticultural Research Station, Thika, Kenya (mimeo).
- TARR, S.A.J. (1955). *The fungi and plant diseases of the Sudan*. Commonwealth Mycological Institute, Kew, 127 pp.
- TARR, S.A.J. (1963). A supplementary list of Sudan fungi and plant diseases. *Mycological Papers* **85**: 1-31.
- TEVERSON, D.M. (1991). Genetics of pathogenicity and resistance in the halo-blight disease of beans in Africa. PhD Thesis, University of Birmingham.
- TRENCH, T.N., ROUX, C. & MELIS, R.J.M. (1986). Recent records of *Phoma exigua* var. *exigua* in Natal. *Phytophylactica* **18** (1), abstr.
- VAN RHEENEN, H.A. & MUIGAI, S.G.S. (1984). Control of bean common mosaic by deployment of the dominant I gene. *Netherlands Journal of Plant Pathology* **90**: 85-94.
- VETTEN, H.J. & ALLEN, D.J. (1991). Recent progress in the identification of viruses of *Phaseolus vulgaris* in Africa. *Annual Report Bean Improvement Cooperative* **34**: 3-4.
- WALLACE, G.B. (1941). Yellow bean mosaic and notes on other bean diseases. *East African agricultural Journal* **7**: 114-115.
- WALLACE, G.B. (1944). Plant Pathology. Specialist research work of the Department of Agriculture, Tanganyika Territory, 1943. *Review of applied Mycology* **24**: 93.
- WALLACE, G.B. & WALLACE, M.M. (1949). A list of plant diseases of economic importance in Tanganyika Territory. *Mycological Papers* **26**: 1-26.
- WALLACE, G.B. & WALLACE, M.M. (1953). A supplement to a list of plant diseases of economic importance in Tanganyika Territory. *Mycological Papers* **51**: 1-7.
- WHITESIDE, J.O. (1966). Revised list of plant diseases in Rhodesia. *Kirkia* **5**: 87-196.
- WIEHE, P.O. (1948). The plant diseases and fungi recorded from Mauritius. *Mycological Papers* **24**: 1-39.
- ZAUMEYER, W.J. & THOMAS, H.R. (1948). Shiny pod (greasy pod) virus and its identity with black root virus. *Phytopathology* **38**: 29 (Abstr.).
- ZAUMEYER, W.J. & THOMAS, H.R. (1957). A monographic study of bean diseases and methods for their control. *Technical Bulletin U. S. Department of Agriculture* **868**: 2nd edn.