



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Agriculture, Ecosystems and Environment xxx (2005) xxx–xxx

**Agriculture  
Ecosystems &  
Environment**
[www.elsevier.com/locate/agee](http://www.elsevier.com/locate/agee)

## Resource flows and nutrient balances for crop and animal production in smallholder farming systems in eastern Uganda

A.O. Esilaba<sup>a,\*</sup>, P. Nyende<sup>c</sup>, G. Nalukenge<sup>a</sup>, J.B. Byalebeka<sup>b</sup>,  
R.J. Delve<sup>c</sup>, H. Ssali<sup>b</sup>

<sup>a</sup> *Centro Internacional de Agricultura Tropical (CIAT), Kawanda, P.O. Box 6247, Kampala, Uganda*

<sup>b</sup> *National Agricultural Research Organisation (NARO), Kawanda Agricultural Research Institute, P.O. Box 7065, Kampala, Uganda*

<sup>c</sup> *Tropical Soil Biology and Fertility Institute of CIAT, P.O. Box 30592, Nairobi, Kenya*

Received 16 June 2003; received in revised form 18 March 2005; accepted 23 March 2005

### Abstract

Resource flow models are useful tools that assist farmers in analysing their soil fertility management strategies and in planning, experimenting and adapting ways to improve the use of scarce local resources. Resource flows and farm nutrient balance studies were carried out in eastern Uganda to ascertain the movement of organic resources and nutrients in and out of the farm system during a participatory learning and action research (PLAR) process. The resource flows were transformed into nutrient flows and partial nutrient balances were calculated using the Resource Kit computer package. Results of a farmers' soil fertility management classification at the start of the PLAR intervention in 1999 revealed that 3% of the farmers were good soil fertility managers (class I), 10% were average soil fertility managers (class II) and 87% were poor soil fertility managers (class III). The results indicate that the net farm nutrient balances in  $\text{kg ha}^{-1}$  per season for all the nutrients [nitrogen (N), phosphorus (P), and potassium (K)] were negative for both the good and the poor soil fertility managers. Class I farm balances irrespective of the season, were however more negative than those of class 3 farms. For the long rains seasons (LR 2000, 2001 and 2002), the average net farm nutrient balances for N, P, and K for class I farms were  $-5.0$ ,  $-0.6$  and  $-8.0 \text{ kg ha}^{-1} \text{ year}^{-1}$ , while for the short rains seasons (SR 2000 and 2001), the nutrient balances were  $-3.5$ ,  $-0.5$  and  $-6.0 \text{ kg ha}^{-1} \text{ year}^{-1}$ , respectively. For the class III farms, the average net farm nutrient balances for N, P, and K in the long rain seasons (LR 2000, 2001 and 2002) were  $-3.3$ ,  $-0.3$  and  $-4.0 \text{ kg ha}^{-1} \text{ year}^{-1}$  while for the short rains seasons (SR 2000 and 2001), the nutrient balances were  $-3.5$ ,  $0.5$  and  $-5.0 \text{ kg ha}^{-1} \text{ year}^{-1}$ , respectively. Soil management interventions for these small-scale farmers should aim at reversing nutrient depletion with a focus on profitable management of the crop production system, which is the major cause of nutrient depletion.

© 2005 Published by Elsevier B.V.

**Keywords:** Farm classification; Farming systems; Nutrient balances; Resource flows; Soil fertility; Eastern Uganda

\* Corresponding author. Present address: KARI Headquarters, Kenya Agricultural Research Institute, P.O. Box 57811, City Square, 00200 Nairobi, Kenya. Tel.: +254 20 583301x20; fax: +254 20 583344.

E-mail addresses: [aoesilaba@kari.org](mailto:aoesilaba@kari.org), [aesilaba@yahoo.co.uk](mailto:aesilaba@yahoo.co.uk) (A.O. Esilaba).

## 35 1. Introduction

36 Soil nutrient balance studies in Africa show  
 37 evidence of widespread nutrient mining leading to  
 38 severe nutrient deficiencies across ecological zones.  
 39 Soil nutrient stocks are not static entities and studies in  
 40 different parts of Africa at different spatial scales show  
 41 that nutrients are being depleted at alarming rates  
 42 (Stoorvogel and Smaling, 1990; Van der Pol, 1992;  
 43 Smaling et al., 1993, 1997; Smaling and Braun, 1996;  
 44 Scoones, 2001). Nutrients are annually taken away in  
 45 crops or lost in processes such as leaching and erosion  
 46 which far exceed the nutrient inputs through  
 47 fertilisers, deposition and biological fixation (Smaling  
 48 and Braun, 1996). Nutrient mining has been estimated  
 49 to average 660 kg of nitrogen (N), 75 kg of  
 50 phosphorus (P) and 450 kg of potassium (K) per  
 51 hectare per year during the last 30 years from about  
 52 200 million hectares of cultivated land in 37 countries  
 53 in Africa (Stoorvogel and Smaling, 1990; Sanchez  
 54 et al., 1997; Smaling et al., 1997). Losses of 130 kg N,  
 55 5 kg P and 25 kg K ha<sup>-1</sup> per year have been reported  
 56 in the East African highlands (Smaling et al., 1997).  
 57 Wortmann and Kaizzi (1998) estimated nutrient  
 58 balances for small-scale farming systems in eastern  
 59 and central Uganda to be negative for all crops except  
 60 for nitrogen (N) and phosphorus (P) in the banana-  
 61 based land use type (LUT).

62 The concern for soil nutrient depletion and low soil  
 63 fertility has led to the development of several  
 64 integrated soil fertility management technologies that  
 65 offer potential for improving soil fertility management  
 66 in Africa (Scoones and Toulmin, 1999). These include  
 67 improved soil erosion control using living barriers or  
 68 micro-catchments, inoculation of grain legumes for  
 69 improved N-fixation, efficient use of manure and other  
 70 locally available organic materials, use of green  
 71 manure and cover crops (Delve and Jama, in press)  
 72 and use of low levels of N and P fertilisers on maize  
 73 (*Zea mays*) and beans (*Phaseolus vulgaris*) (Wort-  
 74 mann et al., 1998; Wortmann and Kaizzi, 1998) in  
 75 eastern Uganda. However, there has been limited  
 76 uptake of these “improved” INM practices.

77 There are a limited number of long-term studies  
 78 monitoring the nutrient status of soils, nutrient  
 79 balances, and crop productivity in Uganda (Bekunda  
 80 et al., 1997; Swift et al., 1994). It is important to  
 81 calculate and monitor nutrient flows to quantify the

82 impact of integrated nutrient management (INM)  
 83 systems on soil fertility and sustainable agricultural  
 84 productivity (Smaling and Braun, 1996; Defoer et al.,  
 85 2000). Monitoring of nutrient stocks and flows is a tool  
 86 for assessing the degree of nutrient mining in an agro-  
 87 ecosystem. When applied to systems where INM  
 88 practices are being introduced, nutrient monitoring  
 89 can be used to assess the effects of INM strategies on  
 90 soil nutrient stocks and flows (Van den Bosch et al.,  
 91 1998).

92 Improved soil nutrient management is important for  
 93 maintaining and improving soil productivity in Uganda  
 94 and strategies are required that more closely address  
 95 farmer requirements and priorities (Deugd et al., 1998).  
 96 In this study resource/nutrient flows were used to work  
 97 with farmers to better understand their current practice,  
 98 their constraints and their opportunities for reversing  
 99 nutrient depletion. Therefore, the objectives of this  
 100 study were to determine resource flows and estimate  
 101 nutrient balances in three different farm typologies and  
 102 to investigate if improved soil fertility management  
 103 impact on sustaining agricultural productivity on the  
 104 smallholder farms in eastern Uganda.

## 2. Materials and methods

### 2.1. Characteristics of the farming system

107 A study was carried out in three villages of  
 108 Magada, Mayuge and Buyemba in Imanyiro sub-  
 109 county of Mayuge District in eastern Uganda. This  
 110 area is located at 0° 35'N, 32°29' E and lies at an  
 111 altitude of 1070–1161 m.a.s.l (meters above sea level)  
 112 covering an area of about 11,113 km<sup>2</sup>. The area has a  
 113 bimodal rainfall pattern varying from 1250 to  
 114 2200 mm (average 1345 mm for 22 years) per annum.  
 115 The first rains (long rain (LR) season) occur between  
 116 March and June and the second rains (short rain (SR)  
 117 season) between August and December. The soils are  
 118 reddish brown sandy loams and sandy clay loams  
 119 (Harrop, 1970) and classified as Orthic Ferralsols  
 120 (FAO, 1977). Most soils in the area have an average  
 121 organic matter content of 11.0–31.0 g kg<sup>-1</sup> but are  
 122 deficient in N and P (Fischler, 1997; Wortmann and  
 123 Kaizzi, 1998).

124 The farming systems show a high degree of  
 125 biological and agronomic diversity and complexity.

126 Average farm size is 1.8–2.0 ha and 90% of the  
 127 farmers are the sole owners of the land. The main  
 128 crops grown in the area are bananas (*Musa spp.*),  
 129 maize, cassava (*Manihot esculenta*), beans, coffee  
 130 (*Coffea canephora*), fruits, vegetables and sweet  
 131 potato (*Ipomea batatas*) (Esilaba et al., 2001b;  
 132 Woelcke and Berger, 2002). The majority of the  
 133 farms have few or no livestock and the mean numbers  
 134 are 1.5 local cows, 0.2 improved cows, 1.7 goats or  
 135 sheep, 0.9 pigs and 12.0 chickens per farm (Wortmann  
 136 et al., 1998; Wortmann and Kaizzi, 1998).

### 137 2.2. The PLAR process

138 A Participatory Learning and Action Research  
 139 (PLAR) process (Defoer et al., 2000) was initiated in  
 140 September 1999 in Imanyiro sub-county, Mayuge  
 141 District. The PLAR process comprises four phases:  
 142 diagnosis and analysis, planning, implementation and  
 143 evaluation. During the diagnostic phase of the PLAR  
 144 process, farmers analysed soil fertility management  
 145 diversity and resource endowment of farms in  
 146 Buyemba, Mayuge and Magada villages (Esilaba  
 147 et al., 2001b). The soil fertility management diversity  
 148 classification were standardised into three categories  
 149 (good, average, and poor managers) using the farmers'  
 150 criteria and were attributed to: (1) use of fertilisers  
 151 (both organic and inorganic), (2) use of soil erosion  
 152 control measures, such as vetiver grass strips, terracing  
 153 and mulching, (3) use of green manure, such as  
 154 mucuna, canavalia, crotalaria and lablab, (4) leaving  
 155 land to fallow and (5) use of agroforestry technologies.  
 156 Farms/households using four or more of these measures  
 157 were considered “good” (class I). Farms using one to  
 158 three measures were considered “average” (class II),  
 159 while those farms not using any of these measures were  
 160 considered “poor” (class III).

161 Twenty farmers representing the three soil fertility  
 162 management classes in the three villages were selected  
 163 as test farmers for intensive monitoring, on-farm  
 164 experimentation and resource flow mapping. Soil  
 165 samples (topsoils at 0–20 cm depth) were collected  
 166 from the proposed on farm experimental sites on 28  
 167 farmers' plots for laboratory analysis according to  
 168 methods by Foster (1971) and Okalebo et al., (1993)  
 169 for available Bray-P. Organic matter was determined  
 170 using an oxidation procedure derived from the  
 171 Walkley and Black method as described by Jackson

Table 1

Soil chemical and physical properties (top soils 0–20 cm depth) of typical farms in the three soil fertility management classes<sup>a</sup>

Parameter	Class I (n = 9)	Class II (n = 5)	Class III (n = 14)
pH (water)	5.5	5.2	5.0
Organic matter (g kg <sup>-1</sup> )	28.4	29.4	21.6
N (g kg <sup>-1</sup> )	1.0	1.0	1.0
P (Bray P-2, mg kg <sup>-1</sup> )	4.6	3.3	1.9
K (cmol kg <sup>-1</sup> )	24.6	22.6	15.1
Na (cmol kg <sup>-1</sup> )	6.2	5.7	3.9
Ca (cmol kg <sup>-1</sup> )	53.6	51.6	35.2
Mg (cmol kg <sup>-1</sup> )	29.7	27.6	19.2
Sand (g kg <sup>-1</sup> )	619	642	711
Clay (g kg <sup>-1</sup> )	272	266	201
Silt (g kg <sup>-1</sup> )	109	92	81

<sup>a</sup> Class I = good soil fertility managers; class II = average soil fertility managers; class III = poor soil fertility managers.

(1958). Total nitrogen (N) was determined by the  
 standard Kjeldahl procedure. The extracting solution  
 used for calcium (Ca), Magnesium (Mg), sodium (Na)  
 and potassium (K) was hydrolyzed lactic acid in  
 ammonia solution based on Egner's extracting  
 solution (Foster, 1971). Data on soil properties for  
 farms in the three soil management classes are  
 presented in Table 1. Data analysis for analysis of  
 variance (ANOVA) on soil properties was conducted  
 using SAS (SAS, 1990) and SPSS (SPSS, 2002) for  
 cross tabulation of soil fertility management classes  
 with resource endowment parameters that included  
 wealth ranking (Esilaba et al., 2001a,b).

### 2.3. Resource flow-mapping

Resource flow maps were used to visualise the  
 farmers' soil fertility management situation of the  
 farm during of the PLAR process. The selected  
 farmers drew resource flow maps (RFMs) to visualise,  
 plan and analyse their current, planned and imple-  
 mented soil fertility management practices and to  
 identify possible improvements at the beginning and  
 end of each season. Test farmers from the three soil  
 fertility management diversity classes drew resource  
 maps indicating the different elements of their farms,  
 including fields, food stores, livestock shelters and  
 compost pits (Budelman and Defoer, 2000a,b). The  
 current and preceding crops were noted for each field  
 and farmers drew arrows to show the flows of  
 resources entering and leaving the farm as well as




Fig. 1. Typical resource flow map.

201 flows between fields and other farm components  
 202 (Fig. 1).

203 Farmers estimated the quantity of resources using  
 204 simple local units of measurement [such as tins (1–  
 205 10 kg containers), debes (20 kg containers) etc.] and  
 206 labelled the direction of flow of the resources  
 207 accordingly using appropriate arrows and symbols.  
 208 Similar information was recorded on several recording  
 209 forms relating to (1) farm level data, (2) field level data  
 210 and to various flows, (3) resources leaving the fields

(produce and crop residues); (4) resources entering the  
 211 fields (fertilisers); (5) resources leaving the household  
 212 and animal production system; (6) resources entering  
 213 the household and animal production system and fed  
 214 into a computer using the Resource Kit software for  
 215 analysis (Defoer et al., 2000; Defoer and Budelman,  
 216 2000).  
 217

Nutrient flow analysis was used in evaluating land  
 218 use, the relative intensity of cropping, the ratios  
 219 between inputs and outputs and comparing systems  
 220



Table 2  
Soil fertility management classification (diversity) continuum over three years<sup>a</sup>

Village	Number of farmers							
	Class I		Class II		Class III		Total	
	1999	2002	1999	2002	1999	2002	1999	2002
Buyemba	7	10	19	35	165	153	191	198 <sup>b</sup>
Mayuge	8	10	28	18	94	104	131	132
Magada	4	4	8	51	235	192	247	247
Total	20	24	55	104	494	449	569	577

<sup>a</sup> Class I = good soil fertility managers; class II = average soil fertility managers; class III = poor soil fertility managers.

<sup>b</sup> Other new farmers had joined the village hence the increase in the number.

221 along these lines (Budelman and Defoer, 2000a). The  
222 unit of analysis was the farm system, which is part of  
223 the village land use system (consisting of several  
224 farms and communally used resources). There are  
225 three sub-systems within the farm: the crop production  
226 system (CPS), the animal production system (APS),  
227 and the household system (HHS). For each of the sub-  
228 systems, links with the elements outside the farm  
229 system are presented as flows entering (i.e. IN) the  
230 farm from outside, and as flows leaving (i.e. OUT) the  
231 farm. Links between the sub-systems (i.e. INT) of the  
232 farm are identified and refer to internal flows (Defoer  
233 et al., 1998; Defoer and Budelman, 2000). After five  
234 seasons of experimentation and resource flow map-  
235 ping, farmers evaluated themselves to establish a  
236 continuum as to whether they had moved from one soil  
237 fertility management class to another or remained in  
238 the same class and establish factors that led to these  
239 scenarios (Table 2).

### 240 3. Results and discussion

#### 241 3.1. Soil fertility management diversity 242 classification

243 The results indicate that out of a total of 569  
244 households only 20 (3.5%) were in class I, 55 (10%) in  
245 class II and the majority (494 or 87%) were in class III  
246 (Table 2). Therefore, most farmers were not carrying  
247 out any improved soil fertility management practices,  
248 despite the previous National Agricultural Research  
249 Organisation (NARO), Uganda and the International

Centre for Tropical Agriculture (CIAT) work in this  
250 area (Fischler, 1997; Wortmann et al., 1998; Wort-  
251 mann and Kaizzi, 1998). When key informants from  
252 the farmer groups were asked to reclassify the farmers  
253 after two years they indicated that 10% of the farmers  
254 from the 1999 class III classification had moved to  
255 class I and II. Some farmers in class I had to be  
256 relegated to class II and III (Table 2). The PLAR  
257 process had more impact on farmers in the newest  
258 research village (Magada).  
259

Wealth rankings (Grandin, 1988) obtained during  
260 the diagnostic phase of the PLAR process (Esilaba  
261 et al., 2001b) were compared with data from the soil  
262 fertility management diversity classification for the  
263 same households in the three villages. The wealth  
264 ranks were standardised into four categories (wealthy,  
265 average, poor and very poor) for correlation with soil  
266 fertility management classes using SPSS (Table 3).  
267 The majority of the respondents (74%) were in soil  
268 fertility class III while class II and I each had 13%,  
269 respectively. There were trends indicating a relationship  
270 between wealth ranks and soil fertility management  
271 classes. Seventy five percent (75%) of the farmers in  
272 soil fertility management class I were wealthy, another  
273 25% average, and none were poor or very poor. For class  
274 2 farms, 67% of the farmers were average, 33% were  
275 very poor and none was wealthy or poor. In class 3, 7%  
276 of the farm households were wealthy, 23% average,  
277 31% were poor and 39% were very poor.  
278

No significant differences ( $P < 0.05$ ) were observed  
279 in the soil chemical and physical properties among the  
280 three soil fertility management classes despite farmers'  
281 assessment (Table 1). The soil pH was generally  
282

Table 3  
Relationship between soil fertility management classification and wealth ranking<sup>a</sup>

Soil fertility management Class <sup>b</sup>	Soil fertility management		Wealth rank <sup>c</sup>			
	Number of farmers		Number of farmers (%)			
	1999	2002	Wealthy	Average	Poor	Very poor
Class I	20 (3)	24 (4)	75	25	0	0
Class II	55 (10)	104 (18)	0	67	0	33
Class III	494 (87)	449 (78)	7	23	31	39

<sup>a</sup> Values in parentheses are the percentage of farmers.

<sup>b</sup> Class I = good soil fertility managers; class II = average soil fertility managers; class III = poor soil fertility managers.

<sup>c</sup> Wealth ranks classified according to the farmers criteria into four classes (wealthy, average, poor and very poor).

283 favourable except on farms in class III, which had a low  
 284 average pH. Total (Kjeldahl) N, soil organic matter  
 285 (SOM), available P are inadequate while exchangeable  
 286 K is relatively adequate (Foster, 1971, 1973; Landon,  
 287 1984). Critical values for soil pH, organic matter, total  
 288 N and K in Uganda are 5.2, 30.0 g kg<sup>-1</sup>, 1.8 g kg<sup>-1</sup>,  
 289 5 mg kg<sup>-1</sup> and 13.3 cmol kg<sup>-1</sup>, respectively (Foster,  
 290 1971). The soil textural class at all sites was dominantly  
 291 loam.

292 3.2. Resource flow mapping and resource flows

293 The resource flow mapping exercise was used as a  
 294 learning tool for the researchers, extension staff and  
 295 farmers to visualise the farm system and its  
 296 subsystems, the flow of resources within the farm  
 297 systems and outside the farm system. The complexity  
 298 of the flows within and outside the farm system was  
 299 evident from the exercise. The farmers together with  
 300 the extension staff were able to examine the quantities  
 301 and direction of the flows of the main agricultural  
 302 resources and possible options to minimise losses and  
 303 concentrate resources in key areas. The results show  
 304 that farmers using different soil fertility management  
 305 measures varied also in terms of resource endowment,  
 306 for example, farm size, land tenure, livestock own-  
 307 ership, off-farm employment and farm structures. The  
 308 average class I total farm size was considerably larger  
 309 (3.3 ha) than class III (1.4 ha) with the implication that  
 310 class I farmers can therefore leave more land under  
 311 fallow (0.7 ha) than class 3 farmers (0.2 ha) to restore  
 312 soil fertility. Most of the maps were characterised by  
 313 one field but with many plots (1–20) and of different  
 314 sizes (Fig. 1). The plot sizes ranged from 0.125 to 2 ha.  
 315 Farmers in this area divide their land into many plots  
 316 because of the need to distribute the risk of crop failure  
 317 by growing a variety of crops. The fields/plots were  
 318 divided, according to the crop growing or intended to  
 319 be grown for that particular season.

320 For all the farm classes, most of the resources within  
 321 the farm system came from crop fields (crop production  
 322 system) into the household system (CPS-HHS) as food,  
 323 and out of the farm system (OUTcps) as sale of surplus  
 324 food (Figs. 2–4). On the other hand, very limited  
 325 resources were returned to the farm and to the crop  
 326 production system (INcps). There were no seasonal  
 327 differences in the direction of flow of the resources but  
 328 there was for the quantities of resource flows.




Fig. 2. Net partial farm balances and partial crop production system (CPS) balances for class I (2000–2002) and III (2000–2002) farms. Net partial farm balances = INcps + INaps + INhhs – OUTcps – OUTaps – OUThhs; Partial crop production system (CPS) balances = INcps + INTaps – cps + INT hhs – cps – OUTcps – INTcps – aps – INTcps – hhs; where IN = input flow; OUT = output flow; INT = internal flow; aps = animal production system; cps = crop production system; hhs = household system; LR = long rains; SR = short rains.

Food crops are either consumed immediately when they come from the field or are temporarily stored for food or sale to the market. However, these are subsistence farmers and therefore most of the produce




Fig. 3. Average N, P and K balances per hectare per season for all locations of typical class I farms for the crop production system (CPS), animal production system (APS), household system (HHS) and out of the farm system (OFS) over five seasons in the 2000, 2001 and 2002 long rains (LR) and 2000 and 2001 short rains (SR).




Fig. 4. Average N, P and K balances per hectare per season for all locations of typical class III for the crop production system (CPS), animal production system (APS), household system (HHS) and out of the farm system (OFS) over five seasons in the 2000, 2001 and 2002 long rains (LR) and 2000 and 2001 short rains (SR).

333 (80%) is consumed on the farm, with the exception of  
 334 coffee. Intercropping is practised to reduce on labour  
 335 demands, maximise land use and reduce risk due to  
 336 drought. Apart from land allocation for crops, there is  
 337 no significant difference in the crops cultivated, crop  
 338 pattern (sole versus intercropping), yields, crop  
 339 residue management and general crop husbandry  
 340 between the two farm classes, despite farmers in class  
 341 I being known to be ‘good’ soil managers. There are  
 342 also no major differences in yield, crop types and size  
 343 of land allocation between the long rains and short rain  
 344 seasons. Nutrients are exported from the farms mainly  
 345 through coffee, food crops and crop residues (banana)  
 346 fed to livestock and also used as compost. There was  
 347 very little evidence of fertilizer use on the farms apart  
 348 from previous on-farm testing by research and

development organisations. Soil fertility is maintained  
 mainly through natural fallows, improved fallows and  
 leguminous cover crops such as *Mucuna*, *Canavalia*  
 and *Tephrosia* (Wortmann et al., 1998; Wortmann and  
 Kaizzi, 1998).

### 3.3. Nutrient flows and balances

Nutrient flow analysis (NFA) was used to compare  
 situations and outcomes in relative terms (Budelman  
 and Defoer, 2000a,b). Nutrient balances of the three  
 production subsystems (crop, animal and household  
 production systems) and out of the farm system for N,  
 P and K are presented in Figs. 3 and 4. For the crop  
 production system, N, P and K balances were negative  
 in all the five seasons for both class I and III. The

363 nutrient balances for the LR seasons are about twice as  
364 much as for the SR seasons. In the LR more crops are  
365 cultivated and therefore there are more harvests. Thus,  
366 in the LR season, farmers have more surplus of produce  
367 for sale and therefore, export more nutrients out of the  
368 farm. The animal production system had zero or  
369 positive nutrient balance in both LR and SR seasons for  
370 both farm classes. The household system had positive N  
371 balance but with either zero or negative P and K  
372 balances for both classes in all the five seasons. This  
373 further emphasises the point that nutrient stocks of  
374 individual plots within farms and village territories can  
375 differ considerably due to management. Thus plots  
376 around the homestead, which receive substantial  
377 amounts of nutrients from animal manure and house-  
378 hold waste, maintain a relatively high level of  
379 productivity (Smaling and Braun, 1996; Hilhorst  
380 et al., 2000). However, the farm system had net  
381 negative balance for all nutrients in all the seasons and  
382 for all the farm classes. The average N, P and K balances  
383 due to crop removal from the partial crop production  
384 system for class I and III farmers in the LR  
385 and SR seasons were  $-7.3 \text{ kg N ha}^{-1}$ ,  $-1.3 \text{ kg P ha}^{-1}$ ,  
386  $-10.0 \text{ kg K ha}^{-1}$  and  $-4.0 \text{ kg N ha}^{-1}$ ,  $-1.5 \text{ kg}$   
387  $\text{P ha}^{-1}$ ,  $-8.5 \text{ kg K ha}^{-1}$  and  $-0.6 \text{ kg N ha}^{-1}$ ,  $-0.6$   
388  $\text{kg P ha}^{-1}$ ,  $-4.3 \text{ kg K ha}^{-1}$  and  $-3.0 \text{ kg N ha}^{-1}$ ,  $-0.5$   
389  $\text{kg P ha}^{-1}$ ,  $-6.0 \text{ kg K ha}^{-1}$ , respectively (Fig. 2a–d).  
390 The balance was negative due to crop removal of maize,  
391 beans sweet potatoes, cassava, and bananas. Nitrogen, P  
392 and K balances for the animal production system were  
393 marginal or zero as no nutrient entered or left the  
394 system. The household system had positive N, P and K  
395 balance because of the food crops that entered the  
396 system from the crop production system. Looking at the  
397 whole farm system, the export of nutrients from the  
398 farm as sales was greater than the imports. The average  
399 farm-level nutrient exports for class I and III farmers  
400 were more in the LR season than in the SR season  
401 (Fig. 2a–d). Of all the nutrients, substantial amounts of  
402 K were exported through banana fruit and residues',  
403 thus making the K balances more negative. Potassium  
404 export through banana either consumed or sold also  
405 poses a problem as much of it remains in the bodies of  
406 the farm inhabitants, while the rest is excreted but not  
407 returned to the fields.

408 The limited nutrients that enter the farm system are  
409 mainly added to the crop production system, with  
410 lower amounts entering into the household system as

411 food or animal feed. Despite these additions, 411  
412 significant losses occur from the CPS. The crop 412  
413 production system, which is the major source of the 413  
414 nutrients leaving the farm, has the highest risk for soil 414  
415 nutrient depletion. Woelcke and Berger (2002) 415  
416 conducted bio-economic modelling studies in the 416  
417 area using different scenarios and they also found 417  
418 similar nutrient balances. For example, the N balances 418  
419 varied from  $-28 \text{ kg N ha}^{-1}$  (subsistence farm house- 419  
420 holds) to  $-77 \text{ kg N ha}^{-1}$  (commercial farm house- 420  
421 holds) in the case of the baseline scenario under 421  
422 current land management and socio-economic condi- 422  
423 tions (Woelcke and Berger, 2002). The commercial 423  
424 households had higher yields and therefore higher 424  
425 amounts of nutrients were exported in the harvested 425  
426 produce. The results of the NFA suggest that there is 426  
427 need for a more targeted approach to soil fertility 427  
428 interventions that differentiate between farm compo- 428  
429 nents and socio-economic conditions (Elias et al., 429  
430 1998). 430

431 Though farmers classified themselves into three 431  
432 soil fertility management classes of good, average and 432  
433 poor soil fertility managers, however, soil chemical 433  
434 and physical properties revealed no significant fertility 434  
435 differences. Furthermore, the nutrient flows and 435  
436 balances also revealed that there is no significant 436  
437 difference in nutrient management for the three soil 437  
438 fertility classifications. It is instead observed that class 438  
439 1 farmers', despite being good managers, lose/export 439  
440 more nutrients from the farm than class 3 farmers. The 440  
441 soil chemical analysis and the nutrient balance studies 441  
442 results do not reflect the farmers' soil fertility 442  
443 assessment and therefore require further analysis. 443  
444 However, De Jager et al. (1998) followed a budget 444  
445 approach in linking household objectives and wealth 445  
446 to nutrient management and mining and found a strong 446  
447 correlation between market orientation of farm 447  
448 households and the nutrient balance. Thus inspite of 448  
449 higher input use in market oriented farms, outputs 449  
450 were so high that the balance was more negative than 450  
451 in subsistence farming. 451

#### 4. Conclusions 452

453 Resource flows and nutrient balances from this 453  
454 study show that soil nutrient depletion is a major 454  
455 problem in the study area. Nutrient mining is more 455



456 intense in the crop production system of the  
457 smallholder farmers in eastern Uganda. Harvesting  
458 of crops for food and the surplus for sale are the most  
459 important sources of nutrient mining in the crop  
460 production system. Therefore attempts to correct the  
461 imbalance need to address these and other socio-  
462 economic factors. Given the high costs of fertilisers,  
463 intensifying use of legume cover crops as intercrops or  
464 improved fallows and strategic management of crop  
465 residues such as through home gardens are some of the  
466 options for minimising nutrient depletion.

467 The PLAR process enabled farmers to diagnose,  
468 plan, implement and evaluate their own activities for  
469 soil fertility improvement on their farms. The resource  
470 flow mapping exercise was an important tool in guiding  
471 farmers in selecting technologies and solutions accord-  
472 ing to the available farm resources as well as stimula-  
473 ting them to take action. This study also demonstrated  
474 that the maps drawn by the farmers were a source of  
475 information in determining resource flows and calcula-  
476 ting nutrient balances that were used as indicators for  
477 improvements in soil fertility management.

## 478 Acknowledgements

479 The authors thank collaborating farmers who  
480 actively participated in this research, the National  
481 Agricultural Research Organization (NARO) in  
482 Uganda, Africa 2000 Network/UNDP Project and  
483 the Department of Agriculture, Animal Industry and  
484 Fisheries in Iganga District for allowing their field  
485 staff to participate fully in all the activities of the  
486 project. CIAT and NARO also gratefully acknowledge  
487 the financial assistance of the German Federal  
488 Ministry of Technical Co-operation (BMZ).

## 489 References

490 Bekunda, M.A., Bationo, A., Ssali, H., 1997. Soil fertility manage-  
491 ment in Africa: A review of selected research trials. In: Buresh,  
492 R.J., Sanchez, P.A., Calhoun, F. (Eds.), Replenishing Soil Ferti-  
493 lity in Africa. Soil Science Society of America Special Pub-  
494 lication No. 51. Soil Science Society of America, American  
495 Society of Agronomy, Madison, Wisconsin, USA, pp. 63–79.  
496 Budelman, A., Defoer, T., 2000a. Not by nutrient alone: a call to  
497 broaden the soil fertility initiative. *Nat. Resour. Forum* 24, 173–  
498 184.

Budelman, A., Defoer, T., 2000b. PLAR and resource flow analysis 499  
in practice. Case studies from Benin, Ethiopia, Kenya, Mali and 500  
Tanzania (Part 2). In: Defoer, T., Budelman, A., (Eds.), Mana- 501  
ging Soil Fertility in the Tropics: A Resource Guide for Parti- 502  
cipatory Learning and Action Research. Royal Tropical 503  
Institute, Amsterdam, The Netherlands/International Institute 504  
for Environment and Development, London, UK, p. 192. 505  
Defoer, T., Budelman, A., 2000. In: Defoer, T., Budelman, A., 506  
(Eds.), *Managing Soil Fertility in the Tropics. A Resource Guide* 507  
for Participatory Learning and Action Research, Royal Tropical 508  
Institute, Amsterdam, The Netherlands/International Institute 509  
for Environment and Development, London, UK. 510  
Defoer, T., Budelman, A., Toulmin, C., Carter, S.E., 2000. Building 511  
common knowledge. Participatory learning and action research 512  
Part (1). In: Defoer, T., Budelman, A., (Eds.). *Managing Soil* 513  
*Fertility in the Tropics: A Resource Guide for Participatory* 514  
*Learning and Action Research*, Royal Tropical Institute, 515  
Amsterdam, The Netherlands/International Institute for Envir- 516  
onment and Development, London, UK, p. 208. 517  
Defoer, T., De Groot, H., Hilhorst, T., Kante, S., Budelman, A., 518  
1998. Participatory action research and quantitative analysis for 519  
nutrient management in southern Mali: a fruitful marriage? 520  
*Agric., Ecosys. Environ.* 71, 215–228. 521  
De Jager, A., Kariuki, I., Matiri, F.M., Odendo, M., Wanyama, J.M., 522  
1998. Monitoring nutrient flows and economic performance in 523  
African farming systems (NUTMON). IV. Linking farm eco- 524  
nomic performance and nutrient balances in three districts in 525  
Kenya. *Agriculture. Ecosys. Environ.* 71, 81–92. 526  
Delve, R.J., Jama, B., in press. Developing organic resource man- 527  
agement options with farmers in eastern Uganda. 528  
Deugd, M., Rolling, N., Smaling, E.M.A., 1998. A praxeology for 529  
integrated nutrient management, facilitating innovation with and 530  
by farmers. *Agric., Ecosys. Environ.* 71, 269–283. 531  
Elias, E., Morse, S., Belshaw, D.G.R., 1998. Nitrogen and phos- 532  
phorus balances of Kindo Koisha farms in southern Ethiopia. 533  
*Agric., Ecosys. Environ.* 71, 93–113. 534  
Esilaba, A.O., Byalebeka, J.B., Nakiganda, A., Mubiru, S., 535  
Ssenyange, D., Delve, R., Mbalule, M., Nalukenge, G., 536  
2001a. Farmer evaluation of integrated nutrient management 537  
strategies in eastern Uganda. A paper presented at The Southern 538  
and Eastern Africa Association of Farming Systems Research 539  
and Extension (SEAAFSRE-E), Eighth Conference held in 540  
Nairobi, Kenya, 20–24 August, 2001. SEAAFSRE-E, Egerton 541  
University and Kenya Agricultural Research Institute, Nairobi, 542  
Kenya. 543  
Esilaba, A.O., Byalebeka, J.B., Nakiganda A.A., Mubiru, S., Sse- 544  
nyange, D., Delve, R., Mbalule, M., Nalukenge, G., 2001b. 545  
*Integrated Nutrient Management in Iganga District, Uganda:* 546  
*Diagnosis by Participatory Learning and Action Research, Net-* 547  
*work on Bean Research in Africa, Occasional Publications* 548  
*Series, No. 35, CIAT, Kampala, Uganda.* 549  
FAO, 1977. *Soil Map of the World, 1:5,000,000. Food and Agri-* 550  
*culture Organization of the United Nations and the United* 551  
*Nations Educational, Vol. VI, Scientific and Cultural Organiza-* 552  
*tion.* 553  
Fischler, M., 1997. Legume green manures in the management of 554  
maize-bean cropping systems in eastern Africa with special 555

- reference to crotalaria (*C. Ochroleuca* G. Don.) PhD. Thesis, Swiss Federal Institute of Technology, Zurich, Switzerland.
- Foster, H.L., 1971. Rapid soil and plant analysis without automatic equipment. I. Routine soil analysis. *East Afric. Agric. Forestry J.* 37, 160–171.
- Foster, H.L., 1973. Fertilizer recommendations for cereals grown on soils derived from volcanic rocks in Uganda. *East Afric. Agric. Forestry J.* 38, 303–313.
- Grandin, B., 1988. *Wealth Ranking in Smallholder Communities: A Field Manual*. Publications Intermediate Technology Publications, Nottingham, England.
- Harrop, J.F., 1970. Soils. In: Jameson, J.D. (Ed.), *Agriculture in Uganda*. Oxford University Press, UK, pp. 43–71.
- Hilhorst, T., Muchena, F., Defoer, T., Hassink, J., De Jager, A., Smaling, E., Toulmin, C., 2000. Managing soil fertility in Africa: diverse settings and changing practice. In: Hilhorst, T., Muchena, F. (Eds.), *Nutrients on the Move—Soil Fertility Dynamics in African Farming Systems*. International Institute for Environment and Development, London, pp. 1–25.
- Jackson, M.L., 1958. *Soil Chemical Analysis*. Constable and Co. Ltd., London.
- Landon, J.R., 1984. *Booker Tropical Soil Manual. A Handbook for Soil Survey and Agricultural Land Evaluation in the Tropics and Subtropics*. Booker Agricultural International Limited.
- Okalebo, J.R., Gathua, K.W., Woome, P.L., 1993. *Laboratory Methods of Soil and Plant Analysis. A Working Manual*. Soil Science Society of East Africa, UNESCO/ROSTA, Nairobi, Kenya.
- Sanchez, P.A., Shepperd, K.D., Soule, M.J., Place, F.M., Buresh, R.J., Izac Anne-Marie, N., Mokwunye, A.U., Kwesiga, F.R., Ndiritu, C.G., Woome, P.L., 1997. Soil fertility replenishment in Africa: An investment in natural resource capital. In: Buresh, R.J., Sanchez, P.A., Calhoun, F. (Eds.), *Replenishing Soil Fertility in Africa*. Soil Science Society of America Special Publication No. 51. Soil Science Society of America, American Society of Agronomy, Madison, Wisconsin, USA, pp. 1–46.
- SAS, 1990. SAS Institute Inc., SAS/STAT. Users Guide, Version 6, fourth ed., Vol. 2, Cary, NC, SAS Institute Inc., USA, p. 846.
- Scoones, I., 2001. *Dynamics and Diversity: Soil Fertility and Farming Livelihoods in Africa: Case Studies from Ethiopia, Mali, and Zimbabwe*. Earthscan Publications Ltd., London, 244.
- Scoones, I., Toulmin, C., 1999. *Policies for Soil Fertility Management in Africa*. Department of International Development (DFID), London, p. 128.
- Smaling, E.M.A., Braun, A.R., 1996. Soil fertility research in sub-Saharan Africa: New dimensions, new challenges. *Commun. Soil Sci. Plant Anal.* 27, 365–386.
- Smaling, E.M.A., Nandwa, S.M., Janssen, B.H., 1997. Soil fertility in Africa is at stake. In: Buresh, R.J., Sanchez, P.A., Calhoun, F. (Eds.), *Replenishing Soil Fertility in Africa*. Soil Science Society of America, American Society of Agronomy, Soil Science Society of America Special Publication No. 51, Madison, Wisconsin, USA, pp. 47–61.
- Smaling, E.M.A., Stoorvogel, J.J., Windmeijer, P.N., 1993. Calculating soil nutrient balances in Africa at different scales II. District scale. *Fert. Res.* 35, 237–250.
- SPSS, 2002. *Statistical Packages for Social Scientists, SPSS Users Guide*, SPSS Inc., Chicago, IL, USA.
- Stoorvogel, J.J., Smaling, E.M.A., 1990. Assessment of soil nutrient depletion in sub-Saharan Africa: 1983–2000, Report 28, The Winand Staring Centre, Wageningen, The Netherlands.
- Swift, M.J., Seward, P.D., Frost, P., Qureshi, J., Muchena, F., 1994. Long-term experiments in Africa: Developing a database for sustainable land use under global change. In: Leigh, R.A., Johnston, A.E. (Eds.), *Long-Term Experiments in Agricultural and Ecological Sciences*. CAB International, Wallingford, England, pp. 229–251.
- Van den Bosch, H., De Jager, A., Vlaming, J., 1998. Monitoring nutrient flows and economic performance in African farming systems (NUTMON) II. Tool development. *Agric., Ecosys. Environ.* 71, 54–64.
- Van der Pol, F., 1992. Soil Mining: An Unseen Contributor to Farm Income in southern Mali. *Bulletin* 325. Royal Tropical Institute, Amsterdam, p. 47.
- Woelcke, J., Berger, T., 2002. Land management and technology adoption in eastern Uganda: An integrated bio-economic modelling approach. A Paper Presented at The Final Workshop on Policies for Improved Land Management in Uganda, April 17–19, 2002. International Food Policy Research Institute (IFPRI), Washington, DC, USA.
- Wortmann, C.S., Fischler, M., Alifugani, F., Kaizzi, C.K., 1998. Accomplishments of participatory research for systems improvement in Iganga District, Uganda 1993–1997, Occasional Publication Series, No. 27, CIAT, Kampala, Uganda, p. 40.
- Wortmann, C.S., Kaizzi, C.K., 1998. Nutrient balances and expected effects of alternative practices in the farming systems of Uganda. *Agric., Ecosys. Environ.* 71, 117–131.