

Elsevier Editorial System(tm) for Agriculture, Ecosystems and Environment
Manuscript Draft

Manuscript Number: AGEE5072R3

Title: Agricultural management affects earthworm and termite diversity across humid to semi-arid tropical zones

Article Type: Research Paper

Keywords: Soil biodiversity; earthworms; termites; agriculture; crop management; soil carbon; climate

Corresponding Author: Mr Fredrick Ouma Ayuke, M.Phil

Corresponding Author's Institution: Wageningen University

First Author: Fredrick O Ayuke, PhD

Order of Authors: Fredrick O Ayuke, PhD; Mirjam M Pulleman, PhD; Bernard Vanlauwe, PhD; Ron G de Goede, PhD; Csaba Csuzdi, PhD; Lijbert Brussaard, PhD

Manuscript Region of Origin: NETHERLANDS

Abstract: Earthworm and termite diversity were studied in 12 long-term agricultural field trials across the sub-humid to semi-arid tropical zones of Eastern and Western Africa. In each trial, treatments with high and low soil organic C were chosen to represent contrasts in long-term soil management effects, including tillage intensity, organic matter and nutrient management and crop rotations. For each trial, a fallow representing a relatively undisturbed reference was also sampled. Earthworm taxonomic richness decreased in the direction fallow > high-C soil > low-C soil and earthworm abundance was higher in fallow than under continuous crop production. Termite abundance was not significantly different between fallow and high and low-C treatments and termite taxonomic richness was higher in fallow soil than in the two cropping systems. We concluded that fewer species of earthworms and termites were favored under agricultural management that led to lower soil C. Results indicated that the soil disturbance induced by continuous crop production was more detrimental to earthworms than to termites, when compared to the fallow.

Wageningen, November 24th, 2010

Dear Dr. Olaf Meesters,

Please find attached our revised manuscript AGEE5072R2, entitled "Agricultural management affects earthworm and termite diversity across humid to semi-arid tropical zones". We thank the editor for the valuable comments, which has still helped us to substantially improve the quality of our manuscript. As indicated in our revision notes (below), we have incorporated most of the suggestions made. When we had divergent view with the comments made by the editor, we clearly explained ourselves. We hope that, after this round of revisions, our manuscript meets the standards for publication in Agriculture, Ecosystems and Environment.

On behalf of all authors,

Ayuke Fredrick Ouma

Dept. of Soil Quality

Wageningen University

The Netherlands

Editor's comments and our response (in red):

EDITOR

This new version still requires some more attention with a view to:

- simplify Highlights to :

? [Earthworms](#) and termites were favored by warm and dry conditions + Done.

? Cropping negatively affected earthworms more than termites + Done.

? Management leading to low soil C supported fewer earthworm and termite taxa + Done.

- stick to common units e.g. "t" instead of Mg in Table 1 + Done

; "min" l. 60, p. 4 + Done (see l. 4, p. 5).

- modify l. 24, p. 5 to: "Earthworms and termites were..." + Done (l. 26, p. 5); l. 31, p. 5 to "Adult earthworms and termites were... + Done (l. 34, p. 5) " omitting l. 38-51 ; + Done.
- erase whole () l. 56, p. 5; + Done.
- place also Table 2 under Supplementary Material; With reference to table 2. + The authors still respectfully disagree with this. This table captures hypotheses 2 and 3 and is the core of the study as reflected in the title. Moreover, by removing the table, the results in sections 3.1, 3.2 and discussion (section 4.2) would become irrelevant. Therefore, the authors feel the table should remain in the manuscript.
- change l. 37, p. 7 to : "However, site-treatment interactions were also significant and there were significant...";+ Done (l. 22-24, p. 7).
- stick to past tense as of l. 48, 50, p. 8 ; etc. + Effected (l. 32-42, p. 8; l. 59, p. 8 to l. 17, p. 9).
- check caption to both Figures correcting "sp." throughout and "vitriolatus", referring to Figure 1 for environmental abbreviations. + Effected.

Agricultural management affects earthworm and termite diversity across humid to semi-arid tropical zones

- Earthworms and termites were favored by warm and dry conditions.
- Cropping negatively affected earthworms more than termites.
- Management leading to low soil C supported fewer earthworm and termite taxa.

1
2
3
4
5 **Agricultural management affects earthworm and termite diversity across humid to semi-**
6
7 **arid tropical zones**
8
9

10 F.O. Ayuke ^{a*, b}, M.M. Pulleman ^a, B. Vanlauwe ^b, R.G.M. de Goede ^a, J. Six ^c., C. Csuzdi ^d, L.
11
12
13 Brussaard ^a
14
15

16
17
18 ^a*Department of Soil Quality, Wageningen University, P.O Box 47, NL-6700 AA Wageningen,*
19
20 *The Netherlands*

21
22 ^b*Tropical Soil Biology and Fertility (TSBF) Institute of CIAT, P.O Box 30677-00100, Nairobi,*
23
24 *Kenya*

25
26
27 ^c*Department of Plant Sciences, University of California, Davis, CA 95616, USA*

28
29 ^d*Systematic Zoology Research Group of HAS and Hungarian Natural History Museum, Baross*
30
31 *str. 13, H-1088 Budapest, Hungary*
32
33

34
35
36 *Corresponding author: (0317) 486524; Fax: (0317) 419000
37
38

39 Email: fredrick.Ayuke@wur.nl or fredrick.ayuke@yahoo.com*
40
41
42
43

44 **ABSTRACT**
45

46
47 Earthworm and termite diversity were studied in 12 long-term agricultural field trials across the
48
49 sub-humid to semi-arid tropical zones of Eastern and Western Africa. In each trial, treatments
50
51 with high and low soil organic C were chosen to represent contrasts in long-term soil
52
53 management effects, including tillage intensity, organic matter and nutrient management and
54
55 crop rotations. For each trial, a fallow representing a relatively undisturbed reference was also
56
57
58 sampled. Earthworm taxonomic richness decreased in the direction fallow > high-C soil > low-C
59
60
61
62
63
64
65

1
2
3
4 soil and earthworm abundance was higher in fallow than under continuous crop production.
5
6 Termite abundance was not significantly different between fallow and high and low-C treatments
7
8 and termite taxonomic richness was higher in fallow soil than in the two cropping systems. We
9
10 concluded that fewer species of earthworms and termites were favored under agricultural
11
12 management that led to lower soil C. Results indicated that the soil disturbance induced by
13
14 continuous crop production was more detrimental to earthworms than to termites, when
15
16 compared to the fallow.
17
18
19
20
21
22
23

24
25 *Key words:* Soil biodiversity, earthworms, termites, agriculture, crop management, soil carbon,
26
27 climate
28
29
30
31
32
33

34 **1. Introduction**

35
36
37 Knowledge of the spatial distribution of different taxa of earthworms and termites in
38
39 agricultural soils across climatic regions and soil types is necessary in determining how, and to
40
41 what extent, the diversity and activities of these important soil ecosystem engineers are affected
42
43 by climate versus agricultural land use and management. Site-specific studies by Kooyman and
44
45 Onck (1987), Eggleton et al. (1996), Okwakol (2000), Birang et al. (2003) and Curry et al.
46
47 (2002) have indicated that deforestation, soil disturbance and increased intensity of agriculture
48
49 are among the management practices that negatively impact soil macrofauna species richness and
50
51 abundance in general. Clearance and tillage of habitats result in destruction of nests, nesting
52
53 sites and burrow systems for termites and earthworms respectively, thus exposing them to harsh
54
55 environmental conditions and/or predators (House and Parmelee, 1985; Barros et al., 2002). Soil
56
57
58
59
60
61
62
63
64
65

1
2
3
4 management practices such as use of organic inputs and crop diversification through rotation
5
6 favor macrofauna diversity due to improvements in the abiotic conditions and increased substrate
7
8 supply (Tian et al., 1997; Ayuke et al., 2003; Curry, 2004; Osler et al., 2008, Fonte et al., 2009).
9
10 Although research has continued to demonstrate that tillage and other agronomic practices (e.g.
11
12 mulching, organic and inorganic fertilizer applications, cover crop management, crop rotation
13
14 and rotation pattern) influence changes in invertebrate communities (Smith et al., 1988; Zehnder
15
16 and Linduska, 1987), such studies have mostly targeted aboveground fauna, and the effects on
17
18 belowground invertebrate communities in cropped soils are little understood, especially in the
19
20 case of termites. This study, therefore, aimed at assessing how agricultural management affects
21
22 earthworm and termite diversity across humid to semi-arid tropical regions. We hypothesized
23
24 that the biodiversity of earthworms and termites will: 1) decrease with increasing temperature
25
26 and decreasing precipitation, 2) be lower in agricultural than in fallow systems, and 3) be higher
27
28 under long-term agricultural management that has led to high-C soils than where it has led to
29
30 low-C soils.
31
32
33
34
35
36
37
38
39
40
41

42 **2. Material and methods**

43
44 The study was conducted in 12 long-term field trials across the sub-humid to semi-arid
45
46 tropical zones of Eastern Africa (Embu, Kabete, Impala and Nyabeda in Kenya and Chitala in
47
48 Malawi) and Western Africa (Tamale in Ghana, Ibadan in Nigeria, Sadore in Niger and
49
50 Farakoba, Saria I, II and III in Burkina Faso). The long-term trials were established in the past
51
52 one to four decades and aimed at testing different management options for arable crop
53
54 production such as organic versus mineral inputs, crop rotation, and tillage. In our sampling
55
56 scheme, the arable treatments that, according to previously available data, had resulted in the
57
58
59
60
61
62
63
64
65

1
2
3
4 highest and lowest soil organic carbon (SOC) contents were included (Table 1). The SOC
5
6 content is known to increase due to improved long-term management practices, such as reduced
7
8 tillage (Six et al., 1999; Pikul et al., 2009), organic matter and nutrient inputs (Ouédraogo et al.,
9
10 2001; Palm et al., 2001; Vanlauwe et al., 2002) and crop rotations (Wu et al., 2003). These
11
12 management practices have been found to also have a beneficial effect on soil macrofauna
13
14 activity (Tian et al., 1997; Mando and Brussaard, 1999; Ayuke et al., 2003; Fonte et al., 2009). A
15
16 fallow located nearby the arable fields representing a reference with continuous plant cover and
17
18 without soil disturbance was also included. Depending on site, the fallows consisted of green
19
20 fallow, forest or shrubland, which may or may not have been used for crop production in the
21
22 past, but had been undisturbed for at least 10 years at the time of sampling. No insecticides were
23
24 applied in the fields we sampled for this study.
25
26
27
28
29
30

31 **(Table 1)**

32 33 34 35 36 37 *2.1. Macrofauna sampling*

38
39
40 Monolith sampling and transect sampling were used for termite assessments while
41
42 earthworms were collected through monolith sampling only (Swift and Bignell, 2001; Bignell et
43
44 al., 2008). Sampling was done six to eight weeks after planting (July-September, 2006 in the
45
46 Western Africa sites and February-June 2007 in Eastern Africa sites). One soil monolith of 25 ×
47
48 25 × 30 cm depth was randomly taken in each replicate plot (n = 3). The soil was separated into
49
50 two depth layers (0-15 and 15-30 cm) and earthworms and termites were collected by hand-
51
52 sorting on plastic trays. Transect sampling for termites was done alongside the monolith
53
54 sampling. In each sampling plot, a 20 × 2 m transect (or 5 × 2 m sections depending on the
55
56 dimension of the plots) was randomly laid out. Each transect or section was sampled sequentially
57
58
59
60
61
62
63
64
65

1
2
3
4 by two trained persons for 30 min, and the following microhabitats which are common sites for
5
6 termites were searched manually: surface soil to 5 cm depth; the soil between large buttress roots
7
8 if any; the inside of branches and twigs; the soil within and beneath rotten logs and smaller
9
10 pieces of dead wood; subterranean nests, mounds, carton sheeting and runways on vegetation
11
12
13
14 (Jones and Eggleton, 2000; Swift and Bignell, 2001).
15
16
17
18

19 *2.2. Macrofauna preservation and identification*

20
21 Termites were preserved in 75% alcohol and earthworms were killed in 75% alcohol
22
23 before fixation in 4% formaldehyde. Termites and earthworms were stored in sealed vials before
24
25 being transported to the laboratory for identification and enumeration. Earthworms and termites
26
27 were identified at the Department of Invertebrate Zoology of the National Museums of Kenya
28
29 (NMK), Nairobi, Kenya, using appropriate keys and reference specimens in the NMK
30
31 collections. Adult earthworms and termites were identified to species or where this proved
32
33 difficult, to numbered morphospecies. Unidentified earthworm specimens were sent to the
34
35 Hungarian Natural History Museum (HNHM), Budapest, Hungary, for further analysis. Voucher
36
37 specimens are being held at the NMK, Nairobi and HNHM, Budapest.
38
39
40
41
42

43 Taxonomic richness (S) was calculated as the number of taxa per monolith and transect
44
45 sample (pooled together from each experimental plot for termites).
46
47
48
49
50

51 *2.3. Statistical analyses*

52
53 The data obtained were subjected to analysis of variance (ANOVA) using GENSTAT
54
55 11.1 (2009). Levene's test was used to assess homogeneity in the data (Field, 2005). Due to non-
56
57 homogeneity of variances in the termite and earthworm data (diversity indices and abundance),
58
59
60
61
62
63
64
65

1
2
3
4 data were square root transformed $(x + 0.5)^{1/2}$. Levels of significance were evaluated using
5
6
7 Fisher's least significant difference (LSD).
8

9 The strength and statistical significance of species-environment relationships were
10 analyzed with the programme CANOCO 3.1 (Ter Braak, 1988). With CANOCO, a preliminary
11 detrended correspondence analysis (DCA) was conducted to determine the length of the gradient,
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The strength and statistical significance of species-environment relationships were analyzed with the programme CANOCO 3.1 (Ter Braak, 1988). With CANOCO, a preliminary detrended correspondence analysis (DCA) was conducted to determine the length of the gradient, test the homogeneity of the data for abundance of earthworms and termites, identified to the lowest taxonomic level possible, and decide on which ordination analysis to apply. In this study, length of gradient determined by DCA was <4 for the first axis, so we opted for the linear ordination technique Redundancy Analysis (RDA). RDA was used to investigate the correlative relationships of environmental variables (latitude, longitude, altitude, average temperature, precipitation, clay, sand and soil carbon) with both earthworm and termite abundance at lowest taxonomic level possible. Forward selection was applied to assess the significance and strength of each environmental variable within the RDA. The first variable selected was that with the highest marginal eigenvalue (i.e. its explanatory fit to the species and site data if added as the only environmental variable in the analysis). Subsequently, the environmental variables were entered one at a time in order of the magnitude of their conditional eigenvalues (i.e. additional fit after adding previous variables), until none of the remaining variables significantly explained additional variation in abundance of earthworms and termites. In the subsequent analysis only those environmental variables with conditional statistical significance in the above RDA were included. Overall contribution of each of the variables to description of the variation in macrofauna data was tested by a Monte-Carlo test based on 999 random permutations. Since the management treatments were nested within the trial sites, we restricted permutations to a split-

1
2
3
4 plot design, such that trial sites were treated as whole plot factors, while management treatments
5
6 within trial sites were treated as split plot factors (Ter Braak and Smilauer, 1998).
7
8
9

10 11 **3. Results**

12 13 *3.1. Earthworm and termite taxonomic richness and abundance*

14
15 Mean earthworm taxonomic richness across all sites was significantly higher in the
16
17 fallow (1.72) than in the high-C treatments (1.02), where, in turn, the richness was significantly
18
19 higher than in the low-C (0.64) treatments. However, site-treatment interactions were also
20
21 significant and there were significant differences in taxonomic richness between sites (Table 2).
22
23 Differences in mean earthworm abundance showed the same pattern as differences in mean
24
25 taxonomic richness, but the difference between high-C and low-C treatments was not significant
26
27 at any of the sites (Table 2).
28
29
30
31
32
33
34

35 Mean termite taxonomic richness across all the sites showed a pattern similar to that of
36
37 earthworms, also following the order fallow > high-C > low-C treatments. Whereas the
38
39 difference between fallow and low-C treatments was significant, this was not the case for the
40
41 difference between fallow and high-C treatments, and between high-C and low-C (Table 2).
42
43 Also, not at all sites the taxonomic richness followed the order high-C > low-C treatment (Table
44
45 2). Hence, site-treatment interactions were not significant. Termite abundance was highly
46
47 variable across replicates and no significant differences were found between treatments (fallow,
48
49 high-C and low- C) (Table 2). Termite abundance, like taxonomic richness, differed significantly
50
51 between sites.
52
53
54
55
56

57 **(Table 2)**
58
59
60
61
62
63
64
65

3.2. Correlations between abundances and environmental parameters

Of the eight variables (latitude, longitude, altitude, average temperature, precipitation, soil clay content, soil sand content, and soil carbon) initially entered into the RDA, six (precipitation, average temperature, soil carbon, altitude, longitude and latitude) were significantly correlated with earthworm abundance ($p < 0.001$). The eigenvalues of the first and second RDA axes constrained to the environmental variables were 0.261 and 0.069, respectively, and the two axes explained 33% of the variation in earthworm taxa distribution (Figure 1A). The sum of all canonical eigenvalues showed that the environmental variables that significantly contributed to the description of the variation in the earthworm fauna (latitude, longitude, altitude, temperature, precipitation and soil carbon content) explained 42% of the total variation observed (data not shown). The first axis was mainly a temperature/altitudinal gradient ($r = -0.83$ and 0.86 respectively), while the second axis was a precipitation/latitudinal gradient ($r = 0.12$ and -0.16 respectively) (Figure 1A). Axis 1 separated cooler, higher-altitude sites from hotter, lower-altitude sites. Earthworm taxa were less abundant on the higher, cooler altitudes that are characteristic of the sites sampled in Eastern Africa (Figure 1A and 1B).

For termites, eigenvalues of the first and second RDA axes constrained to the environmental variables were 0.135 and 0.067, respectively, and the two axes explained 20.2% of variation in termite taxa distribution (Figure 2A). The sum of all canonical eigenvalues showed that the environmental variables that significantly contributed to the description of the variation in the termite fauna (precipitation, longitude, average temperature, soil sand content and latitude), explained 25% of the total variation observed (data not shown). The first axis was a

1
2
3
4 precipitation/latitudinal gradient ($r = 0.81$ and -0.63 respectively), while the second axis was a
5
6 temperature/texture/longitudinal gradient ($r = -0.84$, -0.77 and 0.85 respectively). A higher than
7
8 average number of taxa occurred in the sites with warmer and drier climatic conditions (Figure
9
10 2A). Axis 2 mainly separated Eastern and Western Africa with mean rainfall being higher, and
11
12 temperature lower in Eastern Africa and termite taxa were less abundant in most of the sites here
13
14 (Figure 2B).
15
16
17

18 **(Figure 2)**
19
20
21
22

23 **4. Discussion**

24
25
26 At most sites, soil C in the fallow treatments were relatively higher than in the arable
27
28 (high-C and low-C) treatments. Differences between high-C and low-C in some cases, especially
29
30 in Saria 1 and 3, however, were relatively small (Table 1). Variability in soil C across sites was
31
32 large. Sites in Eastern Africa had higher soil C than sites in Western Africa (Table 1). In view of
33
34 the large variability in the data, results can be considered robust, wherever differences in
35
36 macrofauna diversity between treatments were significant.
37
38
39
40
41
42
43

44 *4.1. Influence of environmental factors*

45
46
47 The number of taxa of both earthworms and termites was low in most of the sites. Mean
48
49 taxonomic richness ranged between 0-5 for earthworms and 0-3 for termites. The number of
50
51 termite taxa collected in our sites was lower than the range of 10-24 recorded by Kooyman and
52
53 Onck (1987) and Eggleton et al. (1996, 2002) across fallow and arable fields of Western Kenya
54
55 and around the Mbalmayo forest reserve in Cameroun, respectively. However, it conforms with
56
57 the range of 3-5 taxa recorded by Okwakol (2000) across arable sites in Uganda.
58
59
60
61
62
63
64
65

1
2
3
4 Highly significant correlations were observed between macrofauna richness, and
5
6 environmental variables, reflecting the influence of climate and soil type. The two RDA biplots
7
8 for earthworms (Figure 1) and termites (Figure 2) separated the relatively cooler, wetter, more
9
10 clayey Eastern Africa sites from the warmer, drier, more sandy Western Africa sites. Contrary to
11
12 our hypothesis, more than average numbers of earthworm and termite taxa were found under
13
14 relatively warmer, drier conditions. This is not in line with the observation that earthworm and
15
16 termite diversity increases with increase in rainfall or soil moisture, as generally found at least in
17
18 temperate climates (Bohlen et al., 1995; Curry, 2004). However, seasonality of rainfall in the
19
20 tropical regions means rainfall amount per season may be more important than annual total. We
21
22 attributed lower taxonomic richness among the sites in Eastern Africa to less favourable
23
24 conditions arising from high rainfall and low temperatures at higher altitudes. These observations
25
26 were consistent with those of Collins (1980) and Gathorne-Hardy et al. (2001), who noted that
27
28 high altitudes harbour less termite taxa, especially of the subfamily Macrotermitinae. It however,
29
30 contradicted findings by Buxton, (1981), who observed an increase in the number of termite
31
32 species with increasing rainfall. Widespread presence of *Microtermes* spp. across all sites
33
34 illustrated how widely adapted these are to varying climatic conditions and management
35
36 practices. This may partly be attributed to their nesting habits and also their ability to utilize a
37
38 wide variety of food resources which include wood, litter, soil and dung. These termites were
39
40 observed to build both aboveground mounds as well as underground tunnels that can extend
41
42 many metres, probably a strategy to avoid harsh conditions while foraging. This is in contrast to
43
44 group I feeders which nest and feed on the same or nearby pieces of wood, and group III feeders
45
46 that commonly make nests on the ground to feed on surrounding humus or humified soil (Bignell
47
48 and Eggleton, 2000; Gathorne-Hardy et al., 2001). Some termite and earthworm taxa, however,
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 were better represented under cool and wet conditions, in agreement with our hypothesis (Figure
5
6 1A and 2A).
7
8
9

10 11 12 *4.2. Effect of contrasting agricultural management* 13 14

15 In agreement with our hypothesis, higher earthworm taxonomic richness and abundance
16 was recorded in the fallow than in the agricultural systems. Similarly, under continuous crop
17 production, higher earthworm taxonomic richness was observed under agricultural management
18 that had resulted in high-C than low-C soils. This indicates that earthworms are sensitive to
19 disturbances associated with agriculture and agricultural land use intensity, and that their role can
20 be significantly enhanced by applying management practices that increase soil C in arable
21 systems.
22
23
24
25
26
27
28
29
30
31

32
33 Contrary to our hypothesis, no difference in termite taxonomic richness and abundance
34 was observed between fallow and high-C arable systems. We attributed this to the wide feeding
35 and foraging habits of most of the species sampled. Very few typical soil feeders (group IV)
36 were found. Rather, a large proportion of the termites sampled in this study belong to feeding
37 group II and to a smaller extent to feeding group I and III. These groups are largely foragers of
38 wood, litter, soil, dung/manure and grass. They were probably scavenging for food all over the
39 plots and, hence, were sampled uniformly across treatments. In agreement with our hypothesis,
40 higher termite abundance was recorded in agricultural management that had resulted in high-C
41 than in low-C soils. We suggest that the foraging groups utilizing organic residues as food were
42 favoured by the addition of organic resources characteristic of many of our high C treatments.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

This study was supported by WOTRO Science for Global Development of The Netherlands Organization for Scientific Research (NWO). We thank our partners, especially the scientists and field technicians of the Tropical Soil Biology and Fertility Institute of the International Centre for Tropical Agriculture (TSBF-CIAT, Kenya), Kenya Agricultural Research Institute (KARI), Kenya Forestry Research Institute (KEFRI), Kenyatta University, Kenya, International Institute of Tropical Agriculture (IITA, Nigeria), International Center for Soil and Agricultural Development (IFDC, Burkina Faso), Institute of Natural Environmental and Agricultural Research (INERA, Burkina Faso), Savanna Agricultural Research Institute (SARI, Ghana), Chitedze Agricultural Research Station (Malawi), International Crops Research Institute for the Semi-Arid Tropics (ICRISAT, Niger), National Museums of Kenya and Natural History Museums (Hungary), who agreed to work with us and have allowed us to use their experimental sites and facilities for this study.

References

- Ayuke, F.O., Rao, M.R., Swift, M.J., Opondo-Mbai, M.L., 2003. Impact of soil fertility management strategies on diversity and populations of soil macrofauna in agroecosystem. Special Issue: Sustainable use of land resources to alleviate poverty in the new millenium. *East Afr. Agric. For. J.* 69 (1, 2), 131-137.
- Barros, E., Pashanasi, B., Constantino, R., Lavelle, P., 2002. Effects of land-use system on the soil macrofauna in western Brazilian Amazonia. *Biol. Fertil. Soils.* 35, 338-377.
- Bignell, D.E., Eggleton, P., 2000. Termites in ecosystems. In: Abe, T. (Ed.), *Termites: Evol. Soc. Symb. Ecol.* 17, 363-387.

- 1
2
3
4 Bignell, D.E., Constantino, R., Csuzdi, C., Karyanto, A., Konate, S., Louzada, J., Susilo, F.X.,
5
6 Tondoh, J.E., zanetti, R., 2008. Macrofauna. In: Moreira, F.M.S., Huising, J.E., Bignell,
7
8 D. (Eds.). A Handbook of Tropical Soil Biology: Sampling and Characterization of
9
10 Below-ground Biodiversity. Earthscan, UK, 3, pp. 43-83.
11
12
13
14 Birang, M., Csuzdi, C., Hauser, S., Zebaze, I., Didden, W.A. and Brussaard, L., 2003.
15
16 Earthworm Community Structure along a Gradient of Land use Intensification in the
17
18 Humid Forest Zone of Southern Cameroun. Wageningen Univ. Papers. 47, 819-824.
19
20
21 Bohlen, P.J., Edwards, W.M., Edwards, C.A., 1995. Earthworm community structure and
22
23 diversity in experimental agricultural watersheds in Northeastern Ohio. Plant Soil 170,
24
25 233-239.
26
27
28 Buxton, R.D., 1981. Changes in the composition and activities of termite communities in relation
29
30 to changing rainfall. *Oecologia* 51, 371-378.
31
32
33 Collins, N.M., 1980. The distribution of soil macrofauna o the West ridge of Gunung (Mount)
34
35 Mulu, Sarawak. *Oecologia* 44, 263-275.
36
37
38 Curry, J.P., 2004. Factors affecting the abundance of earthworms in soils. In: Edwards, C.A.
39
40 (Ed.), *Earthworm Ecology*, 2nd ed. CRC Press, Boca Raton, London, New York, pp. 91-
41
42 111.
43
44
45 Curry, J.P., Byrne, D., Schmidt, O., 2002. Intensive cultivation can drastically reduce
46
47 earthworm populations in arable land. *Eur. J. Soil Biol.* 38, 127-130.
48
49
50 Eggleton, P., Bignell, D.E., Sands, W.A., Mawdsley, N.A., Lawton, J.H., Wood, T.G. and
51
52 Bignell, N.C., 1996. The diversity, abundance and biomass of termites under differing
53
54 levels of disturbance in the Mbalmayo Forest Reserve, southern Cameroun. *Phil. Trans.R.*
55
56 *Soc. Lond. B.* 351, 51-68.
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Eggleton, P., Bignell, D.E., Hauser, S., Dibog, L., Norgrove, L., Birang, M., 2002. Termite
5
6 diversity across an anthropogenic disturbance gradient in the humid forest zone of West
7
8 Africa. *Agric. Ecosyst. Environ.* 90, 189-202.
9
- 10
11 Field, A., 2005. *Discovering Statistics using SPSS*, 2nd ed. SAGE Publication.
12
13
- 14 Fonte, S.J., Winsome, T., Six, J., 2009. Earthworm populations in relation to soil organic matter
15
16 and management in California tomato cropping systems, *Appl. Soil Ecol.* 41, 206-214.
17
18
- 19 Gathorne-Hardy, F.G., Syaukani., Eggleton, P., 2001. The effects of altitude and rainfall on the
20
21 composition of the termites (Isoptera) of the Leuser Ecosystem (Sumatra, Indonesia). *J.*
22
23 *Trop. Ecol.* 17, 379-393.
24
25
- 26 GENSTAT 11.1., 2009. Genstat release, 11th ed. VSN International LTD.
27
28
- 29 House, G.J., Parmelee, R.W., 1985. Comparison of soil arthropods and earthworms from
30
31 conventional and no-tillage agroecosystems. *Soil Tillage Res.* 5, 351-360.
32
33
- 34 Jones, D.T., Eggleton, P., 2000. Sampling termite assemblages in tropical forests: testing a rapid
35
36 biodiversity assessment protocol. *J. Appl. Ecol.* 37, 191-203.
37
38
- 39 Kooyman, C., Onck, R.F.M., 1987. The interaction between termite activity, agricultural
40
41 practices and soil characteristics in Kisii district, Kenya. *Wageningen Agric. Univ.*
42
43 *Papers*, 87-3.
44
45
- 46 Mando, A., Brussaard, L., 1999. Contribution of termites to the breakdown of straw
47
48 under Sahelian conditions. *Bio. Fertil. Soils* 29, 332-334.
49
50
- 51 Okwakol, M.J.N., 2000 Changes in termite (Isoptera) communities due to the clearance and
52
53 cultivation of the tropical forest in Uganda. *Afr. J. Ecol.* 38, 1-7.
54
55
- 56 Osler, G.H.R., Harrison, L., Kanashiro, D.K., Clapperton, M.J., 2008. Soil
57
58
59
60
61
62
63
64
65

1
2
3
4 microarthropod assemblages under different arable crop rotations in Alberta, Canada.
5
6 Appl. Soil Ecol. 38, 71-78.
7

8
9 Ouédraogo, E., Mando, A., Zombie, N.P., 2001. Use of compost to improve soil properties and
10
11 crop productivity under low-input agricultural system in West Africa. Agric. Ecosyst.
12
13 Environ. 84, 259-266.
14

15
16 Palm, C.A., Gachengo, C.N., Delve, R.J. Cadisch, G., Giller, K., 2001. Organic inputs for soil
17
18 fertility management in tropical agroecosystems: application of an organic resource
19
20 database. Agric. Ecosyst. Environ. 83, 27-42.
21
22

23
24 Pikul Jr, J.L., Chilom, G, Rice, J., Eynard, A., Schumacher, T.E., Nichols, K., Johnson, J.M.F.,
25
26 Wright, S., Caesar, T., Ellsbury, M. 2009. Organic matter and water stability of field
27
28 aggregate affected by tillage in South Dakota, Soil Sci. Soc. Am. J. 73 (1), 197-206.
29

30
31 Six, J., Elliott, E.T., Paustian, K., 1999. Agregate & Soil Organic Matter Dynamics
32
33 under Conventional and No-tillage Systems. Soil Sci. Soc. Am. J. 63, 1350-1358.
34

35
36 Smith, A.W., Hammond, R.B., Stinner, B.R., 1988. Influence of rye-cover crop management on
37
38 soybean foliage arthropods. Environ. Entomol. 17, 109-114.
39

40
41 Swift, M.J., Bignell, D., 2001. Standard Methods for Assessment of Soil Biodiversity and Land
42
43 use Practice. ASB Lecture Note 6B, Bogor, Indonesia.
44

45
46 Ter Braak, C.J.F., 1988. CANOCO- A FORTRAN Program for Canonical Community
47
48 Ordination by Partial Detrended Canonical Correspondence Analysis and
49
50 Redundancy Analysis (version 2.1). Report LWA-88-02. Agricultural Mathematics
51
52 Group, Wageningen.
53

54
55 Ter Braak, C.J.F., Smilauer, P., 1998. CANOCO Reference manual and User's Guide to Canoco
56
57
58
59
60
61
62
63
64
65

1
2
3
4 for Windows: Software for Canonical Community Ordination (Version 4).
5
6 Microcomputer Power, Ithaca, NY, USA.
7

8
9 Tian, G., Kang, B.T., Brussaard, L., Swift, M.J., 1997. Soil fauna mediated decomposition of
10
11 plant residues under constrained environmental and residue quality and conditions. In:
12
13 Cadisch, G., Giller, K.E. (eds.), Driven by Nature: Plant Litter Quality and
14
15 Decomposition. CAB International Wallingford, UK, pp. 125-134.
16
17

18
19 Vanlauwe, B., Palm, C.A., Murwira, H.K., Merckx, R., 2002. Organic resource management in
20
21 Sub-Saharan Africa: validation of a residue quality-driven decision support system.
22
23 *Agronomie* 22, 839-846.
24
25

26
27 Wu, T., Schoenau, J.J., Li, Qian, P., Mahli, S.S., Shi, Y., 2003. Effect of tillage and rotation on
28
29 organic carbon forms of chernozemic soils. *J. Plant Nutr. Soil Sci.* 166, 328-335.
30

31
32 Zehnder, G.W., Linduska, J.J. 1987. Influence of conservation tillage practices on populations of
33
34 Colorado potato beetle (Coleoptera: Chrysomelidae) in rotated and non-rotated tomato
35
36 fields. *Entomology* 16, 135-139.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Treatment description of selected sites and their soil organic C content (g kg^{-1} soil) at 0-15 and 15-30 cm depths.

Trial	Embu	Kabete	Impala	Nyabeda	Chitala	†Ibadan	Tamale	Sadore	Farakoba	Saria I	Saria II	Saria III		
Year established	1993	1976	2000	2003	1995	1996	1996	1986	1993	1960	1980	1990		
Treatments	Fallow	Fallow	Fallow type	Woodland since 1993	Bushland since 1976	Shrubland since 2000	Shrubland since 2003	Grass fallow since 1995	Bushland since 1986	Grass fallow since 1996	Shrubland since 1986	Grass fallow since 1993	Grass fallow since 1959	Grass fallow since 1959
													(Common for Sarias)	
	Soil C (g Kg^{-1} soil)	40.0 ^a , 28.4 ^b	23.7 ^a , 20.8 ^b	24.5 ^a , 16.9 ^b	28.2 ^a , 16.6 ^b	13.6 ^a , 9.4 ^b	12.4 ^a , 7.5 ^b	9.0 ^a , 4.8 ^b	1.5 ^a , 1.2 ^b	3.6 ^a , 2.9 ^b	3.3 ^a , 3.1 ^b	See Saria I	See Saria I	
High-C Rotation	Cont. maize	Maize-bean	Maize- <i>Tephrosia candida</i> relay	Maize-soybean	Maize-pigeonpea	Maize-cowpea	Cont. maize	Cont. millet	Cont. sorghum	Sorghum-cowpea	Cont. sorghum	Cont. sorghum		
Tillage	Hand hoeing	Hand hoeing	Hand hoeing	No till	Tractor till	Light surface hoeing	No till	Hand hoeing	Tractor till	Tractor till	Tractor till	Oxen plough		
Organic	<i>Leucaena leucocephala</i> (5 t ha ⁻¹)	10 t ha ⁻¹ manure	<i>T. candida</i> residues (5 t ha ⁻¹)	Maize stover (2 t ha ⁻¹)	Crop residues: stem+ leaves (5 t ha ⁻¹)	<i>Senna siamea</i> (5 t ha ⁻¹)	None	Crop residues	Compost (5 t ha ⁻¹)	Manure (5 t ha ⁻¹ every 2 yrs)	Manure (10 t ha ⁻¹)	Manure (10 t ha ⁻¹)		
Inorganic	None	CAN (120 kg N ha ⁻¹) and TSP (52.8 kg P ha ⁻¹)	None	NPK fertilizer (60:60:60)	(NH ₄) ₂ SO ₄ fertilizer (96 kg N ha ⁻¹ yr ⁻¹)	NPK fertilizer (60:30:30) (60:30:30 ha ⁻¹ yr ⁻¹)	None	13 kg P ha ⁻¹ ; no N	PK fertilizer (25:14)	NPK (100 kg ha ⁻¹) and Urea (50 kg ha ⁻¹ every 2 yrs)	23 kg N ha ⁻¹	NPK (100 kg ha ⁻¹ yr ⁻¹) and Urea (50 kg ha ⁻¹ yr ⁻¹)		
	Soil C (g Kg^{-1} soil)	27.5 ^a , 24.9 ^b	21.9 ^a , 17.2 ^b	27.9 ^a , 21.8 ^b	22.8 ^a , 17.8 ^b	8.8 ^a , 9.9 ^b	10.6 ^a , 6.7 ^b	5.7 ^a , 3.7 ^b	1.4a, 1.2 ^b	3.6 ^a , 2.4 ^b	2.6 ^a , 3.7 ^b	3.5 ^a , 3.3 ^b	2.4 ^a , 3.3 ^b	

Table 1 (Continued) Treatment description of selected sites and their soil organic C content (g kg^{-1} soil) at 0-15 and 15-30 cm depths.

Treatments	Low-C	Rotation	Cont. maize	Maize-bean	Cont.	Maize-soybean	Cont. maize	Maize-cowpea	Cont. maize	Millet-cowpea	Cont. sorghum	Cont. sorghum	Cont. sorghum	Cont. sorghum
	Tillage		Hand hoeing	Hand hoeing	No till	Hand hoeing	Tractor till	Light surface hoeing	Bullock plough	Animal traction	Tractor till	Tractor till	Tractor till	Hand hoeing
	Organic		None	None	None	None	None	None	None	Crop residues	None	Manure (5 t ha^{-1} every 2 yrs)	None	no organic inputs, 4=
	Inorganic		None	None	None	NPK fertilizer (60:60:60)	None	None	None	30 kg N ha^{-1} ; 13 kg P ha^{-1}	None	NPK (100 kg ha^{-1}) and Urea ($50 \text{ kg ha}^{-1} \text{ yr}^{-1}$)	None	NPK (100 kg ha^{-1}) and Urea ($50 \text{ kg ha}^{-1} \text{ yr}^{-1}$)
	Soil C (g Kg^{-1} soil)		$24.6^a, 23.9^b$	$17.7^a, 16.2^b$	$22.0^a, 21.2^b$	$19.9^a, 18.4^b$	$8.2^a, 7.1^b$	$5.7^a, 3.9^b$	$3.1^a, 2.6^b$	$1.0^a, 0.9^b$	$2.7^a, 2.3^b$	$3.3^a, 3.2^b$	$1.8^a, 2.5^b$	$2.2^a, 3.0^b$

Note: Soil C data source: Eastern Africa (from monolith soils, Western Africa (Zida et al., unpublished), †Ibadan (Vanlauwe et al., 2002). Lower case superscripts (a, b) denotes 0-15 and 15-30 cm depths respectively.

Table 2 Macrofauna (earthworm and termite) mean taxonomic richness and abundance across fallow and arable systems of Eastern and Western Africa

Site	Earthworm						Termite											
	Fallow			High-C			Low-C			Fallow			High-C			Low-C		
	Taxonomic richness (S)	Abundance (no m ⁻²)		Taxonomic richness (S)	Abundance (no m ⁻²)		Taxonomic richness (S)	Abundance (no m ⁻²)		Taxonomic richness (S)	Abundance (no m ⁻²)		Taxonomic richness (S)	Abundance (no m ⁻²)				
Embu	2.67 ^a	0.33 ^b	0.67 ^b	1264 ^a	32 ^b	27 ^b	1.33 ^a	1.33 ^a	1.00 ^a	139 ^a	528 ^a	32 ^a						
Kabete	3.00 ^a	2.00 ^{ab}	1.00 ^b	363 ^a	395 ^a	272 ^a	1.00 ^a	0.33 ^a	0.67 ^a	203 ^b	1029 ^a	384 ^{ab}						
Impala	3.33 ^a	0.67 ^b	0.67 ^b	165 ^a	32 ^b	16 ^b	2.00 ^a	1.68 ^a	1.33 ^a	987 ^a	43 ^b	1104 ^a						
Nyabeda	0.67 ^a	0.33 ^a	0.33 ^a	117 ^a	11 ^{ab}	0 ^b	3.00 ^a	2.68 ^a	2.67 ^a	475 ^a	208 ^{ab}	0 ^b						
Chitala	1.00 ^a	1.00 ^a	1.00 ^a	53 ^a	5 ^a	5 ^a	2.33 ^a	2.00 ^a	2.00 ^a	752 ^{ab}	117 ^b	1744 ^a						
Ibadan	1.67 ^b	4.67 ^a	1.67 ^b	96 ^b	933 ^a	117 ^b	2.00 ^a	1.33 ^a	0.00 ^b	320 ^b	1483 ^a	0 ^b						
Tamale	2.00 ^a	1.00 ^{ab}	0.00 ^b	37 ^a	32 ^a	16 ^a	2.00 ^a	1.68 ^a	1.00 ^a	149 ^b	1179 ^a	32 ^b						
Sadore	0.00 ^a	0.00 ^a	0.00 ^a	0 ^a	0 ^a	0 ^a	2.00 ^a	1.00 ^a	2.00 ^a	256 ^a	181 ^a	0 ^a						
Farakoba	2.33 ^a	0.67 ^b	0.33 ^b	453 ^a	59 ^b	11 ^b	0.68 ^b	3.00 ^a	1.00 ^b	59 ^b	693 ^a	160 ^{ab}						
Saria I	1.33 ^a	0.67 ^a	0.00 ^a	69 ^a	27 ^a	21 ^a	1.68 ^a	1.00 ^a	1.00 ^b	112 ^a	0 ^a	0 ^a						
Saria II	1.33 ^a	0.33 ^{ab}	0.00 ^b	69 ^a	11 ^a	0 ^a	1.68 ^a	1.33 ^a	0.68 ^a	112 ^a	256 ^a	0 ^a						
Saria III	1.33 ^{ab}	0.67 ^b	2.00 ^a	69 ^a	80 ^a	59 ^a	1.68 ^a	1.00 ^a	1.33 ^a	112 ^a	11 ^a	112 ^a						
Mean	1.72^a	1.02^b	0.64^c	230^a	135^b	45^b	1.78^a	1.53^{ab}	1.22^b	306^{ab}	477^a	297^b						
SV	df	F-ratio	P-value	df	F-ratio	P-value	df	F-ratio	P-value	df	F-ratio	P-value						
Site	11	7.40	<0.001	11	10.07	<0.001	11	4.83	<0.001	11	2.74	0.005						
Treatment	2	16.02	<0.001	2	20.37	<0.001	2	3.69	0.030	2	2.54	0.086						
Site*treat	22	3.44	<0.001	22	4.73	<0.001	22	1.41	0.142	22	2.35	0.004						
Residual	70			70			70			70								

Means of S or abundance followed by same lower case letters within rows across treatments are not statistically significant at p<0.05.

Figure 1




Figure 2




Figure Captions

Figure 1 RDA biplot showing correlation between earthworm species abundance and selected environmental variables. Earthworm taxa are represented by triangles, while environmental parameters are represented by pointed arrows. Sites are represented by open symbols.

Earthworm species abbreviations: *Nematogenia lacuum* (**Nlac**), *Gordiodrilus robusta* (**Grob**), *Gordiodrilus wemanus* (**Gwem**), *Gordiodrilus marcusii* (**Gmar**), *Polytoreutus annulatus* (**Panul**), *Hyperiodrilus africanus* (**Hafr**), *Hyperiodrilus* sp. nov (**Hspnv**), *Eudrilus buettneri* (**Ebuet**), *Ephyrodrilus afroccidentalis* (**Eafro**), *Eminoscolex violaceus* (**Eviol**), *Stuhlmannia* sp. nov (**Sspnv**), *Lavellea* sp. nov (**Lspnv**), *Millsonia inermis* (**Miner**), *Millsonia guttata* (**Mgutt**), *Dichogaster (Dt.) saliens* (**Dsal**), *Dichogaster (Dt.) affinis* (**Daffn**), *Dichogaster (Dt.) bolau* (**Dbol**), *Dichogaster (Dt.) modiglianii* (**Dmod**), *Dichogaster (Dt.)* sp. nov 1 (**Dspv1**), *Dichogaster (Dt.)* sp. nov 2 (**Dspv2**). **Environmental abbreviations:** Altitude (**Alt**), Average temperature (**AverT**), Carbon (**C**), Latitude (**Lat**), Longitude (**Long**), Precipitation (**PPT**).

Figure 2 RDA biplot showing correlation between termite species abundance and selected environmental variables. Termite taxa are represented by triangles, while environmental parameters are represented by pointed arrows. Sites are represented by open symbols.

Species abbreviations: *Amitermes stephensoni* (**Astep**), *Ancistrotermes cavithorax* (**Acav**), *Captotermes intermedius* (**Cint**), *Macrotermes nr. vitriolatus* (**Mvitr**), *Macrotermes subhyalinus* (**Msub**), *Macrotermes herus* (**Mherus**), *Macrotermes* sp. (**Mterm**), *Microcerotermes parvulus* (**Mparv**), *Microtermes* sp. (**Microt**), *Nasutitermes* sp. (**Nasut**), *Trinevitermes* sp. (**Trinv**), *Odontotermes magdalenae* (**Odom**), *Odontotermes* sp. (**Odont**), *Psammotermes hybostoma* (**Phyb**), *Pseudacanthotermes spiniger* (**Pspn**), *Pseudacanthotermes* sp. (**Pseud**), *Schedorhinotermes lamanianus* (**Slam**), *Termes baculi* (**Tbac**), *Tuberculitermes* sp. (**Tubec**). **Environmental abbreviations:** Average temperature (**AverT**), Latitude (**Lat**), Longitude (**Long**), Precipitation (**PPT**).

Appendix 1 Earthworm taxonomic richness and functional diversity based on monolith method.

Taxonomic group	Functional group ^a	Site											
		Eastern Africa					Western Africa						
		Embu	Kabete	Impala	Nyabeda	Chitala	Ibadan	Tamale	Sadore	Farakoba	Saria I	Saria II	Saria III
Ocnerodrilidae													
<i>Nematogenia lacuum</i>	Endogeic	+	+	-	+	-	-	-	-	-	-	-	-
<i>Gordiodrilus robustus</i>	Endogeic	-	-	-	-	-	+	-	-	-	-	-	-
<i>Gordiodrilus wemanus</i>	Endogeic	-	+	+	+	-	-	-	-	-	-	-	-
<i>Gordiodrilus marcusii</i>	Endogeic	-	-	-	-	-	+	-	-	-	-	-	-
Acanthodrilidae													
<i>Millsonia inermis</i>	Endogeic	-	-	-	-	-	+	-	-	-	+	+	-
<i>Millsonia guttata</i>	Endogeic	-	-	-	-	-	-	+	-	-	-	-	-
<i>Dichogaster (Dt.) saliens</i>	Epigeic	+	-	+	-	-	-	-	-	-	-	-	-
<i>Dichogaster (Dt.) affinis</i>	Epigeic	-	+	+	+	-	+	-	-	-	+	-	+
<i>Dichogaster (Dt.) bolauii</i>	Epigeic	-	+	+	-	-	-	-	-	-	-	-	-
<i>Dichogaster (Dt.) modiglianii</i>	Epigeic	-	-	+	-	-	-	-	-	-	-	-	-
<i>Dichogaster (Dt.)</i> sp. nov 1	Epigeic	-	-	-	-	-	+	-	-	-	-	-	+
<i>Dichogaster (Dt.)</i> sp. nov 2	Epigeic	-	-	-	-	+	-	-	-	-	-	-	-
Eudrilidae													
<i>Polytoreutus annulatus</i>	Epigeic	+	+	-	-	-	-	-	-	-	-	-	-
<i>Hyperiodrilus africanus</i>	Epigeic	-	-	-	-	-	+	+	-	+	-	-	-
<i>Hyperiodrilus</i> sp. nov	Epigeic	-	-	-	-	-	+	+	-	+	-	-	-
<i>Eudrilus buettneri</i>	Epigeic	-	-	-	-	-	+	-	-	+	-	-	-
<i>Ephyriodrilus afroccidentalis</i>	Epigeic	-	-	-	-	-	+	-	-	-	-	-	-
<i>Eminoscolex violaceus</i>	Epigeic	-	-	+	+	-	-	-	-	-	-	-	-
<i>Stuhlamannia</i> sp. nov	Epigeic	-	+	-	-	-	+	-	-	-	-	-	-
<i>Lavellea</i> sp. nov	Epigeic	-	-	-	-	-	+	-	-	+	-	-	-
Taxonomic richness (S)		3	6	6	4	1	11	3	0	4	2	1	2

^aBased on classification by Swift and Bignell (2001). +/- denotes presence or absence, respectively.

Appendix 2 Termite taxonomic richness and functional groups based on monolith and transect methods.

Taxonomic group	Functional group ^a	Food type ^b	Possible pest ^c	Site											
				Eastern Africa					Western Africa						
				Embu	Kabete	Impala	Nyabeda	Chitala	Ibadan	Tamale	Sadore	Farakoba	Saria I	Saria II	Saria III
Rhinotermitidae-Rhinotermitinae															
<i>Captotermes intermedius</i>	I	WLG	Yes	-	-	-	-	-	-	+	-	-	-	-	-
<i>Schedorhinotermes lamanianus</i>	I	WLG	Yes	-	-	-	-	+	-	-	-	-	-	-	-
Rhinotermitidae-Psammotermitinae															
<i>Psammotermes hybostoma</i>	I	WLG	Yes	-	-	-	-	-	-	-	+	-	-	-	-
Termitidae-Nasutitermitinae															
<i>Nasutitermes</i> sp.	II	WLG	Yes	-	-	-	-	+	-	-	-	-	-	-	-
<i>Trinevitermes</i> sp.	II	LG	Yes	-	-	-	-	-	-	+	-	+	+	+	+
Termitidae-Macrotermitinae															
<i>Ancistrotermes cavithorax</i>	II	WLG	Yes	-	-	-	-	-	+	+	-	+	-	-	-
<i>Macrotermes nr. vitriolatus</i>	II	FWLG	Yes	-	-	-	-	+	-	-	-	-	-	-	-
<i>Macrotermes subhyalinus</i>	II	FWLG	Yes	-	-	-	-	-	-	-	-	-	+	+	+
<i>Macrotermes herus</i>	II	FWLG	Yes	-	-	-	+	-	-	-	-	-	-	-	-
<i>Microtermes pusillas</i>	II	FWLG	Yes	-	-	-	+	-	-	-	-	-	-	-	-
<i>Macrotermes</i> spp.	II	FWLG	Yes	-	-	-	-	+	-	-	-	-	-	-	-
<i>Microtermes</i> spp.	II	FWLG	Yes	+	+	+	+	+	+	+	+	+	+	+	+
<i>Odontotermes magdalenae</i>	II	FWLG	Yes	-	-	-	-	-	-	-	-	-	+	+	+
<i>Odontotermes</i> spp.	II	FWLG	Yes	+	+	-	-	+	-	+	+	+	-	-	-
<i>Pseudacanthotermes spiniger</i>	II	FLG	No	-	-	+	+	-	-	-	-	-	-	-	-
<i>Pseudacanthotermes</i> spp.	II	FLSD	Yes	+	+	-	-	-	+	-	-	+	-	-	-
Termitidae-Termitinae															
<i>Amitermes-Amitermes stephensoni</i>	II	WLSL	Yes	-	-	-	-	-	-	-	-	-	+	+	+
<i>Microcerotermes parvulus</i>	II	WLG	Yes	-	-	-	-	+	-	-	+	+	+	+	+
<i>Termes-Termes baculi</i>	III	WS	No	-	-	-	-	-	+	-	-	-	-	-	-
<i>Cubitermes-Tubeculitermes</i> sp.	IV	S	No	-	-	+	+	-	-	-	-	-	-	-	-
Taxonomic richness (S)				3	3	3	5	7	4	5	4	6	6	6	6


Footnote for table 4

(W=wood; L=leaf litter; S=soil; D=Dung/manure; F=fungus grower; G=dead/dry grass. ^abased on classification by Bignell and Eggleton (2000) and Eggleton et al. (2002). ^bbased on field notes/observation. ^cbased on observations by Kooyman and Onck (1987). +/- denotes presence or absence, respectively.

AGEE5072R2

...

From: "Agriculture, Ecosystems & Environment" <agee@elsevier.com>
...
View Contact
To: fredrick.ayuke@yahoo.com

 Ref.: Ms. No. AGE5072R2

Agricultural management affects earthworm and termite diversity across humid to semi-arid tropical zones

Agriculture, Ecosystems and Environment

Dear Mr Ayuke,

The Editor has now evaluated the [revised version](#) of the manuscript: Agricultural management affects earthworm and termite diversity across humid to semi-arid tropical zones.

However, before it can be reconsidered for publication, the comments below have to be taken into account. Please submit the revision within 60 days of receiving this letter and explain in a [covering letter](#) how and where the comments have been incorporated in the text.

To submit a revision, go to <http://ees.elsevier.com/agee/> and log in as an Author.

You will see a menu item called "Submissions Needing Revision". You will find your submission record there. Please ensure that you upload the source files. This is especially important in case of figure files, please upload these one by one in EES. For example: fig. 1a, 1b and 1c as one file and figures 2 and 3 as separate files. Uploading a PDF file at this stage will create delays should your manuscript be finally accepted for publication.

I will look forward to receiving the revised version of your manuscript.

Olaf Meesters
Journal Manager

Important note: If a reviewer has provided a review or other materials as attachments, those items will not be in this letter. Please ensure therefore that you log on to the journal site and check if any attachments have been provided.

Editor's comments:

EDITOR

This new version still requires some more attention with a view to:

- simplify Highlights to :

? [Earthworms](#) and termites were favored by warm and dry conditions + **Done**.

- ? Cropping negatively affected earthworms more than termites + **Done**.
- ? Management leading to low soil C supported fewer earthworm and termite taxa + **Done**.
- stick to common units e.g. "t" instead of Mg in Table 1 + **Done**
- ; "min" l. 60, p. 4 + **Done** (see l. 4, p. 5).
- modify l. 24, p. 5 to: "Earthworms and termites were..." + **Done** (l. 26, p. 5); l. 31, p. 5 to "Adult earthworms and termites were..." + **Done** (l. 34, p. 5) " omitting l. 38-51 ; + **Done**.
- erase whole () l. 56, p. 5; + **Done**.
- place also Table 2 under Supplementary Material; With reference to table 2. + **The authors still respectfully disagree with this. This table captures hypotheses 2 and 3 and is the core of the study as reflected in the title. Moreover, by removing the table, the results in sections 3.1, 3.2 and discussion (section 4.2) would become irrelevant. Therefore, the authors feel the table should remain in the manuscript.**
- change l. 37, p. 7 to : "However, site-treatment interactions were also significant and there were significant...";+ **Done** (l. 22-24, p. 7).
- stick to past tense as of l. 48, 50, p. 8 ; etc. + **Effectuated** (l. 32-42, p. 8; l. 59, p. 8 to l. 17, p. 9).
- check caption to both Figures correcting "sp." throughout and "vitriolatus", referring to Figure 1 for environmental abbreviations. + **Effectuated**.